

$\Upsilon(2S)$ $I^G(J^{PC}) = 0^-(1^{--})$ **$\Upsilon(2S)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1023.26 ± 0.31 OUR AVERAGE			
10023.5 ± 0.5	¹ ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
10023.1 ± 0.4	BARBER 84	REDE	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10023.6 ± 0.5	2,3 BARU	86B REDE	$e^+ e^- \rightarrow$ hadrons
¹ Reanalysis of BARU 86B using new electron mass (COHEN 87).			
² Reanalysis of ARTAMONOV 84.			
³ Superseded by ARTAMONOV 00.			

 $m\Upsilon(3S) - m\Upsilon(2S)$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$331.50 \pm 0.02 \pm 0.13$	LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$

 $\Upsilon(2S)$ WIDTH

VALUE (keV)	DOCUMENT ID
31.98 ± 2.63 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"

 $\Upsilon(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \Upsilon(1S)\pi^+\pi^-$	(17.85 ± 0.26) %	
$\Gamma_2 \Upsilon(1S)\pi^0\pi^0$	(8.6 ± 0.4) %	
$\Gamma_3 \tau^+\tau^-$	(2.00 ± 0.21) %	
$\Gamma_4 \mu^+\mu^-$	(1.93 ± 0.17) %	S=2.2
$\Gamma_5 e^+e^-$	(1.91 ± 0.16) %	
$\Gamma_6 \Upsilon(1S)\pi^0$	< 4 $\times 10^{-5}$	CL=90%
$\Gamma_7 \Upsilon(1S)\eta$	(2.9 ± 0.4) $\times 10^{-4}$	S=2.0
$\Gamma_8 J/\psi(1S)$ anything	< 6 $\times 10^{-3}$	CL=90%
$\Gamma_9 J/\psi(1S)\eta_c$	< 5.4 $\times 10^{-6}$	CL=90%
$\Gamma_{10} J/\psi(1S)\chi_{c0}$	< 3.4 $\times 10^{-6}$	CL=90%
$\Gamma_{11} J/\psi(1S)\chi_{c1}$	< 1.2 $\times 10^{-6}$	CL=90%
$\Gamma_{12} J/\psi(1S)\chi_{c2}$	< 2.0 $\times 10^{-6}$	CL=90%
$\Gamma_{13} J/\psi(1S)\eta_c(2S)$	< 2.5 $\times 10^{-6}$	CL=90%
$\Gamma_{14} J/\psi(1S)X(3940)$	< 2.0 $\times 10^{-6}$	CL=90%
$\Gamma_{15} J/\psi(1S)X(4160)$	< 2.0 $\times 10^{-6}$	CL=90%
$\Gamma_{16} \chi_{c1}$ anything	(2.2 ± 0.5) $\times 10^{-4}$	

Γ_{17}	$\chi_{c1}(1P)^0 X_{tetra}$	$< 3.67 \times 10^{-5}$	CL=90%
Γ_{18}	χ_{c2} anything	$(2.3 \pm 0.8) \times 10^{-4}$	
Γ_{19}	$\psi(2S)\eta_c$	$< 5.1 \times 10^{-6}$	CL=90%
Γ_{20}	$\psi(2S)\chi_{c0}$	$< 4.7 \times 10^{-6}$	CL=90%
Γ_{21}	$\psi(2S)\chi_{c1}$	$< 2.5 \times 10^{-6}$	CL=90%
Γ_{22}	$\psi(2S)\chi_{c2}$	$< 1.9 \times 10^{-6}$	CL=90%
Γ_{23}	$\psi(2S)\eta_c(2S)$	$< 3.3 \times 10^{-6}$	CL=90%
Γ_{24}	$\psi(2S)X(3940)$	$< 3.9 \times 10^{-6}$	CL=90%
Γ_{25}	$\psi(2S)X(4160)$	$< 3.9 \times 10^{-6}$	CL=90%
Γ_{26}	$Z_c(3900)^+ Z_c(3900)^-$	$< 1.0 \times 10^{-6}$	CL=90%
Γ_{27}	$Z_c(4200)^+ Z_c(4200)^-$	$< 1.67 \times 10^{-5}$	CL=90%
Γ_{28}	$Z_c(3900)^{\pm} Z_c(4200)^{\mp}$	$< 7.3 \times 10^{-6}$	CL=90%
Γ_{29}	$X(4050)^+ X(4050)^-$	$< 1.35 \times 10^{-5}$	CL=90%
Γ_{30}	$X(4250)^+ X(4250)^-$	$< 2.67 \times 10^{-5}$	CL=90%
Γ_{31}	$X(4050)^{\pm} X(4250)^{\mp}$	$< 2.72 \times 10^{-5}$	CL=90%
Γ_{32}	$Z_c(4430)^+ Z_c(4430)^-$	$< 2.03 \times 10^{-5}$	CL=90%
Γ_{33}	$X(4055)^{\pm} X(4055)^{\mp}$	$< 1.11 \times 10^{-5}$	CL=90%
Γ_{34}	$X(4055)^{\pm} Z_c(4430)^{\mp}$	$< 2.11 \times 10^{-5}$	CL=90%
Γ_{35}	$\overline{^2H}$ anything	$(2.78^{+0.30}_{-0.26}) \times 10^{-5}$	S=1.2
Γ_{36}	hadrons	$(94 \pm 11) \%$	
Γ_{37}	ggg	$(58.8 \pm 1.2) \%$	
Γ_{38}	γgg	$(1.87 \pm 0.28) \%$	
Γ_{39}	$\phi K^+ K^-$	$(1.6 \pm 0.4) \times 10^{-6}$	
Γ_{40}	$\omega \pi^+ \pi^-$	$< 2.58 \times 10^{-6}$	CL=90%
Γ_{41}	$K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(2.3 \pm 0.7) \times 10^{-6}$	
Γ_{42}	$\phi f_2'(1525)$	$< 1.33 \times 10^{-6}$	CL=90%
Γ_{43}	$\omega f_2(1270)$	$< 5.7 \times 10^{-7}$	CL=90%
Γ_{44}	$\rho(770) a_2(1320)$	$< 8.8 \times 10^{-7}$	CL=90%
Γ_{45}	$K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$	$(1.5 \pm 0.6) \times 10^{-6}$	
Γ_{46}	$K_1(1270)^{\pm} K^{\mp}$	$< 3.22 \times 10^{-6}$	CL=90%
Γ_{47}	$K_1(1400)^{\pm} K^{\mp}$	$< 8.3 \times 10^{-7}$	CL=90%
Γ_{48}	$b_1(1235)^{\pm} \pi^{\mp}$	$< 4.0 \times 10^{-7}$	CL=90%
Γ_{49}	$\rho \pi$	$< 1.16 \times 10^{-6}$	CL=90%
Γ_{50}	$\pi^+ \pi^- \pi^0$	$< 8.0 \times 10^{-7}$	CL=90%
Γ_{51}	$\omega \pi^0$	$< 1.63 \times 10^{-6}$	CL=90%
Γ_{52}	$\pi^+ \pi^- \pi^0 \pi^0$	$(1.30 \pm 0.28) \times 10^{-5}$	
Γ_{53}	$K_S^0 K^+ \pi^- + \text{c.c.}$	$(1.14 \pm 0.33) \times 10^{-6}$	
Γ_{54}	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$< 4.22 \times 10^{-6}$	CL=90%
Γ_{55}	$K^*(892)^- K^+ + \text{c.c.}$	$< 1.45 \times 10^{-6}$	CL=90%
Γ_{56}	$f_1(1285)$ anything	$(2.2 \pm 1.6) \times 10^{-3}$	
Γ_{57}	$f_1(1285) X_{tetra}$	$< 6.47 \times 10^{-5}$	CL=90%
Γ_{58}	Sum of 100 exclusive modes	$(2.90 \pm 0.30) \times 10^{-3}$	

Radiative decays

Γ_{59}	$\gamma\chi_{b1}(1P)$	(6.9 \pm 0.4) %	
Γ_{60}	$\gamma\chi_{b2}(1P)$	(7.15 \pm 0.35) %	
Γ_{61}	$\gamma\chi_{b0}(1P)$	(3.8 \pm 0.4) %	
Γ_{62}	$\gamma f_0(1710)$	< 5.9 $\times 10^{-4}$	CL=90%
Γ_{63}	$\gamma f'_2(1525)$	< 5.3 $\times 10^{-4}$	CL=90%
Γ_{64}	$\gamma f_2(1270)$	< 2.41 $\times 10^{-4}$	CL=90%
Γ_{65}	$\gamma f_J(2220)$		
Γ_{66}	$\gamma\eta_c(1S)$	< 2.7 $\times 10^{-5}$	CL=90%
Γ_{67}	$\gamma\chi_{c0}$	< 1.0 $\times 10^{-4}$	CL=90%
Γ_{68}	$\gamma\chi_{c1}$	< 3.6 $\times 10^{-6}$	CL=90%
Γ_{69}	$\gamma\chi_{c2}$	< 1.5 $\times 10^{-5}$	CL=90%
Γ_{70}	$\gamma\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi$	< 8 $\times 10^{-7}$	CL=90%
Γ_{71}	$\gamma\chi_{c1}(3872) \rightarrow \pi^+\pi^-\pi^0 J/\psi$	< 2.4 $\times 10^{-6}$	CL=90%
Γ_{72}	$\gamma X(3915) \rightarrow \omega J/\psi$	< 2.8 $\times 10^{-6}$	CL=90%
Γ_{73}	$\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi$	< 1.2 $\times 10^{-6}$	CL=90%
Γ_{74}	$\gamma X(4350) \rightarrow \phi J/\psi$	< 1.3 $\times 10^{-6}$	CL=90%
Γ_{75}	$\gamma\eta_b(1S)$	(5.5 \pm 1.1) $\times 10^{-4}$	S=1.2
Γ_{76}	$\gamma\eta_b(1S) \rightarrow \gamma$ Sum of 26 exclusive modes	< 3.7 $\times 10^{-6}$	CL=90%
Γ_{77}	$\gamma X_{b\bar{b}} \rightarrow \gamma$ Sum of 26 exclusive modes	< 4.9 $\times 10^{-6}$	CL=90%
Γ_{78}	$\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 1.95 $\times 10^{-4}$	CL=95%
Γ_{79}	$\gamma A^0 \rightarrow \gamma$ hadrons	< 8 $\times 10^{-5}$	CL=90%
Γ_{80}	$\gamma a_1^0 \rightarrow \gamma\mu^+\mu^-$	< 8.3 $\times 10^{-6}$	CL=90%

Lepton Family number (*LF*) violating modes

Γ_{81}	$e^\pm\tau^\mp$	<i>LF</i>	< 3.2 $\times 10^{-6}$	CL=90%
Γ_{82}	$\mu^\pm\tau^\mp$	<i>LF</i>	< 3.3 $\times 10^{-6}$	CL=90%

[a] 1.5 GeV $< m_X <$ 5.0 GeV

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 13 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 11.8$ for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$x_7 \quad \begin{array}{|c} \hline 2 \\ \hline \end{array} \\ x_1$$

$\Upsilon(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_4\Gamma_5/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
6.5±1.5±1.0	KOBEL	92	$e^+e^- \rightarrow \mu^+\mu^-$

$\Gamma(\Upsilon(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
105.4±1.0±4.2	11.8K	¹ AUBERT	08BP BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$

¹ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{36}\Gamma_5/\Gamma$		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.577±0.009 OUR AVERAGE			
0.581±0.004±0.009	¹ ROSNER 06	CLEO	$10.0 e^+e^- \rightarrow \text{hadrons}$
0.552±0.031±0.017	¹ BARU 96	MD1	$e^+e^- \rightarrow \text{hadrons}$
0.54 ± 0.04 ± 0.02	¹ JAKUBOWSKI 88	CBAL	$e^+e^- \rightarrow \text{hadrons}$
0.58 ± 0.03 ± 0.04	² GILES 84B	CLEO	$e^+e^- \rightarrow \text{hadrons}$
0.60 ± 0.12 ± 0.07	² ALBRECHT 82	DASP	$e^+e^- \rightarrow \text{hadrons}$
0.54 ± 0.07 +0.09 -0.05	² NICZYPORUK 81C	LENA	$e^+e^- \rightarrow \text{hadrons}$
0.41 ± 0.18	² BOCK 80	CNTR	$e^+e^- \rightarrow \text{hadrons}$

¹ Radiative corrections evaluated following KURAEV 85.

² Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

 $\Upsilon(2S) \text{ PARTIAL WIDTHS}$

$\Gamma(e^+e^-)$	Γ_5
VALUE (keV)	DOCUMENT ID
0.612±0.011 OUR EVALUATION	

 $\Upsilon(2S) \text{ BRANCHING RATIOS}$

$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ
Abbreviation MM in the COMMENT field below stands for missing mass.	
17.85±0.26 OUR FIT	
17.92±0.26 OUR AVERAGE	
16.8 ± 1.1 ± 1.3	906k ¹ LEES 11c BABR $e^+e^- \rightarrow \pi^+\pi^-X$
17.80±0.05±0.37	170k ² LEES 11L BABR $\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
18.02±0.02±0.61	851k ³ BHARI 09 CLEO $e^+e^- \rightarrow \pi^+\pi^- \text{MM}$
17.22±0.17±0.75	11.8K ⁴ AUBERT 08BP BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
19.2 ± 0.2 ± 1.0	52.6k ⁵ ALEXANDER 98 CLE2 $\pi^+\pi^-\ell^+\ell^-, \pi^+\pi^- \text{MM}$
18.1 ± 0.5 ± 1.0	11.6k ALBRECHT 87 ARG $e^+e^- \rightarrow \pi^+\pi^- \text{MM}$
16.9 ± 4.0	GELPHMAN 85 CBAL $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
19.1 ± 1.2 ± 0.6	BESSON 84 CLEO $\pi^+\pi^- \text{MM}$

18.9 \pm 2.6	FONSECA 84	CUSB $e^+ e^- \rightarrow \ell^+ \ell^- \pi^+ \pi^-$
21 \pm 7	7	NICZYPORUK 81B LENA $e^+ e^- \rightarrow \ell^+ \ell^- \pi^+ \pi^-$

¹ LEES 11C reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything})] = (1.78 \pm 0.02 \pm 0.11) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything}) = (10.6 \pm 0.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

³ A weighted average of the inclusive and exclusive results.

⁴ Using $B(\Upsilon(2S) \rightarrow e^+ e^-) = (1.91 \pm 0.16)\%$, $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$ and, $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011 \text{ keV}$.

⁵ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.07)\%$.

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
8.6 \pm 0.4 OUR AVERAGE				
8.43 \pm 0.16 \pm 0.42	38k	¹ BHARI 09	CLEO	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
9.2 \pm 0.6 \pm 0.8	275	² ALEXANDER 98	CLE2	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
9.5 \pm 1.9 \pm 1.9	25	ALBRECHT 87	ARG	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
8.0 \pm 1.5		GELPHMAN 85	CBAL	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
10.3 \pm 2.3		FONSECA 84	CUSB	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$

¹ Authors assume $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$.

² Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.07)\%$.

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_2/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

0.462 \pm 0.037	¹ BHARI 09	CLEO	$e^+ e^- \rightarrow \Upsilon(2S)$
-------------------	-----------------------	------	------------------------------------

¹ Not independent of other values reported by BHARI 09.

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00 \pm 0.21 OUR AVERAGE				

2.00 \pm 0.12 \pm 0.18	22k	¹ BESSON 07	CLEO	$e^+ e^- \rightarrow \Upsilon(2S) \rightarrow \tau^+ \tau^-$
1.7 \pm 1.5 \pm 0.6		HAAS 84B	CLEO	$e^+ e^- \rightarrow \tau^+ \tau^-$

¹ BESSON 07 reports $[\Gamma(\Upsilon(2S) \rightarrow \tau^+ \tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \mu^+ \mu^-)] = 1.04 \pm 0.04 \pm 0.05$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

Γ_4/Γ


VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.0193±0.0017 OUR AVERAGE	Error includes scale factor of 2.2. See the ideogram below.				
0.0203±0.0003±0.0008	120k	ADAMS	05	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0122±0.0028±0.0019	1	KOBEL	92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$
0.0138±0.0025±0.0015		KAARSBERG	89	CSB2	$e^+e^- \rightarrow \mu^+\mu^-$
0.009 ± 0.006 ± 0.006	2	ALBRECHT	85	ARG	$e^+e^- \rightarrow \mu^+\mu^-$
0.018 ± 0.008 ± 0.005		HAAS	84B	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.038	90	NICZYPORUK	81C	LENA	$e^+e^- \rightarrow \mu^+\mu^-$

¹ Taking into account interference between the resonance and continuum.

² Re-evaluated using $B(\gamma(1S) \rightarrow \mu^+\mu^-) = 0.026$.

WEIGHTED AVERAGE

0.0193±0.0017 (Error scaled by 2.2)



$$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$$

$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$

Γ_3/Γ_4

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.04±0.04±0.05	22k	BESSON	07	CLEO $e^+e^- \rightarrow \tau(2S)$

$\Gamma(\tau(1S)\pi^0)/\Gamma_{\text{total}}$

Γ_6/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4	90	¹ TAMPONI	13	BELL	$e^+ e^- \rightarrow \gamma(1S)\pi^0$
< 18	90	² HE	08A	CLEO	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma\gamma$
< 110	90	ALEXANDER	98	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma\gamma$
< 800	90	LURZ	87	CBAL	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma\gamma$

¹ TAMPONI 13 reports $[\Gamma(\gamma(2S) \rightarrow \gamma(1S)\pi^0)/\Gamma_{\text{total}}] / [B(\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-)] < 2.3 \times 10^{-4}$ which we multiply by our best value $B(\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-) = 17.85 \times 10^{-2}$.

² Authors assume $B(\gamma(1S) \rightarrow e^+ e^-) + B(\gamma(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$.

$\Gamma(\gamma(1S)\pi^0)/\Gamma(\gamma(1S)\pi^+\pi^-)$

Γ_6/Γ_1

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.3	90	TAMPONI	13	$e^+ e^- \rightarrow \gamma(1S)\pi^0$

$\Gamma(\gamma(1S)\eta)/\Gamma_{\text{total}}$

Γ_7/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.9 ± 0.4 OUR FIT					Error includes scale factor of 2.0.

2.9 ± 0.4 OUR AVERAGE Error includes scale factor of 1.9. See the ideogram below.

$2.39 \pm 0.31 \pm 0.14$	112	¹ LEES	11L	BABR	$\gamma(2S) \rightarrow \ell^+ \ell^- \eta$
$2.1^{+0.7}_{-0.6} \pm 0.3$	14	² HE	08A	CLEO	$e^+ e^- \rightarrow \ell^+ \ell^- \eta$

• • • We use the following data for averages but not for fits. • • •

$3.55 \pm 0.32 \pm 0.05$	241	³ TAMPONI	13	BELL	$e^+ e^- \rightarrow \gamma(1S)\eta$
--------------------------	-----	----------------------	----	------	--------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •


< 9	90	^{1,4} AUBERT	08BP	BABR	$e^+ e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$
< 28	90	ALEXANDER	98	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \eta$
< 50	90	ALBRECHT	87	ARG	$e^+ e^- \rightarrow \pi^+\pi^-\ell^+\ell^- \text{MM}$
< 70	90	LURZ	87	CBAL	$e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, 3\pi^0)$
< 100	90	BESSON	84	CLEO	$e^+ e^- \rightarrow \pi^+\pi^-\ell^+\ell^- \text{MM}$
< 20	90	FONSECA	84	CUSB	$e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, \pi^+\pi^-\pi^0)$

¹ Using $B(\gamma(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\gamma(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

² Authors assume $B(\gamma(1S) \rightarrow e^+ e^-) + B(\gamma(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$.

³ TAMPONI 13 reports $[\Gamma(\gamma(2S) \rightarrow \gamma(1S)\eta)/\Gamma_{\text{total}}] / [B(\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-)] = (1.99 \pm 0.14 \pm 0.11) \times 10^{-3}$ which we multiply by our best value $B(\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-) = (17.85 \pm 0.26) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Using $\Gamma_{ee}(\gamma(2S)) = 0.612 \pm 0.011 \text{ keV}$.



$\Gamma(\gamma(1S)\eta)/\Gamma(\gamma(1S)\pi^+\pi^-)$

Γ_7/Γ_1

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.64±0.25 OUR FIT					Error includes scale factor of 2.0.
1.99±0.14±0.11	241	TAMPONI	13	BELL	$e^+e^- \rightarrow \gamma(1S)\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.35±0.17±0.08	1	LEES	11L	BABR	$\gamma(2S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\mu^+\mu^-$
< 5.2	90	2	AUBERT	08BP	$BABR e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

¹ Not independent of other values reported by LEES 11L.

² Not independent of other values reported by AUBERT 08BP.

$\Gamma(\gamma(1S)\pi^0)/\Gamma(\gamma(1S)\eta)$

Γ_6/Γ_7

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.13	90	TAMPONI	13	BELL $e^+e^- \rightarrow \gamma(1S)\pi^0$

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$

Γ_8/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.006	90	MASCHMANN	90	CBAL $e^+e^- \rightarrow \text{hadrons}$

$\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$

Γ_9/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 5.4×10^{-6}	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$	$CL\%$
$<3.4 \times 10^{-6}$	90

$\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$	$CL\%$
$<1.2 \times 10^{-6}$	90

$\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$	$CL\%$
$<2.0 \times 10^{-6}$	90

$\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$	$CL\%$
$<2.5 \times 10^{-6}$	90

$\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$	$CL\%$
$<2.0 \times 10^{-6}$	90

$\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$	$CL\%$
$<2.0 \times 10^{-6}$	90

$\Gamma(\chi_{c1} \text{anything})/\Gamma_{\text{total}}$	$EVTS$
$2.24 \pm 0.44 \pm 0.20$	376

$\Gamma(\chi_{c1}(1P)^0 X_{\text{tetra}})/\Gamma_{\text{total}}$	$CL\%$
$<36.7 \times 10^{-6}$	90

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 4.4×10^{-6} to 36.7×10^{-6} .

$\Gamma(\chi_{c2} \text{anything})/\Gamma_{\text{total}}$	$EVTS$
$2.28 \pm 0.73 \pm 0.34$	

$\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$	$CL\%$
$<5.1 \times 10^{-6}$	90

$\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$	$CL\%$
$<4.7 \times 10^{-6}$	90

$DOCUMENT\ ID$	$TECN$	$COMMENT$
YANG	14	$e^+e^- \rightarrow J/\psi X$

$DOCUMENT\ ID$	$TECN$	$COMMENT$
YANG	14	$e^+e^- \rightarrow J/\psi X$

$DOCUMENT\ ID$	$TECN$	$COMMENT$
YANG	14	$e^+e^- \rightarrow J/\psi X$

$DOCUMENT\ ID$	$TECN$	$COMMENT$
YANG	14	$e^+e^- \rightarrow J/\psi X$

$DOCUMENT\ ID$	$TECN$	$COMMENT$
YANG	14	$e^+e^- \rightarrow J/\psi X$

$DOCUMENT\ ID$	$TECN$	$COMMENT$
JIA	17	$\gamma(2S) \rightarrow \gamma J/\psi(1S)$

$DOCUMENT\ ID$	$TECN$	$COMMENT$
1 JIA	17A	$e^+e^- \rightarrow \text{hadrons}$

$DOCUMENT\ ID$	$TECN$	$COMMENT$
JIA	17	$\gamma(2S) \rightarrow \gamma J/\psi(1S)$

$DOCUMENT\ ID$	$TECN$	$COMMENT$
YANG	14	$e^+e^- \rightarrow \psi(2S)X$

$DOCUMENT\ ID$	$TECN$	$COMMENT$
YANG	14	$e^+e^- \rightarrow \psi(2S)X$

$\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>
$<2.5 \times 10^{-6}$	90

 Γ_{21}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
YANG	BELL	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>
$<1.9 \times 10^{-6}$	90

 Γ_{22}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
YANG	BELL	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>
$<3.3 \times 10^{-6}$	90

 Γ_{23}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
YANG	BELL	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>
$<3.9 \times 10^{-6}$	90

 Γ_{24}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
YANG	BELL	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>
$<3.9 \times 10^{-6}$	90

 Γ_{25}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
YANG	BELL	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(Z_c(3900)^+ Z_c(3900)^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>
$<1.0 \times 10^{-6}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1 JIA	BELL	$\gamma(2S) \rightarrow J/\psi \pi^\pm X$

 Γ_{26}/Γ

¹ Assuming $B(Z_c(3900)^\pm \rightarrow J/\psi \pi^\pm) = 1$.

 $\Gamma(Z_c(4200)^+ Z_c(4200)^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>
$<16.7 \times 10^{-6}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1 JIA	BELL	$\gamma(1S) \rightarrow J/\psi \pi^\pm X$

¹ Assuming $B(Z_c(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1$

 Γ_{27}/Γ

<u>VALUE</u>	<u>CL%</u>
$<7.3 \times 10^{-6}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1 JIA	BELL	$\gamma(2S) \rightarrow J/\psi \pi^\pm X$

¹ Assuming $B(Z_c(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1 = B(Z_c(3900)^\pm \rightarrow J/\psi \pi^\pm)$.

 Γ_{28}/Γ $\Gamma(X(4050)^+ X(4050)^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>
$<13.5 \times 10^{-6}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1 JIA	BELL	$\gamma(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

¹ Assuming $B(X(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$

 Γ_{29}/Γ $\Gamma(X(4250)^+ X(4250)^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>
$<26.7 \times 10^{-6}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1 JIA	BELL	$\gamma(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

¹ Assuming $B(X(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$

 Γ_{30}/Γ

$\Gamma(X(4050)^{\pm} X(4250)^{\mp})/\Gamma_{\text{total}}$					Γ_{31}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<27.2 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \chi_{c1}(1P)\pi^{\pm}X$	
¹ Assuming $B(X(4050)^{\pm} \rightarrow \chi_{c1}(1P)\pi^{\pm}) = 1 = B(X(4250)^{\pm} \rightarrow \chi_{c1}(1P)\pi^{\pm})$					

$\Gamma(Z_c(4430)^+ Z_c(4430)^-)/\Gamma_{\text{total}}$					Γ_{32}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<20.3 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \psi(2S)\pi^{\pm}X$	
¹ Assuming $B(Z_c(4430)^{\pm} \rightarrow \psi(2P)\pi^{\pm}) = 1$					

$\Gamma(X(4055)^{\pm} X(4055)^{\mp})/\Gamma_{\text{total}}$					Γ_{33}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<11.1 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \psi(2S)\pi^{\pm}X$	
¹ Assuming $B(X(4055)^{\pm} \rightarrow \psi(2S)\pi^{\pm}) = 1$					

$\Gamma(X(4055)^{\pm} Z_c(4430)^{\mp})/\Gamma_{\text{total}}$					Γ_{34}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<21.1 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \psi(2S)\pi^{\pm}X$	
¹ Assuming $B(X(4055)^{\pm} \rightarrow \psi(2S)\pi^{\pm}) = 1 = B(Z_c(4430)^{\pm} \rightarrow \psi(2S)\pi^{\pm})$					

$\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$					Γ_{35}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.78^{+0.30}_{-0.26}$ OUR AVERAGE				Error includes scale factor of 1.2.	
$2.64 \pm 0.11^{+0.26}_{-0.21}$		LEES	14G BABR	$e^+ e^- \rightarrow \overline{2H} X$	
$3.37 \pm 0.50 \pm 0.25$	58	ASNER	07 CLEO	$e^+ e^- \rightarrow \overline{2H} X$	

$\Gamma(ggg)/\Gamma_{\text{total}}$					Γ_{37}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
58.8 ± 1.2	6M	1 BESSON	06A CLEO	$\gamma(2S) \rightarrow \text{hadrons}$	
¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$ from BESSON 06A and PDG 08 values of $B(\pi^+\pi^-\gamma(1S)) = (18.1 \pm 0.4)\%$, $B(\pi^0\pi^0\gamma(1S)) = (8.6 \pm 0.4)\%$, $B(\mu^+\mu^-) = (1.93 \pm 0.17)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.					

$\Gamma(\gamma gg)/\Gamma(ggg)$					Γ_{38}/Γ_{37}
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
$3.18 \pm 0.04 \pm 0.47$	6M	BESSON	06A CLEO	$\gamma(2S) \rightarrow (\gamma +) \text{ hadrons}$	

$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$					Γ_{39}/Γ
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.58 \pm 0.33 \pm 0.18$	58	SHEN	12A BELL	$\gamma(1S) \rightarrow 2(K^+ K^-)$	

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>
<2.58	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$

 Γ_{40}/Γ $\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>
$2.32 \pm 0.40 \pm 0.54$	135

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 Γ_{41}/Γ $\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>
<1.33	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow 2(K^+ K^-)$

 Γ_{42}/Γ $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>
<0.57	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$

 Γ_{43}/Γ $\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>
<0.88	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$

 Γ_{44}/Γ $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>
$1.53 \pm 0.52 \pm 0.19$	32

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 Γ_{45}/Γ $\Gamma(K_1(1270)^{\pm} K^{\mp})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>
<3.22	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 Γ_{46}/Γ $\Gamma(K_1(1400)^{\pm} K^{\mp})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>
<0.83	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 Γ_{47}/Γ $\Gamma(b_1(1235)^{\pm} \pi^{\mp})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>
<0.40	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$

 Γ_{48}/Γ $\Gamma(\rho\pi)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>
<1.16	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13 BELL	$\gamma(2S) \rightarrow \pi^+ \pi^- \pi^0$

 Γ_{49}/Γ $\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>
<0.80	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13 BELL	$\gamma(2S) \rightarrow \pi^+ \pi^- \pi^0$

 Γ_{50}/Γ

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>
<1.63	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13	BELL $\gamma(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

 Γ_{51}/Γ $\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>
$13.0 \pm 1.9 \pm 2.1$	261 ± 37

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13	BELL $\gamma(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

 Γ_{52}/Γ $\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>EVTS</u>
$1.14 \pm 0.30 \pm 0.13$	40 ± 10	

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13	BELL $\gamma(2S) \rightarrow K_S^0 K^- \pi^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.2	90	¹ DOBBS	12A	$\gamma(2S) \rightarrow K_S^0 K^- \pi^+$
--------	----	--------------------	-----	--

¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>
<4.22	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13	BELL $\gamma(2S) \rightarrow K_S^0 K^- \pi^+$

 Γ_{54}/Γ $\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>
<1.45	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
SHEN	13	BELL $\gamma(2S) \rightarrow K_S^0 K^- \pi^+$

 Γ_{55}/Γ $\Gamma(f_1(1285)\text{anything})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>
$2.20 \pm 1.50 \pm 0.63$	$2.9k$

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

 Γ_{56}/Γ $\Gamma(f_1(1285)X_{\text{tetra}})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>
$<64.7 \times 10^{-6}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
¹ JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

 Γ_{57}/Γ

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 7.8×10^{-6} to 64.7×10^{-6} .

 $\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-2})	<u>EVTS</u>
0.29 ± 0.03	

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1,2 DOBBS	12A	$\gamma(2S) \rightarrow \text{hadrons}$

 Γ_{58}/Γ

¹ DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

² Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

$\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$

VALUE	EVTS
0.069 ± 0.004 OUR AVERAGE	
0.0693 ± 0.0012 ± 0.0041	407k
0.069 ± 0.005 ± 0.009	
0.091 ± 0.018 ± 0.022	
0.065 ± 0.007 ± 0.012	
0.080 ± 0.017 ± 0.016	
0.059 ± 0.014	

 Γ_{59}/Γ

DOCUMENT ID	TECN	COMMENT
ARTUSO 05	CLEO	$e^+e^- \rightarrow \gamma X$
EDWARDS 99	CLE2	$\gamma(2S) \rightarrow \gamma\chi(1P)$
ALBRECHT 85E	ARG	$e^+e^- \rightarrow \gamma\text{conv. } X$
NERNST 85	CBAL	$e^+e^- \rightarrow \gamma X$
HAAS 84	CLEO	$e^+e^- \rightarrow \gamma\text{conv. } X$
KLOPFEN... 83	CUSB	$e^+e^- \rightarrow \gamma X$

 $\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$

VALUE	EVTS
0.0715 ± 0.0035 OUR AVERAGE	
0.0724 ± 0.0011 ± 0.0040	410k
0.074 ± 0.005 ± 0.008	
0.098 ± 0.021 ± 0.024	
0.058 ± 0.007 ± 0.010	
0.102 ± 0.018 ± 0.021	
0.061 ± 0.014	

 Γ_{60}/Γ

DOCUMENT ID	TECN	COMMENT
ARTUSO 05	CLEO	$e^+e^- \rightarrow \gamma X$
EDWARDS 99	CLE2	$\gamma(2S) \rightarrow \gamma\chi(1P)$
ALBRECHT 85E	ARG	$e^+e^- \rightarrow \gamma\text{conv. } X$
NERNST 85	CBAL	$e^+e^- \rightarrow \gamma X$
HAAS 84	CLEO	$e^+e^- \rightarrow \gamma\text{conv. } X$
KLOPFEN... 83	CUSB	$e^+e^- \rightarrow \gamma X$

 $\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$

VALUE	EVTS
0.038 ± 0.004 OUR AVERAGE	
0.0375 ± 0.0012 ± 0.0047	198k
0.034 ± 0.005 ± 0.006	
0.064 ± 0.014 ± 0.016	
0.036 ± 0.008 ± 0.009	
0.044 ± 0.023 ± 0.009	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.035 ± 0.014	

 Γ_{61}/Γ

DOCUMENT ID	TECN	COMMENT
ARTUSO 05	CLEO	$e^+e^- \rightarrow \gamma X$
EDWARDS 99	CLE2	$\gamma(2S) \rightarrow \gamma\chi(1P)$
ALBRECHT 85E	ARG	$e^+e^- \rightarrow \gamma\text{conv. } X$
NERNST 85	CBAL	$e^+e^- \rightarrow \gamma X$
HAAS 84	CLEO	$e^+e^- \rightarrow \gamma\text{conv. } X$
KLOPFEN... 83	CUSB	$e^+e^- \rightarrow \gamma X$

 $\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%
<59	90

DOCUMENT ID	TECN	COMMENT
1 ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5.9	90	2 ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma\pi^+\pi^-$
-------	----	---------------	-----	---

¹ Re-evaluated assuming $B(f_0(1710) \rightarrow K^+ K^-) = 0.19$.

² Includes unknown branching ratio of $f_0(1710) \rightarrow \pi^+\pi^-$.

 Γ_{62}/Γ $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%
<53	90

DOCUMENT ID	TECN	COMMENT
1 ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma K^+ K^-$

¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$.

 Γ_{63}/Γ $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%
<24.1	90

DOCUMENT ID	TECN	COMMENT
1 ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma\pi^+\pi^-$

¹ Using $B(f_2(1270) \rightarrow \pi\pi) = 0.84$.

 Γ_{64}/Γ

$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$ Γ_{65}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.8 90 ¹ ALBRECHT 89 ARG $\gamma(2S) \rightarrow \gamma K^+ K^-$

¹ Includes unknown branching ratio of $f_J(2220) \rightarrow K^+ K^-$.

 $\Gamma(\gamma \eta_c(1S))/\Gamma_{\text{total}}$ Γ_{66}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.7 \times 10^{-5}$	90	WANG	11B	BELL $\gamma(2S) \rightarrow \gamma X$

 $\Gamma(\gamma \chi_{c0})/\Gamma_{\text{total}}$ Γ_{67}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.0 \times 10^{-4}$	90	WANG	11B	BELL $\gamma(2S) \rightarrow \gamma X$

 $\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$ Γ_{68}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.6 \times 10^{-6}$	90	WANG	11B	BELL $\gamma(2S) \rightarrow \gamma X$

 $\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$ Γ_{69}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.5 \times 10^{-5}$	90	WANG	11B	BELL $\gamma(2S) \rightarrow \gamma X$

 $\Gamma(\gamma \chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi)/\Gamma_{\text{total}}$ Γ_{70}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<0.8 \times 10^{-6}$	90	WANG	11B	BELL $\gamma(2S) \rightarrow \gamma X$

 $\Gamma(\gamma \chi_{c1}(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi)/\Gamma_{\text{total}}$ Γ_{71}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.4 \times 10^{-6}$	90	WANG	11B	BELL $\gamma(2S) \rightarrow \gamma X$

 $\Gamma(\gamma X(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$ Γ_{72}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.8 \times 10^{-6}$	90	WANG	11B	BELL $\gamma(2S) \rightarrow \gamma X$

 $\Gamma(\gamma \chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$ Γ_{73}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.2 \times 10^{-6}$	90	WANG	11B	BELL $\gamma(2S) \rightarrow \gamma X$

 $\Gamma(\gamma X(4350) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$ Γ_{74}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.3 \times 10^{-6}$	90	WANG	11B	BELL $\gamma(2S) \rightarrow \gamma X$

$\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$ Γ_{75}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	-------------	--------------------	-------------	----------------

5.5 $^{+1.1}_{-0.9}$ OUR AVERAGE Error includes scale factor of 1.2.

$6.1^{+0.6+0.9}_{-0.7-0.6}$ 29k FULSUM 18 BELL $\gamma(2S) \rightarrow \gamma X$

$3.9 \pm 1.1^{+1.1}_{-0.9}$ $13 \pm 5k$ ¹AUBERT 09AQ BABR $\gamma(2S) \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21 90 LEES 11J BABR $\gamma(2S) \rightarrow X\gamma$

< 8.4 90 ¹BONVICINI 10 CLEO $\gamma(2S) \rightarrow \gamma X$

< 5.1 90 ²ARTUSO 05 CLEO $e^+e^- \rightarrow \gamma X$

¹ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV.

² Superseded by BONVICINI 10.

 $\Gamma(\gamma\eta_b(1S) \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$ Γ_{76}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.7 \times 10^{-6}$	90	SANDILYA	13	$\gamma(2S) \rightarrow \gamma$ hadrons

 $\Gamma(\gamma X_{b\bar{b}} \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$ Γ_{77}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 4.9	90	SANDILYA	13	BELL	$\gamma(2S) \rightarrow \gamma$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

$46.2^{+29.7}_{-14.2} \pm 10.6$ 10 ¹DOBBS 12 $\gamma(2S) \rightarrow \gamma$ hadrons

¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ Γ_{78}/Γ

(1.5 GeV < m_X < 5.0 GeV)

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.95	95	ROSNER	07A	CLEO $e^+e^- \rightarrow \gamma X$

 $\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$ Γ_{79}/Γ

(0.3 GeV < m_{A^0} < 7 GeV)

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<8 \times 10^{-5}$	90	¹ LEES	11H	BABR $\gamma(2S) \rightarrow \gamma$ hadrons

¹ For a narrow scalar or pseudoscalar A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} .

 $\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{80}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8.3	90	¹ AUBERT	09Z	BABR $e^+e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$

¹ For a narrow scalar or pseudoscalar a_1^0 with mass in the range 212–9300 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of $m_{a_1^0}$ range from 0.26×10^{-6} to 8.3×10^{-6} .

LEPTON FAMILY NUMBER (*LF*) VIOLATING MODES **$\Gamma(e^\pm\tau^\mp)/\Gamma_{\text{total}}$**

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{81}/Γ
<3.2	90	LEES	10B	BABR	$e^+e^- \rightarrow e^\pm\tau^\mp$

 $\Gamma(\mu^\pm\tau^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{82}/Γ
< 3.3	90	LEES	10B	BABR	$e^+e^- \rightarrow \mu^\pm\tau^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<14.4	95	LOVE	08A	CLEO	$e^+e^- \rightarrow \mu^\pm\tau^\mp$

 $\Upsilon(2S)$ Cross-Particle Branching Ratios **$B(\Upsilon(2S) \rightarrow \pi^+\pi^-) \times B(\Upsilon(3S) \rightarrow \Upsilon(2S)X)$**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{83}/Γ
$1.78 \pm 0.02 \pm 0.11$	906k	LEES	11C	BABR	$e^+e^- \rightarrow \pi^+\pi^- X$

 $\Upsilon(2S)$ REFERENCES

FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)
JIA	18	PR D97 112004	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)
SANDILYA	13	PRL 111 112001	S. Sandilya <i>et al.</i>	(BELLE Collab.)
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
TAMPONI	13	PR D87 011104	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>	
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)

ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE $e^+ e^-$ Physics 412	W. Buchmueller, S. Cooper Editors: A. Ali and P. Soeding, World Scientific, Singapore	(HANN, DESY, MIT)
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
LURZ	87	ZPHY C36 383	B. Lurz <i>et al.</i>	(Crystal Ball Collab.)
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85	ZPHY C28 45	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
GELPHMAN	85	PR D32 2893	D. Gelfphman <i>et al.</i>	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BARBER	84	PL 135B 498	D.P. Barber <i>et al.</i>	
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
FONSECA	84	NP B242 31	V. Fonseca <i>et al.</i>	(CUSB Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
HAAS	84B	PR D30 1996	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
NICZYPORUK	81B	PL 100B 95	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
NICZYPORUK	81C	PL 99B 169	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)
