

X(2370)

$$I^G(J^{PC}) = ??(???)$$

OMITTED FROM SUMMARY TABLE

X(2370) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2341.6 ± 6.5 ± 5.7		¹ ABLIKIM	20Q	BES3 $J/\psi \rightarrow \gamma K \bar{K} \eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2376.3 ± 8.7 ^{+3.2} _{-4.3}	565	ABLIKIM	11C	BES3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$

¹The state observed by ABLIKIM 11C at 2120 MeV is not observed with 90% CL upper limit of 1.49×10^{-5} for $J/\psi \rightarrow \gamma X(2120) \rightarrow \gamma K^+ K^- \eta'$ and 6.38×10^{-6} for $K_S^0 K_S^0 \eta'$.

X(2370) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
117 ± 10 ± 8	¹ ABLIKIM	20Q	BES3 $J/\psi \rightarrow \gamma K \bar{K} \eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
83 ± 17 ⁺⁴⁴ ₋₆	ABLIKIM	11C	BES3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$

¹The state observed by ABLIKIM 11C at 2120 MeV is not observed with 90% CL upper limit of 1.49×10^{-5} for $J/\psi \rightarrow \gamma X(2120) \rightarrow \gamma K^+ K^- \eta'$ and 6.38×10^{-6} for $K_S^0 K_S^0 \eta'$.

X(2370) DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^+ K^- \eta'$	seen
Γ_2 $K_S^0 K_S^0 \eta'$	seen
Γ_3 $\pi^+ \pi^- \eta'$	seen

X(2370) BRANCHING RATIOS

$\Gamma(K^+ K^- \eta')/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	ABLIKIM	20Q	BES3 $J/\psi \rightarrow \gamma K^+ K^- \eta'$	
$\Gamma(K_S^0 K_S^0 \eta')/\Gamma_{\text{total}}$				Γ_2/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	ABLIKIM	20Q	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$	
$\Gamma(\pi^+ \pi^- \eta')/\Gamma_{\text{total}}$				Γ_3/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	ABLIKIM	11C	BES3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$	

X(2370) REFERENCES

ABLIKIM	20Q	EPJ C80 746	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
