

$\eta_b(1S)$


$$I^G(J^{PC}) = 0^+(0^{-+})$$

Quantum numbers shown are quark-model predictions. Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$.

$\eta_b(1S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
9398.7 ± 2.0 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
9394.8 ^{+2.7} _{-3.1}	4.5 2.7	29k	FULSOM 18 BELL	$\Upsilon(2S) \rightarrow \gamma X$
9400.7 ± 1.7	± 1.6	33.1k	TAMPONI 15 BELL	$e^+ e^- \rightarrow \gamma \eta + \text{hadrons}$
9402.4 ± 1.5	± 1.8	34k	¹ MIZUK 12 BELL	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- + \text{hadrons}$
9391.8 ± 6.6	± 2.0	2.3k	² BONVICINI 10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
9394.2 ^{+4.8} _{-4.9}	± 2.0	13k	² AUBERT 09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
9388.9 ^{+3.1} _{-2.3}	± 2.7	19k	² AUBERT 08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
9393.2 ± 3.4	± 2.3	10	^{2,3} DOBBS 12	$\Upsilon(2S) \rightarrow \gamma \text{hadrons}$
9300 ± 20	± 20		HEISTER 02D ALEP	181–209 $e^+ e^-$


¹ With floating width. Not independent of the corresponding mass difference measurement.
² Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding γ energy or mass difference measurements.
³ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.



$m_{\Upsilon(1S)} - m_{\eta_b}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
62.3 ± 3.2 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.			
$57.9 \pm 1.5 \pm 1.8$	34k	¹ MIZUK	12 BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- + \text{hadrons}$
$68.5 \pm 6.6 \pm 2.0$	$2.3 \pm 0.5k$	² BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
$66.1^{+4.8}_{-4.9} \pm 2.0$	$13 \pm 5k$	² AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
$71.4^{+2.3}_{-3.1} \pm 2.7$	$19 \pm 3k$	² AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$67.1 \pm 3.4 \pm 2.3$	10^{+5}_{-4}	^{2,3} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma \text{hadrons}$

- ¹ With floating width. Not independent of the corresponding mass measurement.
² Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding γ energy or mass measurements.
³ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.



γ ENERGY IN $\Upsilon(3S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$920.6^{+2.8}_{-3.2}$ OUR AVERAGE				
$918.6 \pm 6.0 \pm 1.9$	$2.3 \pm 0.5k$	¹ BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
$921.2^{+2.1}_{-2.8} \pm 2.4$	$19 \pm 3k$	¹ AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$

¹ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.

γ ENERGY IN $\Upsilon(2S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
609.3^{+4.6}_{-4.5} ± 1.9	13 ± 5k	¹ AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$

¹ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.

$\eta_b(1S)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10 ⁺⁵₋₄ OUR AVERAGE				
8 ⁺⁶ ₋₅ ± 5	33.1k	¹ TAMPONI	15 BELL	$e^+ e^- \rightarrow \gamma \eta + \text{hadrons}$
10.8 ^{+4.0+4.5} _{-3.7-2.0}	34k	¹ MIZUK	12 BELL	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- + \text{hadrons}$

¹ With floating mass.

$\eta_b(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 hadrons	seen	
Γ_2 $3h^+ 3h^-$	not seen	
Γ_3 $2h^+ 2h^-$	not seen	
Γ_4 $4h^+ 4h^-$	not seen	
Γ_5 $\gamma\gamma$	not seen	
Γ_6 $\mu^+ \mu^-$	$< 9 \times 10^{-3}$	90%
Γ_7 $\tau^+ \tau^-$	$< 8 \%$	90%

$\eta_b(1S)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(3h^+ 3h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2 \Gamma_5 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<470	95	ABDALLAH	06 DLPH	161–209 $e^+ e^-$
<132	95	HEISTER	02D ALEP	181–209 $e^+ e^-$

$\Gamma(2h^+ 2h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_3 \Gamma_5 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<190	95	ABDALLAH	06 DLPH	161–209 $e^+ e^-$
< 48	95	HEISTER	02D ALEP	181–209 $e^+ e^-$

$\Gamma(4h^+4h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_4\Gamma_5/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<660	95	ABDALLAH 06	DLPH	161–209 e^+e^-

$\eta_b(1S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	34k	MIZUK 12	BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- + \text{hadrons}$

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<9 × 10⁻³	90	¹ AUBERT 09Z	BABR	$e^+e^- \rightarrow \Upsilon(2S,3S) \rightarrow \gamma\eta_b$

¹Obtained using $B(\Upsilon(2S) \rightarrow \gamma\eta_b) = (4.2_{-1.0}^{+1.1} \pm 0.9) \times 10^{-4}$ and $B(\Upsilon(3S) \rightarrow \gamma\eta_b) = (4.8 \pm 0.5 \pm 0.6) \times 10^{-4}$. This limit is equivalent to $B(\eta_b \rightarrow \mu^+\mu^-) = (-0.25 \pm 0.51 \pm 0.33)\%$ measurement.

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8 × 10⁻²	90	AUBERT 09P	BABR	$e^+e^- \rightarrow \gamma\tau^+\tau^-$

$\eta_b(1S)$ REFERENCES

FULSOM 18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)
TAMPONI 15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS 12	PRL 109 082001	S. Dobbs <i>et al.</i>	
MIZUK 12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)
BONVICINI 10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
AUBERT 09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABDALLAH 06	PL B634 340	J.M. Abdallah <i>et al.</i>	(DELPHI Collab.)
HEISTER 02D	PL B530 56	A. Heister <i>et al.</i>	(ALEPH Collab.)