


# $f_2(1950)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

## $f_2(1950)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1936 ± 12</b>	<b>OUR AVERAGE</b>		Error includes scale factor of 1.3. See the ideogram below.
1940 ± 50	BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
1980 ± 22	<sup>1</sup> BARBERIS	00C	450 $pp \rightarrow pp4\pi$
1940 ± 22	<sup>2</sup> BARBERIS	00C	450 $pp \rightarrow pp2\pi2\pi^0$
1960 ± 30	BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+\pi^-)$
1918 ± 12	ANTINORI	95 OMEG	300,450 $pp \rightarrow pp2(\pi^+\pi^-)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1978.2 ± 1.8 <sup>+28.4</sup> <sub>-16.9</sub>	<sup>3</sup> ALBRECHT	20 RVUE	0.9 $\bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$
2038 <sup>+13</sup> <sub>-11</sub> <sup>+12</sup> <sub>-73</sub>	<sup>4</sup> UEHARA	09 BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
1930 ± 25	<sup>5</sup> BINON	05 GAMS	33 $\pi^-p \rightarrow \eta\eta n$
1980 ± 2 ± 14	ABE	04 BELL	10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
1867 ± 46	<sup>6</sup> AMSLER	02 CBAR	0.9 $\bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
2010 ± 25	ANISOVICH	00J SPEC	
1980 ± 50	ANISOVICH	99B SPEC	1.35–1.94 $p\bar{p} \rightarrow \eta\eta\pi^0$
~ 1990	<sup>7</sup> OAKDEN	94 RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
1950 ± 15	<sup>8</sup> ASTON	91 LASS	11 $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$



- <sup>1</sup> Decaying into  $\pi^+\pi^-2\pi^0$ .
- <sup>2</sup> Decaying into  $2(\pi^+\pi^-)$ .
- <sup>3</sup> T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).
- <sup>4</sup> Taking into account  $f_4(2050)$ .
- <sup>5</sup> First solution, PWA is ambiguous.
- <sup>6</sup> T-matrix pole.
- <sup>7</sup> From solution B of amplitude analysis of data on  $\bar{p}p \rightarrow \pi\pi$ . See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.
- <sup>8</sup> Cannot determine spin to be 2.

### $f_2(1950)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>464 ± 24 OUR AVERAGE</b>			
380 <sup>+120</sup> <sub>-90</sub>	BAI	00A	BES $J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
520 ± 50	<sup>1</sup> BARBERIS	00C	450 $pp \rightarrow pp4\pi$
485 ± 55	<sup>2</sup> BARBERIS	00C	450 $pp \rightarrow pp4\pi$
460 ± 40	BARBERIS	97B	OMEG 450 $pp \rightarrow pp2(\pi^+\pi^-)$
390 ± 60	ANTINORI	95	OMEG 300,450 $pp \rightarrow pp2(\pi^+\pi^-)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
237.6 ± 1.6 <sup>+41.6</sup> <sub>-15.5</sub>	<sup>3</sup> ALBRECHT	20	RVUE 0.9 $\bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$
441 <sup>+27</sup> <sub>-25</sub> <sup>+28</sup> <sub>-192</sub>	<sup>4</sup> UEHARA	09	BELL 10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
450 ± 50	<sup>5</sup> BINON	05	GAMS 33 $\pi^-p \rightarrow \eta\eta n$
297 ± 12 ± 6	ABE	04	BELL 10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
385 ± 58	<sup>6</sup> AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
495 ± 35	ANISOVICH	00J	SPEC
500 ± 100	ANISOVICH	99B	SPEC 1.35–1.94 $p\bar{p} \rightarrow \eta\eta\pi^0$
~ 100	<sup>7</sup> OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
250 ± 50	<sup>8</sup> ASTON	91	LASS 11 $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$

- <sup>1</sup> Decaying into  $\pi^+\pi^-2\pi^0$ .
- <sup>2</sup> Decaying into  $2(\pi^+\pi^-)$ .
- <sup>3</sup> T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).
- <sup>4</sup> Taking into account  $f_4(2050)$ .
- <sup>5</sup> First solution, PWA is ambiguous.
- <sup>6</sup> T-matrix pole.
- <sup>7</sup> From solution B of amplitude analysis of data on  $\bar{p}p \rightarrow \pi\pi$ . See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.
- <sup>8</sup> Cannot determine spin to be 2.

## $f_2(1950)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K^*(892)\bar{K}^*(892)$	seen
$\Gamma_2$ $\pi\pi$	
$\Gamma_3$ $\pi^+\pi^-$	seen
$\Gamma_4$ $\pi^0\pi^0$	seen
$\Gamma_5$ $4\pi$	seen
$\Gamma_6$ $\pi^+\pi^-\pi^+\pi^-$	
$\Gamma_7$ $a_2(1320)\pi$	
$\Gamma_8$ $f_2(1270)\pi\pi$	
$\Gamma_9$ $\eta\eta$	seen
$\Gamma_{10}$ $K\bar{K}$	seen
$\Gamma_{11}$ $\gamma\gamma$	seen
$\Gamma_{12}$ $\rho\bar{\rho}$	seen

### $f_2(1950)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{10}\Gamma_{11}/\Gamma$
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

• • • We do not use the following data for averages, fits, limits, etc. • • •

122 ± 4 ± 26                      <sup>1</sup> ABE                      04    BELL    10.6  $e^+e^- \rightarrow e^+e^-K^+K^-$

<sup>1</sup> Assuming spin 2.

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_{11}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

• • • We do not use the following data for averages, fits, limits, etc. • • •

162 <sup>+69</sup> <sub>-42</sub> <sup>+1137</sup> <sub>-204</sub>                      <sup>1</sup> UEHARA                      09    BELL    10.6  $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$

<sup>1</sup> Taking into account  $f_4(2050)$ .

### $f_2(1950)$ BRANCHING RATIOS

$\Gamma(K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>

seen                                      ASTON                      91    LASS    0    11  $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$

$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$	$\Gamma_7/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen                                      BARBERIS                      00B                      450  $pp \rightarrow p_f\eta\pi^+\pi^-p_s$

not seen                                      BARBERIS                      00C                      450  $pp \rightarrow p_f4\pi p_s$

possibly seen                                      BARBERIS                      97B    OMEG    450  $pp \rightarrow p\rho2(\pi^+\pi^-)$

$\Gamma(\eta\eta)/\Gamma(4\pi)$				$\Gamma_9/\Gamma_5$
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<5.0 \times 10^{-3}$	90	BARBERIS	00E 450 $p p \rightarrow p_f \eta \eta p_s$	

$\Gamma(\eta\eta)/\Gamma(\pi^+\pi^-)$				$\Gamma_9/\Gamma_3$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>0.14 ± 0.05</b>	AMSLER	02	CBAR 0.9 $\bar{p} p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$	

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$				$\Gamma_{12}/\Gamma$
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	111	ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma p \bar{p}$

### $f_2(1950)$ REFERENCES

ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)
BINON	05	PAN 68 960	F. Binon <i>et al.</i>	
		Translated from YAF 68 998.		
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>	
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+) JP
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
ASTON	91	NPBPS B21 5	D. Aston <i>et al.</i>	(LASS Collab.)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)