



$$I(J^P) = ?(?^?) \quad \text{Status: } ***$$

Ξ_c^{++} MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$3621.55 \pm 0.23 \pm 0.30$	2k	¹ AAIJ	20J LHCb	pp at 13 TeV
$3620.6 \pm 1.5 \pm 0.4 \pm 0.3$	91	² AAIJ	18BA LHCb	pp at 13 TeV
$3621.40 \pm 0.72 \pm 0.27 \pm 0.14$	313	³ AAIJ	17BC LHCb	pp at 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ AAIJ 20J combines mass measurements $3621.53 \pm 0.24 \pm 0.29$ MeV from $\Xi_c^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+$ and $3621.95 \pm 0.60 \pm 0.49$ MeV from $\Xi_c^{++} \rightarrow \Xi_c^+ \pi^+$. Supersedes AAIJ 18BA and AAIJ 17BC.

² The third error in AAIJ 18BA value is from the uncertainty of the Ξ_c^+ mass.

³ The third error in AAIJ 17BC value is from the uncertainty of the Λ_c^+ mass. The width of the signal is 6.6 ± 0.8 MeV, consistent with the experimental resolution.

Ξ_c^{++} MEAN LIFE

VALUE (10^{-15} s)	EVTS	DOCUMENT ID	TECN	COMMENT
$256_{-22}^{+24} \pm 14$	304	AAIJ	18G LHCb	pp at 13 TeV

Ξ_c^{++} DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \Lambda_c^+ K^- \pi^+ \pi^+$	seen
$\Gamma_2 \quad \Xi_c^+ \pi^+, \Xi_c^+ \rightarrow p K^- \pi^+$	seen
$\Gamma_3 \quad D^+ p K^- \pi^+$	

$\Gamma(\Lambda_c^+ K^- \pi^+ \pi^+)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAIJ	17BC LHCb	12 std significance

$\Gamma(\Xi_c^+ \pi^+, \Xi_c^+ \rightarrow p K^- \pi^+)/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	91	AAIJ	18BA LHCb	5.9 std significance

$\Gamma(\Xi_c^+ \pi^+, \Xi_c^+ \rightarrow p K^- \pi^+)/\Gamma(\Lambda_c^+ K^- \pi^+ \pi^+) \quad \Gamma_2/\Gamma_1$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$2.2 \pm 0.6 \pm 0.1$	⁴ AAIJ	18BA LHCb	Ratio 91 over 289 events

⁴ AAIJ 18BA reports $[\Gamma(\Xi_{cc}^{++} \rightarrow \Xi_c^+ \pi^+, \Xi_c^+ \rightarrow p K^- \pi^+)/\Gamma(\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+)] / [B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = (3.5 \pm 0.9 \pm 0.3) \times 10^{-2}$ which we multiply by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (6.28 \pm 0.32) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^+ p K^- \pi^+)/\Gamma(\Lambda_c^+ K^- \pi^+ \pi^+)$				Γ_3/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-2}$	90	AAIJ	19AO LHCB	pp at 13 TeV

Ξ_{cc}^{++} REFERENCES

AAIJ	20J	JHEP 2002 049	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	19AO	JHEP 1910 124	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18BA	PRL 121 162002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18G	PRL 121 052002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BC	PRL 119 112001	R. Aaij <i>et al.</i>	(LHCb Collab.)