

$\Sigma(1580) 3/2^-$  $I(J^P) = 1(\frac{3}{2}^-)$  Status: \*

## OMITTED FROM SUMMARY TABLE

Seen in the isospin-1  $\bar{K}N$  cross section at BNL (LI 73, CARROLL 76) and in a partial-wave analysis of  $K^-p \rightarrow \Lambda\pi^0$  for c.m. energies 1560–1600 MeV by LITCHFIELD 74. LITCHFIELD 74 finds  $J^P = 3/2^-$ . Not seen by ENGLER 78, CAMERON 78C, OLMSTED 04, nor by PRAKHOV 04.

Neither ZHANG 13A nor SARANTSEV 19 see any evidence for this state.

 $\Sigma(1580)$  POLE POSITION

## REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1607^{+13}_{-11}$	<sup>1</sup> KAMANO	15	DPWA Multichannel
--------------------	---------------------	----	-------------------

<sup>1</sup> From the preferred solution A in KAMANO 15. Solution B reports  $M = 1492^{+4}_{-7}$  MeV.

## –2×IMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$253^{+30}_{-18}$	<sup>2</sup> KAMANO	15	DPWA Multichannel
-------------------	---------------------	----	-------------------

<sup>2</sup> From the preferred solution A in KAMANO 15. Solution B reports  $M = 138^{+8}_{-14}$  MeV.

 $\Sigma(1580)$  POLE RESIDUES

The “normalized residue” is the residue divided by  $\Gamma_{pole}/2$ .

Normalized residue in  $N\bar{K} \rightarrow \Sigma(1580) \rightarrow N\bar{K}$ 

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
---------	-----------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.00778	51	<sup>3</sup> KAMANO	15	DPWA Multichannel
---------	----	---------------------	----	-------------------

<sup>3</sup> From the preferred solution A in KAMANO 15.

Normalized residue in  $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Sigma\pi$ 

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
---------	-----------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0625	–6	<sup>4</sup> KAMANO	15	DPWA Multichannel
--------	----	---------------------	----	-------------------

<sup>4</sup> From the preferred solution A in KAMANO 15.

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Lambda\pi$** 

<u>MODULUS</u>	<u>PHASE (<math>^{\circ}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
----------------	--------------------------------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.059	156	<sup>5</sup> KAMANO	15	DPWA Multichannel
-------	-----	---------------------	----	-------------------

<sup>5</sup> From the preferred solution A in KAMANO 15.

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Sigma(1385)\pi$ , S-wave**

<u>MODULUS</u>	<u>PHASE (<math>^{\circ}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
----------------	--------------------------------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0368	-18	<sup>6</sup> KAMANO	15	DPWA Multichannel
--------	-----	---------------------	----	-------------------

<sup>6</sup> From the preferred solution A in KAMANO 15.

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Sigma(1385)\pi$ , D-wave**

<u>MODULUS</u>	<u>PHASE (<math>^{\circ}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
----------------	--------------------------------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0103	123	<sup>7</sup> KAMANO	15	DPWA Multichannel
--------	-----	---------------------	----	-------------------

<sup>7</sup> From the preferred solution A in KAMANO 15.

 **$\Sigma(1580)$  MASS**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	--------------------	-------------	----------------

 **$\approx 1580$  OUR ESTIMATE**

$1583 \pm 4$	<sup>8</sup> CARROLL	76	DPWA Isospin-1 total $\sigma$
$1582 \pm 4$	<sup>9</sup> LITCHFIELD	74	DPWA $K^- p \rightarrow \Lambda\pi^0$

<sup>8</sup> CARROLL 76 sees a total-cross-section bump with  $(J+1/2) \Gamma_{el} / \Gamma_{total} = 0.06$ .

<sup>9</sup> The main effect observed by LITCHFIELD 74 is in the  $\Lambda\pi$  final state; the  $\bar{K}N$  and  $\Sigma\pi$  couplings are estimated from a multichannel fit including total-cross-section data of LI 73.

 **$\Sigma(1580)$  WIDTH**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	--------------------	-------------	----------------

15	<sup>10</sup> CARROLL	76	DPWA Isospin-1 total $\sigma$
$11 \pm 4$	<sup>11</sup> LITCHFIELD	74	DPWA $K^- p \rightarrow \Lambda\pi^0$

<sup>10</sup> CARROLL 76 sees a total-cross-section bump with  $(J+1/2) \Gamma_{el} / \Gamma_{total} = 0.06$ .

<sup>11</sup> The main effect observed by LITCHFIELD 74 is in the  $\Lambda\pi$  final state; the  $\bar{K}N$  and  $\Sigma\pi$  couplings are estimated from a multichannel fit including total-cross-section data of LI 73.

 **$\Sigma(1580)$  DECAY MODES**

Mode
$\Gamma_1$ $N\bar{K}$
$\Gamma_2$ $\Lambda\pi$
$\Gamma_3$ $\Sigma\pi$
$\Gamma_4$ $\Sigma(1385)\pi$ , S-wave

$\Gamma_5$	$\Sigma(1385)\pi$ , <i>D</i> -wave
$\Gamma_6$	$N\bar{K}^*(892)$ , $S=1/2$ , <i>D</i> -wave
$\Gamma_7$	$N\bar{K}^*(892)$ , $S=3/2$ , <i>S</i> -wave
$\Gamma_8$	$N\bar{K}^*(892)$ , $S=3/2$ , <i>D</i> -wave

---

## $\Sigma(1580)$ BRANCHING RATIOS

See “Sign conventions for resonance couplings” in the Note on  $\Lambda$  and  $\Sigma$  Resonances.

### $\Gamma(N\bar{K})/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
$+0.03 \pm 0.01$	<sup>12</sup> LITCHFIELD 74	DPWA	$\bar{K}N$ multichannel
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
0.003	<sup>13</sup> KAMANO 15	DPWA	Multichannel

<sup>12</sup> The main effect observed by LITCHFIELD 74 is in the  $\Lambda\pi$  final state; the  $\bar{K}N$  and  $\Sigma\pi$  couplings are estimated from a multichannel fit including total-cross-section data of LI 73.

<sup>13</sup> From the preferred solution A in KAMANO 15.

### $\Gamma(\Lambda\pi)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
0.490	<sup>14</sup> KAMANO 15	DPWA	Multichannel

<sup>14</sup> From the preferred solution A in KAMANO 15.

### $\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
0.387	<sup>15</sup> KAMANO 15	DPWA	Multichannel

<sup>15</sup> From the preferred solution A in KAMANO 15.

### $\Gamma(\Sigma(1385)\pi, S\text{-wave})/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
0.12	<sup>16</sup> KAMANO 15	DPWA	Multichannel

<sup>16</sup> From the preferred solution A in KAMANO 15.

### $\Gamma(\Sigma(1385)\pi, D\text{-wave})/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
0.001	<sup>17</sup> KAMANO 15	DPWA	Multichannel

<sup>17</sup> From the preferred solution A in KAMANO 15.

### $\Gamma(N\bar{K}^*(892), S=1/2, D\text{-wave})/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
not seen	<sup>18</sup> KAMANO 15	DPWA	Multichannel

<sup>18</sup> From the preferred solution A in KAMANO 15.

$\Gamma(N\bar{K}^*(892), S=3/2, S\text{-wave})/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen <sup>19</sup>KAMANO 15 DPWA Multichannel<sup>19</sup>From the preferred solution A in KAMANO 15. $\Gamma(N\bar{K}^*(892), S=3/2, D\text{-wave})/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen <sup>20</sup>KAMANO 15 DPWA Multichannel<sup>20</sup>From the preferred solution A in KAMANO 15. $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$  in  $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Lambda\pi$   $(\Gamma_1\Gamma_2)^{1/2}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen CAMERON 78C HBC  $K_L^0 p \rightarrow \Lambda\pi^+$ not seen ENGLER 78 HBC  $K_L^0 p \rightarrow \Lambda\pi^+$ +0.10±0.02 <sup>21</sup>LITCHFIELD 74 DPWA  $K^- p \rightarrow \Lambda\pi^0$ <sup>21</sup>The main effect observed by LITCHFIELD 74 is in the  $\Lambda\pi$  final state; the  $\bar{K}N$  and  $\Sigma\pi$  couplings are estimated from a multichannel fit including total-cross-section data of LI 73. $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$  in  $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Sigma\pi$   $(\Gamma_1\Gamma_3)^{1/2}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen CAMERON 78C HBC  $K_L^0 p \rightarrow \Sigma^0\pi^+$ not seen ENGLER 78 HBC  $K_L^0 p \rightarrow \Sigma^0\pi^+$ +0.03±0.04 <sup>22</sup>LITCHFIELD 74 DPWA  $\bar{K}N$  multichannel<sup>22</sup>The main effect observed by LITCHFIELD 74 is in the  $\Lambda\pi$  final state; the  $\bar{K}N$  and  $\Sigma\pi$  couplings are estimated from a multichannel fit including total-cross-section data of LI 73. **$\Sigma(1580)$  REFERENCES**

SARANTSEV	19	EPJ A55 180	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
KAMANO	15	PR C92 025205	H. Kamano <i>et al.</i>	(ANL, OSAK)
ZHANG	13A	PR C88 035205	H. Zhang <i>et al.</i>	(KSU)
OLMSTED	04	PL B588 29	J. Olmsted <i>et al.</i>	(BNL Crystal Ball Collab.)
PRAKHOV	04	PR C69 042202	S. Prakhov <i>et al.</i>	(BNL Crystal Ball Collab.)
CAMERON	78C	NP B132 189	W. Cameron <i>et al.</i>	(BGNA, EDIN, GLAS+) I
ENGLER	78	PR D18 3061	A. Engler <i>et al.</i>	(CMU, ANL)
CARROLL	76	PRL 37 806	A.S. Carroll <i>et al.</i>	(BNL) I
LITCHFIELD	74	PL 51B 509	P.J. Litchfield	(CERN) IJP
LI	73	Purdue Conf. 283	K.K. Li	(BNL) I