

$K_0^*(1950)$

$$I(J^P) = \frac{1}{2}(0^+)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K^- \pi^+$ system. Needs confirmation.

$K_0^*(1950)$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | CHG | COMMENT |
|---|------------------------|------|--------|---|
| 1944 ± 18 OUR AVERAGE | | | | |
| 1942 ± 22 ± 21 | LEES | 21A | BABR | $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' K^+ K^-$ |
| 1945 ± 10 ± 20 | ¹ ASTON | 88 | LASS 0 | 11 $K^- p \rightarrow K^- \pi^+ n$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1917 ± 12 | ² ZHOU | 06 | RVUE | $K p \rightarrow K^- \pi^+ n$ |
| 1820 ± 40 | ³ ANISOVICH | 97C | RVUE | 11 $K^- p \rightarrow K^- \pi^+ n$ |

¹ We take the central value of the two solutions and the larger error given.

² S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$, $K_0^*(1430)$.

³ T-matrix pole. Reanalysis of ASTON 88 data.

$K_0^*(1950)$ WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | CHG | COMMENT |
|---|------------------------|------|--------|---|
| 100 ± 40 OUR AVERAGE Error includes scale factor of 1.3. | | | | |
| 80 ± 32 ± 20 | LEES | 21A | BABR | $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' K^+ K^-$ |
| 201 ± 34 ± 79 | ⁴ ASTON | 88 | LASS 0 | 11 $K^- p \rightarrow K^- \pi^+ n$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 145 ± 38 | ⁵ ZHOU | 06 | RVUE | $K p \rightarrow K^- \pi^+ n$ |
| 250 ± 100 | ⁶ ANISOVICH | 97C | RVUE | 11 $K^- p \rightarrow K^- \pi^+ n$ |

⁴ We take the central value of the two solutions and the larger error given.

⁵ S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$, $K_0^*(1430)$.

⁶ T-matrix pole. Reanalysis of ASTON 88 data.

$K_0^*(1950)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|----------------------------|--------------------------------|
| $\Gamma_1 \quad K^- \pi^+$ | (52 ± 14) % |

$K_0^*(1950)$ BRANCHING RATIOS

| $\Gamma(K^- \pi^+)/\Gamma_{\text{total}}$ | Γ_1/Γ | | | |
|---|--------------------|------|--------|------------------------------------|
| VALUE | DOCUMENT ID | TECN | CHG | COMMENT |
| 0.52 ± 0.08 ± 0.12 | ⁷ ASTON | 88 | LASS 0 | 11 $K^- p \rightarrow K^- \pi^+ n$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| ~ 0.60 | ⁸ ZHOU | 06 | RVUE | $K p \rightarrow K^- \pi^+ n$ |

⁷ We take the central value of the two solutions and the larger error given.

⁸ S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$, $K_0^*(1430)$.

$K_0^*(1950)$ REFERENCES

| | | | | |
|-----------|-----|----------------|--------------------------------|--------------------------|
| LEES | 21A | PR D104 072002 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| ZHOU | 06 | NP A775 212 | Z.Y. Zhou, H.Q. Zheng | |
| ANISOVICH | 97C | PL B413 137 | A.V. Anisovich, A.V. Sarantsev | |
| ASTON | 88 | NP B296 493 | D. Aston <i>et al.</i> | (SLAC, NAGO, CINC, INUS) |
