

$\psi(4160)$ 

$$J^{PC} = 0^{-}(1^{-}-)$$

### $\psi(4160)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4191 ± 5 OUR AVERAGE</b>			
4191 $\begin{smallmatrix} +9 \\ -8 \end{smallmatrix}$	AAIJ	13BC LHCB	$B^+ \rightarrow K^+ \mu^+ \mu^-$
4191.7 ± 6.5	<sup>1</sup> ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
4193 ± 7	<sup>2</sup> MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
4151 ± 4	<sup>3</sup> SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4155 ± 5	<sup>4</sup> SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4159 ± 20	BRANDELIK	78C DASP	$e^+ e^-$

<sup>1</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (293 \pm 57)^\circ$ .

<sup>2</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects.

<sup>3</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

<sup>4</sup> From a fit to BES (BAI 02C) data.

### $\psi(4160)$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>70 ± 10 OUR AVERAGE</b>			
65 $\begin{smallmatrix} +22 \\ -16 \end{smallmatrix}$	AAIJ	13BC LHCB	$B^+ \rightarrow K^+ \mu^+ \mu^-$
71.8 ± 12.3	<sup>1</sup> ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
79 ± 14	<sup>2</sup> MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
107 ± 10	<sup>3</sup> SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
107 ± 16	<sup>4</sup> SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
78 ± 20	BRANDELIK	78C DASP	$e^+ e^-$

<sup>1</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (293 \pm 57)^\circ$ .

<sup>2</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects.

<sup>3</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

<sup>4</sup> From a fit to BES (BAI 02C) data.

**$\psi(4160)$  DECAY MODES**

Due to the complexity of the  $c\bar{c}$  threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective  $\sqrt{s}$  near this particle’s central mass value, more (less) than  $2\sigma$  above zero, without regard to any peaking behavior in  $\sqrt{s}$  or absence thereof. See mode listing(s) for details and references.

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $e^+ e^-$	$(6.9 \pm 3.3) \times 10^{-6}$	
$\Gamma_2$ $\mu^+ \mu^-$	seen	
$\Gamma_3$ $D \bar{D}$	seen	
$\Gamma_4$ $D^0 \bar{D}^0$	seen	
$\Gamma_5$ $D^+ D^-$	seen	
$\Gamma_6$ $D^* \bar{D} + \text{c.c.}$	seen	
$\Gamma_7$ $D^*(2007)^0 \bar{D}^0 + \text{c.c.}$	seen	
$\Gamma_8$ $D^*(2010)^+ D^- + \text{c.c.}$	seen	
$\Gamma_9$ $D^* \bar{D}^*$	seen	
$\Gamma_{10}$ $D^*(2007)^0 \bar{D}^*(2007)^0$	seen	
$\Gamma_{11}$ $D^*(2010)^+ D^*(2010)^-$	seen	
$\Gamma_{12}$ $D^0 D^- \pi^+ + \text{c.c.}$ (excl. $D^*(2007)^0 \bar{D}^0 + \text{c.c.}$ , $D^*(2010)^+ D^- + \text{c.c.}$ )	not seen	
$\Gamma_{13}$ $D \bar{D}^* \pi + \text{c.c.}$ (excl. $D^* \bar{D}^*$ )	seen	
$\Gamma_{14}$ $D^0 D^{*-} \pi^+ + \text{c.c.}$ (excl. $D^*(2010)^+ D^*(2010)^-$ )	not seen	
$\Gamma_{15}$ $D_s^+ D_s^-$	not seen	
$\Gamma_{16}$ $D_s^{*+} D_s^- + \text{c.c.}$	seen	
$\Gamma_{17}$ $J/\psi \pi^+ \pi^-$	$< 3 \times 10^{-3}$	90%
$\Gamma_{18}$ $J/\psi \pi^0 \pi^0$	$< 3 \times 10^{-3}$	90%
$\Gamma_{19}$ $J/\psi K^+ K^-$	$< 2 \times 10^{-3}$	90%
$\Gamma_{20}$ $J/\psi \eta$	$< 8 \times 10^{-3}$	90%
$\Gamma_{21}$ $J/\psi \pi^0$	$< 1 \times 10^{-3}$	90%
$\Gamma_{22}$ $J/\psi \eta'$	$< 5 \times 10^{-3}$	90%
$\Gamma_{23}$ $J/\psi \pi^+ \pi^- \pi^0$	$< 1 \times 10^{-3}$	90%
$\Gamma_{24}$ $\psi(2S) \pi^+ \pi^-$	$< 4 \times 10^{-3}$	90%
$\Gamma_{25}$ $\chi_{c1} \gamma$	$< 5 \times 10^{-3}$	90%
$\Gamma_{26}$ $\chi_{c2} \gamma$	$< 1.3 \%$	90%
$\Gamma_{27}$ $\chi_{c1} \pi^+ \pi^- \pi^0$	$< 2 \times 10^{-3}$	90%
$\Gamma_{28}$ $\chi_{c2} \pi^+ \pi^- \pi^0$	$< 8 \times 10^{-3}$	90%
$\Gamma_{29}$ $h_c(1P) \pi^+ \pi^-$	$< 5 \times 10^{-3}$	90%
$\Gamma_{30}$ $h_c(1P) \pi^0 \pi^0$	$< 2 \times 10^{-3}$	90%
$\Gamma_{31}$ $h_c(1P) \eta$	$< 2 \times 10^{-3}$	90%

$\Gamma_{32}$	$h_c(1P)\pi^0$	$< 4$	$\times 10^{-4}$	90%
$\Gamma_{33}$	$\phi\pi^+\pi^-$	$< 2$	$\times 10^{-3}$	90%
$\Gamma_{34}$	$\gamma\chi_{c1}(3872)$	$< 1.8$	$\times 10^{-3}$	90%
$\Gamma_{35}$	$\gamma\chi_{c0}(3915) \rightarrow \gamma J/\psi\pi^+\pi^-$	$< 1.36$	$\times 10^{-4}$	90%
$\Gamma_{36}$	$\gamma X(3930) \rightarrow \gamma J/\psi\pi^+\pi^-$	$< 1.18$	$\times 10^{-4}$	90%
$\Gamma_{37}$	$\gamma X(3940) \rightarrow \gamma J/\psi\pi^+\pi^-$	$< 1.47$	$\times 10^{-4}$	90%
$\Gamma_{38}$	$\gamma\chi_{c0}(3915) \rightarrow \gamma\gamma J/\psi$	$< 1.26$	$\times 10^{-4}$	90%
$\Gamma_{39}$	$\gamma X(3930) \rightarrow \gamma\gamma J/\psi$	$< 8.8$	$\times 10^{-5}$	90%
$\Gamma_{40}$	$\gamma X(3940) \rightarrow \gamma\gamma J/\psi$	$< 1.79$	$\times 10^{-4}$	90%
$\Gamma_{41}$	$\omega\pi^0$	not seen		
$\Gamma_{42}$	$\omega\eta$	not seen		
$\Gamma_{43}$	$K^+K^-$			
$\Gamma_{44}$	$K_S^0 K^\pm \pi^\mp$			
$\Gamma_{45}$	$p\bar{p}p\bar{p}$	not seen		
$\Gamma_{46}$	$\Lambda\bar{\Lambda}$	$< 1.5$	$\times 10^{-6}$	90%

### $\psi(4160)$ PARTIAL WIDTHS

#### $\Gamma(e^+e^-)$ $\Gamma_1$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.48±0.22</b>	<sup>1</sup> ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.4 to 1.1	<sup>2</sup> MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
0.83±0.08	<sup>3</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
0.84±0.13	<sup>4</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
0.77±0.23	BRANDELIK	78C DASP	$e^+e^-$

<sup>1</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (293 \pm 57)^\circ$ .

<sup>2</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different  $e^+e^-$  partial widths. We quote only the range of values.

<sup>3</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

<sup>4</sup> From a fit to BES (BAI 02c) data.

#### $\Gamma(\mu^+\mu^-)$ $\Gamma_2$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>2.45±1.24±0.94</b>	<sup>1,2</sup> ABLIKIM	20AG BES3	$e^+e^- \rightarrow \mu^+\mu^-$

<sup>1</sup> From a fit to the  $e^+e^- \rightarrow \mu^+\mu^-$  cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming  $\Gamma(\mu^+\mu^-) = \Gamma(e^+e^-)$ .

<sup>2</sup> From solution 1 of 8 with equal fit quality. Other solutions range from  $2.08 \pm 0.99 \pm 0.80$  to  $2.45 \pm 1.24 \pm 0.94$  keV.

### $\psi(4160) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma(\text{total})$

#### $\Gamma(J/\psi\eta') \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{22}\Gamma_1/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.17 ± 0.04	86	<sup>1,2</sup> ABLIKIM	20A	BES3	$e^+ e^- \rightarrow \eta' J/\psi$
1.07 ± 0.09	86	<sup>1,3</sup> ABLIKIM	20A	BES3	$e^+ e^- \rightarrow \eta' J/\psi$

<sup>1</sup> Based on a fit to  $\sigma(e^+ e^- \rightarrow \eta' J/\psi)$  from  $\sqrt{s} = 4.18$  to 4.60 GeV assuming interfering  $\psi(4160)$  and  $\psi(4260)$  contributions. At  $\sqrt{s} = 4.18$  GeV,  $\sigma(e^+ e^- \rightarrow \eta' J/\psi) = 2.4 \pm 0.3 \pm 0.2$  pb.

<sup>2</sup> Solution I of the fit, corresponding to a phase of  $-0.03 \pm 0.44$  rad.

<sup>3</sup> Solution II of the fit, corresponding to a phase of  $2.54 \pm 0.04$  rad.

#### $\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{25}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<2.2	90	<sup>1</sup> HAN	15	BELL	10.58 $e^+ e^- \rightarrow \chi_{c1}\gamma$
------	----	------------------	----	------	---

<sup>1</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .

#### $\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{26}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.1	90	<sup>1</sup> HAN	15	BELL	10.58 $e^+ e^- \rightarrow \chi_{c2}\gamma$
------	----	------------------	----	------	---

<sup>1</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .

#### $\Gamma(K_S^0 K^\pm \pi^\mp) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{44}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.71 ± 0.13 ± 0.12	<sup>1</sup> ABLIKIM	19AE	BES3	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$
0.0095 ± 0.0088 ± 0.0004	<sup>2</sup> ABLIKIM	19AE	BES3	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>1</sup> Solution I of the fit including the  $\psi(4160)$  with mass  $4191 \pm 5$  MeV and width  $70 \pm 10$  MeV from PDG 16 and the  $\psi(4230)$  with mass  $4219.6 \pm 3.3 \pm 5.1$  MeV and width  $56.0 \pm 3.6 \pm 6.9$  MeV from GAO 17.

<sup>2</sup> Solution II of the fit including the  $\psi(4160)$  with mass  $4191 \pm 5$  MeV and width  $70 \pm 10$  MeV from PDG 16 and the  $\psi(4230)$  with mass  $4219.6 \pm 3.3 \pm 5.1$  MeV and width  $56.0 \pm 3.6 \pm 6.9$  MeV from GAO 17.

### $\psi(4160) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma^2(\text{total})$

#### $\Gamma(J/\psi\eta) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{20}/\Gamma \times \Gamma_1/\Gamma$

VALUE (units $10^{-8}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.8 ± 0.9 ± 0.9	<sup>1</sup> WANG	13B	BELL	$e^+ e^- \rightarrow J/\psi\eta\gamma$
12.8 ± 1.7 ± 2.0	<sup>2</sup> WANG	13B	BELL	$e^+ e^- \rightarrow J/\psi\eta\gamma$

<sup>1</sup> Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

<sup>2</sup> Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

## $\psi(4160)$ BRANCHING RATIOS

### $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	<sup>1</sup> AAIJ	13BC LHCB	$B^+ \rightarrow K^+ \mu^+ \mu^-$
<sup>1</sup> AAIJ 13BC report $B(B^+ \rightarrow K^+ \psi(4160)) B(\psi(4160) \rightarrow \mu^+ \mu^-) = (3.5_{-0.8}^{+0.9}) \times 10^{-9}$ .			

### $\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$ $\Gamma_3/\Gamma_9$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.02 ± 0.03 ± 0.02</b>	AUBERT	09M BABR	$e^+ e^- \rightarrow \gamma D^{(*)} \bar{D}^{(*)}$

### $\Gamma(D^0 \bar{D}^0)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^0 \bar{D}^0$
<b>seen</b>	PAKHLOVA 08	BELL	$e^+ e^- \rightarrow D^0 \bar{D}^0 \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^0 \bar{D}^0 \gamma$

### $\Gamma(D^+ D^-)/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^+ D^-$
<b>seen</b>	PAKHLOVA 08	BELL	$e^+ e^- \rightarrow D^+ D^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^+ D^- \gamma$

### $\Gamma(D^*(2007)^0 \bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$ $\Gamma_7/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	AUBERT	09M BABR	$e^+ e^- \rightarrow D^{*0} \bar{D}^0 \gamma$
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0} \bar{D}^0$

### $\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma_{\text{total}}$ $\Gamma_8/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	<sup>1</sup> ZHUKOVA	18 BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$
<b>seen</b>	AUBERT	09M BABR	$e^+ e^- \rightarrow D^{*+} D^- \gamma$
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	PAKHLOVA	07 BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$

<sup>1</sup> Supersedes PAKHLOVA 07.

### $\Gamma(D^* \bar{D} + \text{c.c.})/\Gamma(D^* \bar{D}^*)$ $\Gamma_6/\Gamma_9$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.34 ± 0.14 ± 0.05</b>	AUBERT	09M BABR	$e^+ e^- \rightarrow \gamma D^{(*)} \bar{D}^{(*)}$

### $\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$ $\Gamma_{10}/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	AUBERT	09M BABR	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0}$

$\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	<sup>1</sup> ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
<b>seen</b>	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^{*-}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<b>seen</b>	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
-------------	-------------	------	--

<sup>1</sup> Supersedes PAKHLOVA 07.

$\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.))}/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	PAKHLOVA 08A	BELL	$e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

$\Gamma(D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*)/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D \bar{D}^* \pi$

$\Gamma(D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-))/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$

$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
<b>not seen</b>	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
<b>not seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$

$\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
<b>seen</b>	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^-$

$\Gamma(J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	COAN 06	CLEO	4.12–4.2 $e^+ e^- \rightarrow$ hadrons

$\Gamma(J/\psi \pi^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	COAN 06	CLEO	4.12–4.2 $e^+ e^- \rightarrow$ hadrons

$\Gamma(J/\psi K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN 06	CLEO	4.12–4.2 $e^+ e^- \rightarrow$ hadrons

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$   $\Gamma_{20}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<8	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons
----	----	------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen		<sup>1</sup> ABLIKIM	15L	BES3 $e^+e^- \rightarrow J/\psi\eta$
seen		WANG	13B	BELL $e^+e^- \rightarrow J/\psi\eta\gamma$

<sup>1</sup> An enhancement around 4.2 GeV is observed.

$\Gamma(J/\psi\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{21}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons
----	----	------	----	--

$\Gamma(J/\psi\eta')/\Gamma_{\text{total}}$   $\Gamma_{22}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<5	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons
----	----	------	----	--

$\Gamma(J/\psi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{23}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons
----	----	------	----	--

$\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{24}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<4	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons
----	----	------	----	--

$\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$   $\Gamma_{25}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons
----	----	------	----	--

$\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$   $\Gamma_{26}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<13	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons
-----	----	------	----	--

$\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{27}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<2	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons
----	----	------	----	--

$\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<8	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons
----	----	------	----	--

$\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<5	90	<sup>1</sup> PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$
----	----	---------------------	----	---

<sup>1</sup> At  $\sqrt{s} = 4170$  MeV, PEDLAR 11 measures  $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 15.6 \pm 2.3 \pm 1.9 \pm 3.0$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

$\Gamma(h_c(1P)\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{30}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	<sup>1</sup> PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^0\pi^0$

<sup>1</sup> At  $\sqrt{s} = 4170$  MeV, PEDLAR 11 measures  $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0\pi^0) = 3.0 \pm 3.3 \pm 1.1 \pm 0.6$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

$\Gamma(h_c(1P)\eta)/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<2	90		<sup>1</sup> PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen 41 <sup>2</sup> ABLIKIM 17R BES3  $e^+e^- \rightarrow h_c(1P)\eta$

<sup>1</sup> At  $\sqrt{s} = 4170$  MeV, PEDLAR 11 measures  $\sigma(e^+e^- \rightarrow h_c(1P)\eta) = 4.7 \pm 1.7 \pm 1.0 \pm 0.9$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

<sup>2</sup> An enhancement around 4.2 GeV is observed.

$\Gamma(h_c(1P)\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	<sup>1</sup> PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^0$

<sup>1</sup> At  $\sqrt{s} = 4170$  MeV, PEDLAR 11 measures  $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0) = -0.7 \pm 1.8 \pm 0.7 \pm 0.1$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{33}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN	06	CLEO $4.12\text{--}4.2 e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma\chi_{c1}(3872))/\Gamma_{\text{total}}$   $\Gamma_{34}/\Gamma$

VALUE	CL%	DOCUMENT ID	COMMENT
<1.8 $\times 10^{-3}$	90	<sup>1,2</sup> XIAO	13 $\psi(4160) \rightarrow \gamma J/\psi \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.012 90 <sup>1,3</sup> XIAO 13  $\psi(4160) \rightarrow \gamma J/\psi \pi^+ \pi^-$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

<sup>2</sup> XIAO 13 reports  $[\Gamma(\psi(4160) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] < 0.68 \times 10^{-4}$  which we divide by our best value  $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) = 3.8 \times 10^{-2}$ .

<sup>3</sup> XIAO 13 reports  $[\Gamma(\psi(4160) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \gamma J/\psi)] < 1.05 \times 10^{-4}$  which we divide by our best value  $B(\chi_{c1}(3872) \rightarrow \gamma J/\psi) = 8 \times 10^{-3}$ .

$\Gamma(\gamma\chi_{c0}(3915) \rightarrow \gamma J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$

VALUE	CL%	DOCUMENT ID	COMMENT
<1.36 $\times 10^{-4}$	90	<sup>1</sup> XIAO	13 $\psi(4160) \rightarrow \gamma J/\psi \pi^+ \pi^-$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.



$\Gamma(\gamma X(3930) \rightarrow \gamma J/\psi \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{36} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.18 \times 10^{-4}$       90      <sup>1</sup> XIAO      13       $\psi(4160) \rightarrow \gamma J/\psi \pi^+ \pi^-$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

$\Gamma(\gamma X(3940) \rightarrow \gamma J/\psi \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{37} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.47 \times 10^{-4}$       90      <sup>1</sup> XIAO      13       $\psi(4160) \rightarrow \gamma J/\psi \pi^+ \pi^-$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

$\Gamma(\gamma \chi_{c0}(3915) \rightarrow \gamma \gamma J/\psi) / \Gamma_{\text{total}}$   $\Gamma_{38} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.26 \times 10^{-4}$       90      <sup>1</sup> XIAO      13       $\psi(4160) \rightarrow \gamma \gamma J/\psi$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

$\Gamma(\gamma X(3930) \rightarrow \gamma \gamma J/\psi) / \Gamma_{\text{total}}$   $\Gamma_{39} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<0.88 \times 10^{-4}$       90      <sup>1</sup> XIAO      13       $\psi(4160) \rightarrow \gamma \gamma J/\psi$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

$\Gamma(\gamma X(3940) \rightarrow \gamma \gamma J/\psi) / \Gamma_{\text{total}}$   $\Gamma_{40} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.79 \times 10^{-4}$       90      <sup>1</sup> XIAO      13       $\psi(4160) \rightarrow \gamma \gamma J/\psi$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

$\Gamma(\omega \pi^0) / \Gamma_{\text{total}}$   $\Gamma_{41} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**not seen**      ABLIKIM      22K      BES3       $e^+ e^- \rightarrow \omega \pi^0$

$\Gamma(\omega \eta) / \Gamma_{\text{total}}$   $\Gamma_{42} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**not seen**      ABLIKIM      22K      BES3       $e^+ e^- \rightarrow \omega \eta$

$\Gamma(K^+ K^-) / \Gamma_{\text{total}}$   $\Gamma_{43} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2 \times 10^{-5}$       90      <sup>1</sup> DRUZHININ      15      RVUE       $e^+ e^- \rightarrow \psi(3770)$

<sup>1</sup> DRUZHININ 15 uses BABAR and CLEO data taking into account interference of the processes  $e^+ e^- \rightarrow K^+ K^-$  and  $e^+ e^- \rightarrow K_S^0 K_L^0$ .

$\Gamma(p \bar{p} p \bar{p}) / \Gamma_{\text{total}}$   $\Gamma_{45} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**not seen**      ABLIKIM      21D      BES3      4.0–4.6  $e^+ e^- \rightarrow p \bar{p} p \bar{p}$

$\Gamma(\Lambda \bar{\Lambda}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{46} \Gamma_1 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

$<0.7 \times 10^{-3}$       90      <sup>1</sup> ABLIKIM      21AS      BES3       $e^+ e^- \rightarrow \psi(4160)$

<sup>1</sup> From a measurement of the  $e^+ e^- \rightarrow \Lambda \bar{\Lambda}$  cross section between 3.5 and 4.6 GeV.

## $\psi(4160)$ REFERENCES

ABLIKIM	22K	JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AS	PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21D	PR D103 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20A	PR D101 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20AG	PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AE	PR D99 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	17R	PR D96 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
GAO	17	PR D95 092007	X.Y. Gao, C.P. Shen, C.Z. Yuan	
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
ABLIKIM	15L	PR D91 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DRUZHININ	15	PR D92 054024	V.P. Druzhinin	(NOVO)
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
AAIJ	13BC	PRL 111 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
XIAO	13	PR D87 057501	T. Xiao <i>et al.</i>	(NWES, WAYN)
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
SETH	05A	PR D72 017501	K.K. Seth	
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)