



$I(J^P) = 0(0^-)$   
 $I, J, P$  need confirmation.

Quantum numbers shown are quark-model predictions.

## $B_c^+$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>6274.47 ± 0.27 ± 0.17</b>	<sup>1</sup> AAIJ	20R	LHCb $p\bar{p}$ at 7, 8, 13 TeV
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
6274.28 ± 1.40 ± 0.32	<sup>2</sup> AAIJ	17L	LHCb Repl. by AAIJ 20R
6274.0 ± 1.8 ± 0.4	<sup>3</sup> AAIJ	14AQ	LHCb Repl. by AAIJ 20R
6276.28 ± 1.44 ± 0.36	<sup>4</sup> AAIJ	13AS	LHCb Repl. by AAIJ 20R
6273.7 ± 1.3 ± 1.6	<sup>5</sup> AAIJ	12AV	LHCb Repl. by AAIJ 20R
6275.6 ± 2.9 ± 2.5	<sup>6</sup> AALTONEN	08M	CDF $p\bar{p}$ at 1.96 TeV
6300 ± 14 ± 5	<sup>6</sup> ABAZOV	08T	D0 $p\bar{p}$ at 1.96 TeV
6285.7 ± 5.3 ± 1.2	<sup>6</sup> ABULENCIA	06C	CDF Repl. by AALTONEN 08M
6400 ± 390 ± 130	<sup>7</sup> ABE	98M	CDF $p\bar{p}$ at 1.8 TeV
6320 ± 60	<sup>8</sup> ACKERSTAFF	98O	OPAL $e^+e^- \rightarrow Z$

<sup>1</sup> AAIJ 20R uses the  $B_c^+ \rightarrow J/\psi\pi^+$ ,  $J/\psi\pi^+\pi^-\pi^+$ ,  $J/\psi p\bar{p}\pi^+$ ,  $J/\psi D_s^+$ ,  $J/\psi D^0 K^+$  and  $B_s^0\pi^+$  modes.

<sup>2</sup> Measured using  $B_c^+ \rightarrow J/\psi D^0 K^+$  decays.

<sup>3</sup> Uses  $B_c^+ \rightarrow J/\psi p\bar{p}\pi^+$  decays.

<sup>4</sup> AAIJ 13AS uses the  $B_c^+ \rightarrow J/\psi D_s^+$ .

<sup>5</sup> AAIJ 12AV uses the  $B_c^+ \rightarrow J/\psi\pi^+$  mode and also measures the mass difference  $M(B_c^+) - M(B^+) = 994.6 \pm 1.3 \pm 0.6$  MeV/c<sup>2</sup>.

<sup>6</sup> Measured using a fully reconstructed decay mode of  $B_c \rightarrow J/\psi\pi$ .

<sup>7</sup> ABE 98M observed  $20.4^{+6.2}_{-5.5}$  events in the  $B_c^+ \rightarrow J/\psi(1s)\ell\nu_\ell$  with a significance of > 4.8 standard deviations. The mass value is estimated from  $m(J/\psi(1S)\ell)$ .

<sup>8</sup> ACKERSTAFF 98O observed 2 candidate events in the  $B_c^+ \rightarrow J/\psi(1S)\pi^+$  channel with an estimated background of  $0.63 \pm 0.20$  events.

## $m_{B_c^+} - m_{B_s^0}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>907.75 ± 0.37 ± 0.27</b>	<sup>1</sup> AAIJ	20R	LHCb $p\bar{p}$ at 7, 8, 13 TeV
<sup>1</sup> AAIJ 20R uses the $B_c^+ \rightarrow J/\psi\pi^+$ , $J/\psi\pi^+\pi^-\pi^+$ , $J/\psi p\bar{p}\pi^+$ , $J/\psi D_s^+$ , $J/\psi D^0 K^+$ and $B_s^0\pi^+$ modes.			

## $B_c^+$ MEAN LIFE

VALUE (10 <sup>-12</sup> s)	DOCUMENT ID	TECN	COMMENT
<b>0.510 ± 0.009 OUR EVALUATION</b>	(Produced by HFLAV)		
<b>0.510 ± 0.009 OUR AVERAGE</b>			

0.541 ± 0.026 ± 0.014	<sup>1</sup> SIRUNYAN	18BY CMS	$p\bar{p}$ at 8 TeV
0.5134 ± 0.0110 ± 0.0057	<sup>2,3</sup> AAIJ	15G LHCb	$p\bar{p}$ at 7, 8 TeV
0.509 ± 0.008 ± 0.012	<sup>4</sup> AAIJ	14G LHCb	$p\bar{p}$ at 8 TeV

0.452	$\pm 0.048$	$\pm 0.027$	<sup>3</sup> AALTONEN	13	CDF	$p\bar{p}$ at 1.96 TeV
0.448	$+0.038$	$-0.036$	$\pm 0.032$	<sup>5</sup> ABAZOV	09H D0	$p\bar{p}$ at 1.96 TeV
0.463	$+0.073$	$-0.065$	$\pm 0.036$	<sup>5</sup> ABULENCIA	060 CDF	$p\bar{p}$ at 1.96 TeV
0.46	$+0.18$	$-0.16$	$\pm 0.03$	<sup>5</sup> ABE	98M CDF	$p\bar{p}$ 1.8 TeV

<sup>1</sup> The lifetime is measured using the decays  $B_c^+ \rightarrow J/\psi\pi^+$  and  $B^+ \rightarrow J/\psi K^+$ .

<sup>2</sup> Also measures the width difference  $\Delta\Gamma = \Gamma_{B_c^+} - \Gamma_{B^+} = 4.46 \pm 0.14 \pm 0.07 \text{ mm}^{-1} \text{ c.}$

<sup>3</sup> Uses fully reconstructed  $B_c^+ \rightarrow J/\psi\pi^+$  decays.

<sup>4</sup> Measured using  $B_c^+ \rightarrow J/\psi\mu^+\nu_\mu X$  decays.

<sup>5</sup> The lifetime is measured from the  $J/\psi e$  decay vertices.

## $B_c^+$ DECAY MODES $\times \mathbf{B}(\bar{b} \rightarrow B_c)$

The following quantities are not pure branching ratios; rather the fractions

$\Gamma_i/\Gamma \times \mathbf{B}(\bar{b} \rightarrow B_c)$ .  $B_c^-$  modes are charge conjugates of the modes below.

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $J/\psi(1S)\ell^+\nu_\ell$ anything		seen
$\Gamma_2$ $J/\psi(1S)\mu^+\nu_\mu$		seen
$\Gamma_3$ $J/\psi(1S)\tau^+\nu_\tau$		seen
$\Gamma_4$ $J/\psi(1S)\pi^+$		seen
$\Gamma_5$ $J/\psi(1S)K^+$		seen
$\Gamma_6$ $J/\psi(1S)\pi^+\pi^+\pi^-$		seen
$\Gamma_7$ $J/\psi(1S)K^+\pi^-\pi^+$		
$\Gamma_8$ $J/\psi(1S)K^+K^-K^+$		
$\Gamma_9$ $J/\psi(1S)a_1(1260)$		not seen
$\Gamma_{10}$ $J/\psi(1S)K^+K^-\pi^+$		seen
$\Gamma_{11}$ $J/\psi(1S)\pi^+\pi^+\pi^-\pi^-$		seen
$\Gamma_{12}$ $\psi(2S)\pi^+$		seen
$\Gamma_{13}$ $\psi(2S)\pi^+\pi^-\pi^+$		
$\Gamma_{14}$ $\psi(2S)K^+K^-\pi^+$		
$\Gamma_{15}$ $J/\psi(1S)D^0K^+$		seen
$\Gamma_{16}$ $J/\psi(1S)D^*(2007)^0K^+$		seen
$\Gamma_{17}$ $J/\psi(1S)D^*(2010)^+K^{*0}$		seen
$\Gamma_{18}$ $J/\psi(1S)D^+K^{*0}$		seen
$\Gamma_{19}$ $J/\psi(1S)D_s^+$		seen
$\Gamma_{20}$ $J/\psi(1S)D_s^{*+}$		seen
$\Gamma_{21}$ $J/\psi(1S)p\bar{p}\pi^+$		seen
$\Gamma_{22}$ $\chi_{c0}\pi^+$	$(2.4^{+0.9}_{-0.8}) \times 10^{-5}$	
$\Gamma_{23}$ $p\bar{p}\pi^+$		not seen
$\Gamma_{24}$ $D^0K^+$		seen

$\Gamma_{25}$	$D^0 \pi^+$	not seen		
$\Gamma_{26}$	$D^{*0} \pi^+$	not seen		
$\Gamma_{27}$	$D^{*0} K^+$	not seen		
$\Gamma_{28}$	$D_s^+ \bar{D}^0$	$< 7.2 \times 10^{-4}$	90%	
$\Gamma_{29}$	$D_s^+ D^0$	$< 3.0 \times 10^{-4}$	90%	
$\Gamma_{30}$	$D^+ \bar{D}^0$	$< 1.9 \times 10^{-4}$	90%	
$\Gamma_{31}$	$D^+ D^0$	$< 1.4 \times 10^{-4}$	90%	
$\Gamma_{32}$	$D_s^{*+} \bar{D}^0$	$< 5.3 \times 10^{-4}$	90%	
$\Gamma_{33}$	$D_s^+ \bar{D}^*(2007)^0$	$< 4.6 \times 10^{-4}$	90%	
$\Gamma_{34}$	$D_s^{*+} D^0$	$< 9 \times 10^{-4}$	90%	
$\Gamma_{35}$	$D_s^+ D^*(2007)^0$	$< 6.6 \times 10^{-4}$	90%	
$\Gamma_{36}$	$D^*(2010)^+ \bar{D}^0$	$< 3.8 \times 10^{-4}$	90%	
$\Gamma_{37}$	$D^*(2010)^+ \bar{D}^0, D^{*+} \rightarrow D^+ \pi^0 / \gamma$	not seen		
$\Gamma_{38}$	$D^+ \bar{D}^*(2007)^0$	$< 6.5 \times 10^{-4}$	90%	
$\Gamma_{39}$	$D^*(2007)^+ D^0$	$< 2.0 \times 10^{-4}$	90%	
$\Gamma_{40}$	$D^*(2010)^+ D^0, D^{*+} \rightarrow D^+ \pi^0 / \gamma$	not seen		
$\Gamma_{41}$	$D^+ D^*(2007)^0$	$< 3.7 \times 10^{-4}$	90%	
$\Gamma_{42}$	$D_s^{*+} \bar{D}^*(2007)^0$	$< 1.3 \times 10^{-3}$	90%	
$\Gamma_{43}$	$D_s^{*+} D^*(2007)^0$	$< 1.3 \times 10^{-3}$	90%	
$\Gamma_{44}$	$D^*(2010)^+ \bar{D}^*(2007)^0$	$< 1.0 \times 10^{-3}$	90%	
$\Gamma_{45}$	$D^*(2010)^+ D^*(2007)^0$	$< 7.7 \times 10^{-4}$	90%	
$\Gamma_{46}$	$D^+ K^{*0}$	not seen		
$\Gamma_{47}$	$D^+ \bar{K}^{*0}$	not seen		
$\Gamma_{48}$	$D_s^+ K^{*0}$	not seen		
$\Gamma_{49}$	$D_s^+ \bar{K}^{*0}$	not seen		
$\Gamma_{50}$	$D_s^+ \phi$	not seen		
$\Gamma_{51}$	$K^+ K^0$	not seen		
$\Gamma_{52}$	$B_s^0 \pi^+ / B(\bar{b} \rightarrow B_s)$	seen		
$\Gamma_{53}$	$B_s^0 \pi^+$			

 **$B_c^+$  BRANCHING RATIOS**

$$\Gamma(J/\psi(1S)\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}} \times \mathbf{B}(\bar{b} \rightarrow B_c)$$

$$\Gamma_1/\Gamma \times \mathbf{B}$$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>8.2 \pm 1.3</math> OUR AVERAGE</b>		Error includes scale factor of 1.4.		
8.8 $\pm 1.0 \pm 0.2$	1,2	AALTONEN	16A CDF	$p\bar{p}$ at 1.96 TeV
5.2 $\pm 2.4$ 5.2 $\pm 2.1$	3	ABE	98M CDF	$p\bar{p}$ 1.8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	<sup>4</sup> ACKERSTAFF	980 OPAL	$e^+ e^- \rightarrow Z$
<19	90	<sup>5</sup> ABREU	97E DLPH	$e^+ e^- \rightarrow Z$
<12	90	<sup>6</sup> BARATE	97H ALEP	$e^+ e^- \rightarrow Z$

- <sup>1</sup> AALTONEN 16A reports  $[\Gamma(B_c^+ \rightarrow J/\psi(1S)\ell^+\nu_\ell \text{anything})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.211 \pm 0.012^{+0.021}_{-0.020}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = (40.8 \pm 0.7) \times 10^{-2}$ ,  $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.
- <sup>2</sup> AALTONEN 16A also measures the cross-section  $\sigma(B_c) \times B(B_c \rightarrow J/\psi\mu\nu_\mu) = 0.60 \pm 0.09 \text{ nb}$  and estimates the total cross-section  $\sigma(B_c)$  to be in the range  $25 \pm 4$  to  $52 \pm 8 \text{ nb}$  for  $p_T(B_c) > 6 \text{ GeV}/c$  and  $|y(B_c)| < 1$ .
- <sup>3</sup> ABE 98M result is derived from the measurement of  $[\sigma(B_c) \times B(B_c \rightarrow J/\psi(1S)\ell\nu_\ell)] / [\sigma(B^+) \times B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.132^{+0.041}_{-0.037} (\text{stat}) \pm 0.031 (\text{sys})^{+0.032}_{-0.020} (\text{lifetime})$  by using PDG 98 values of  $B(b \rightarrow B^+)$  and  $B(B^+ \rightarrow J/\psi(1S)K^+)$ .
- <sup>4</sup> ACKERSTAFF 98O reports  $B(Z \rightarrow B_c X) / B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S)\ell\nu_\ell) < 6.95 \times 10^{-5}$  at 90%CL. We rescale to our PDG 98 values of  $B(Z \rightarrow b\bar{b})$ .
- <sup>5</sup> ABREU 97E value listed is for an assumed  $\tau_{B_c} = 0.4 \text{ ps}$  and improves to  $1.6 \times 10^{-4}$  for  $\tau_{B_c} = 1.4 \text{ ps}$ .
- <sup>6</sup> BARATE 97H reports  $B(Z \rightarrow B_c X) / B(Z \rightarrow qq) \cdot B(B_c \rightarrow J/\psi(1S)\ell\nu_\ell) < 5.2 \times 10^{-5}$  at 90%CL. We rescale to our PDG 96 values of  $B(Z \rightarrow b\bar{b})$ . A  $B_c^+ \rightarrow J/\psi(1S)\mu^+\nu_\mu$  candidate event is found, compared to all the known background sources  $2 \times 10^{-3}$ , which gives  $m_{B_c} = 5.96^{+0.25}_{-0.19} \text{ GeV}$  and  $\tau_{B_c} = 1.77 \pm 0.17 \text{ ps}$ .

$\Gamma(J/\psi(1S)\tau^+\nu_\tau)/\Gamma(J/\psi(1S)\mu^+\nu_\mu)$	$\Gamma_3/\Gamma_2$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.71±0.17±0.18</b>	<sup>1</sup> AAIJ	18C	$p p$ at 7, 8 TeV

<sup>1</sup> AAIJ 18C uses  $\tau^+ \rightarrow \mu^+\nu_\mu\bar{\nu}_\tau$  mode to obtain the ratio value.

$\Gamma(J/\psi(1S)\pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$	$\Gamma_4/\Gamma \times B$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
seen		<sup>1</sup> AABOUD	21 ATLAS	$p p$ at 8 TeV
seen		<sup>2</sup> AAIJ	15M LHCb	$p p$ at 8 TeV
seen		<sup>3</sup> KHACHATRYAN...15AA CMS		$p p$ at 7 TeV
seen		AALTONEN	13 CDF	$p\bar{p}$ at 1.96 TeV
seen		<sup>4</sup> AAIJ	12AV LHCb	$p p$ at 7 TeV
seen		AALTONEN	08M CDF	$p\bar{p}$ at 1.96 TeV
seen		ABAZOV	08T D0	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.4 \times 10^{-4}$	90	<sup>5</sup> ACKERSTAFF 98O OPAL $e^+e^- \rightarrow Z$
$<3.4 \times 10^{-4}$	90	<sup>6</sup> ABREU 97E DLPH $e^+e^- \rightarrow Z$
$<8.2 \times 10^{-5}$	90	<sup>7</sup> BARATE 97H ALEP $e^+e^- \rightarrow Z$
$<2.0 \times 10^{-5}$	95	<sup>8</sup> ABE 96R CDF $p\bar{p}$ 1.8 TeV

<sup>1</sup> AABOUD 21 reports a measurement of  $B(B_c^+ \rightarrow J/\psi\pi^+) / B(B^+ \rightarrow J/\psi K^+) \cdot f_c/f_u = (0.34 \pm 0.04^{+0.06}_{-0.02} \pm 0.01) \%$ , at  $p_T > 13 \text{ GeV}$  and  $|y| < 2.3$ .

<sup>2</sup> AAIJ 15M reports a measurement of  $B(B_c^+ \rightarrow J/\psi\pi^+) / B(B^+ \rightarrow J/\psi K^+) \cdot f_c/f_u = (0.683 \pm 0.018 \pm 0.009) \%$  at  $p_T(B) < 20 \text{ GeV}$  and  $2.0 < y(B) < 4.5$ .

<sup>3</sup> KHACHATRYAN 15AA reports a measurement of  $B(B_c^+ \rightarrow J/\psi\pi^+) / B(B^+ \rightarrow J/\psi K^+) \cdot f_c/f_u = (0.48 \pm 0.05 \pm 0.03 \pm 0.05) \%$ , at  $p_T > 15 \text{ GeV}$  and  $|\eta(B)| < 1.6$ .

- <sup>4</sup> AAIJ 12AV reports a measurement of  $B(B_c^+ \rightarrow J/\psi\pi^+)/B(B^+ \rightarrow J/\psi K^+) f_c/f_u = (0.68 \pm 0.10 \pm 0.03 \pm 0.05)\%$  at  $p_T(B) > 4$  GeV and  $2.5 < \eta(B) < 4.5$ .
- <sup>5</sup> ACKERSTAFF 980 reports  $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S)\pi^+) < 1.06 \times 10^{-4}$  at 90%CL. We rescale to our PDG 98 values of  $B(Z \rightarrow b\bar{b})$ .
- <sup>6</sup> ABREU 97E value listed is for an assumed  $\tau_{B_c} = 0.4$  ps and improves to  $2.7 \times 10^{-4}$  for  $\tau_{B_c} = 1.4$  ps.
- <sup>7</sup> BARATE 97H reports  $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \cdot B(B_c \rightarrow J/\psi(1S)\pi) < 3.6 \times 10^{-5}$  at 90%CL. We rescale to our PDG 96 values of  $B(Z \rightarrow b\bar{b})$ .
- <sup>8</sup> ABE 96R reports  $B(b \rightarrow B_c X)/B(b \rightarrow B^+ X) \cdot B(B_c^+ \rightarrow J/\psi(1S)\pi^+)/B(B^+ \rightarrow J/\psi(1S)K^+) < 0.053$  at 95%CL for  $\tau_{B_c} = 0.8$  ps. It changes from 0.15 to 0.04 for  $0.17 \text{ ps} < \tau_{B_c} < 1.6$  ps. We rescale to our PDG 96 values of  $B(b \rightarrow B^+) = 0.378 \pm 0.022$  and  $B(B^+ \rightarrow J/\psi(1S)K^+) = 0.00101 \pm 0.00014$ .

### $\Gamma(J/\psi(1S)\pi^+)/\Gamma(J/\psi(1S)\mu^+\nu_\mu)$ $\Gamma_4/\Gamma_2$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.69 ± 0.28 ± 0.46</b>	<sup>1</sup> AAIJ	14W LHCb	$p p$ at 7 TeV

<sup>1</sup> AAIJ 14W reports also a measurement  $B(B_c^+ \rightarrow J/\psi\pi^+)/B(B_c^+ \rightarrow J/\psi\mu^+\nu_\mu) = 0.271 \pm 0.016 \pm 0.016$  in the region  $m_{J/\psi\mu^+} > 5.3$  GeV.

### $\Gamma(J/\psi(1S)K^+)/\Gamma(J/\psi(1S)\pi^+)$ $\Gamma_5/\Gamma_4$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.079 ± 0.007 ± 0.003</b>		AAIJ	16AF LHCb	$p p$ at 7, 8 TeV
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
0.069 ± 0.019 ± 0.005	50	AAIJ	13BY LHCb	Repl. by AAIJ 16AF

### $\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_6/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
seen		AAIJ	12Y LHCb	$p p$ at 7 TeV
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
$< 5.7 \times 10^{-4}$	90	<sup>1</sup> ABREU	97E DLPH	$e^+e^- \rightarrow Z$
<sup>1</sup> ABREU 97E value listed is independent of $0.4 \text{ ps} < \tau_{B_c} < 1.4$ ps.				

### $\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+)$ $\Gamma_6/\Gamma_4$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>2.4 ± 0.4 OUR AVERAGE</b>			
$2.55 \pm 0.80 \pm 0.33^{+0.04}_{-0.01}$	KHACHATRY...15AA CMS	$p p$ at 7 TeV	
$2.41 \pm 0.30 \pm 0.33$	AAIJ	12Y LHCb	$p p$ at 7 TeV

### $\Gamma(J/\psi(1S)a_1(1260))/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_9/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.2 \times 10^{-3}$	90	<sup>1</sup> ACKERSTAFF 980	OPAL	$e^+e^- \rightarrow Z$

<sup>1</sup> ACKERSTAFF 980 reports  $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S)a_1(1260)) < 5.29 \times 10^{-4}$  at 90%CL. We rescale to our PDG 98 values of  $B(Z \rightarrow b\bar{b})$ .

### $\Gamma(J/\psi(1S)K^+K^-\pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{10}/\Gamma \times B$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	<sup>1</sup> AAIJ	13CA LHCb	$p p$ at 7, 8 TeV

<sup>1</sup> A signal yield of  $78 \pm 14$  decays is reported with a significance of 6.2 standard deviations using an integrated luminosity of  $3 \text{ fb}^{-1}$  data.

### $\Gamma(J/\psi(1S)K^+K^-\pi^+)/\Gamma(J/\psi(1S)\pi^+)$ $\Gamma_{10}/\Gamma_4$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.53±0.10±0.05</b>	<sup>1</sup> AAIJ	13CA LHCb	$p p$ at 7, 8 TeV

<sup>1</sup> A signal yield of  $78 \pm 14$  decays is reported with a significance of 6.2 standard deviations using an integrated luminosity of  $3 \text{ fb}^{-1}$  data.

### $\Gamma(J/\psi(1S)\pi^+\pi^+\pi^+\pi^-\pi^-)/\Gamma(J/\psi(1S)\pi^+)$ $\Gamma_{11}/\Gamma_4$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.74±0.44±0.24</b>	<sup>1</sup> AAIJ	14P LHCb	$p p$ at 7, 8 TeV

<sup>1</sup> A signal yield of  $32 \pm 8$  decays is reported with a significance of 4.5 standard deviations.

### $\Gamma(\psi(2S)\pi^+)/\Gamma(J/\psi(1S)\pi^+)$ $\Gamma_{12}/\Gamma_4$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.268±0.032±0.007±0.006</b>	<sup>1</sup> AAIJ	15AY LHCb	$p p$ at 7, 8 TeV
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
$0.250 \pm 0.068 \pm 0.014 \pm 0.006$	<sup>1</sup> AAIJ	13AM LHCb	Repl. by AAIJ 15AY

<sup>1</sup> The last uncertainty is due to the uncertainty of the  $B(\psi(2S) \rightarrow \mu^+ \mu^-)/B(J/\psi \rightarrow \mu^+ \mu^-)$  ratio measurement.

### $\Gamma(\psi(2S)\pi^+)/\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)$ $\Gamma_{12}/\Gamma_6$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>(3.5±0.6±0.2) × 10<sup>-2</sup></b>	AAIJ	22P LHCb	$p p$ at 7, 8, 13 TeV

### $\Gamma(\psi(2S)K^+K^-\pi^+)/\Gamma(J/\psi(1S)K^+K^-\pi^+)$ $\Gamma_{14}/\Gamma_{10}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>(3.7±1.2±0.1) × 10<sup>-2</sup></b>	AAIJ	22P LHCb	$p p$ at 7, 8, 13 TeV

### $\Gamma(J/\psi(1S)K^+K^-K^+)/\Gamma(J/\psi(1S)K^+K^-\pi^+)$ $\Gamma_8/\Gamma_{10}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>(7.0±1.8±0.2) × 10<sup>-2</sup></b>	AAIJ	22P LHCb	$p p$ at 7, 8, 13 TeV

### $\Gamma(J/\psi(1S)K^+\pi^-\pi^+)/\Gamma(J/\psi(1S)K^+\pi^-\pi^+)$ $\Gamma_7/\Gamma_{10}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.35±0.06±0.01</b>	AAIJ	22P LHCb	$p p$ at 7, 8, 13 TeV

### $\Gamma(J/\psi(1S)K^+K^-\pi^+)/\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)$ $\Gamma_{10}/\Gamma_6$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.185±0.013±0.006</b>	AAIJ	22P LHCb	$p p$ at 7, 8, 13 TeV

### $\Gamma(\psi(2S)\pi^+\pi^-\pi^+)/\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)$ $\Gamma_{13}/\Gamma_6$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>(1.9±0.4±0.1) × 10<sup>-2</sup></b>	AAIJ	22P LHCb	$p p$ at 7, 8, 13 TeV

### $\Gamma(J/\psi(1S)D^0K^+)/\Gamma(J/\psi(1S)\pi^+)$ $\Gamma_{15}/\Gamma_4$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.432±0.136±0.028</b>	AAIJ	17L LHCb	$p p$ at 7, 8 TeV

$$\Gamma(J/\psi(1S)D^*(2007)^0 K^+)/\Gamma(J/\psi(1S)D^0 K^+) \quad \Gamma_{16}/\Gamma_{15}$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.1±1.8±0.4</b>	AAIJ	17L	LHCb $p p$ at 7, 8 TeV

$$\Gamma(J/\psi(1S)D^*(2010)^+ K^{*0})/\Gamma(J/\psi(1S)D^0 K^+) \quad \Gamma_{17}/\Gamma_{15}$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.10±1.08±0.34</b>	AAIJ	17L	LHCb $p p$ at 7, 8 TeV

$$\Gamma(J/\psi(1S)D^+ K^{*0})/\Gamma(J/\psi(1S)D^0 K^+) \quad \Gamma_{18}/\Gamma_{15}$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.63±0.39±0.08</b>	AAIJ	17L	LHCb $p p$ at 7, 8 TeV

$$\Gamma(J/\psi(1S)D_s^+)/\Gamma(J/\psi(1S)\pi^+) \quad \Gamma_{19}/\Gamma_4$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.8 ±0.4 OUR AVERAGE</b>			
2.76±0.33±0.33	<sup>1</sup> AAD	220	ATLS $p p$ at 13 TeV
2.90±0.57±0.24	AAIJ	13AS	LHCb $p p$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3.8 ±1.1 ±0.4	AAD	16H	ATLS $p p$ at 7, 8 TeV

<sup>1</sup> Supersedes the measurement of AAD 16H.

$$\Gamma(J/\psi(1S)D_s^{*+})/\Gamma(J/\psi(1S)\pi^+) \quad \Gamma_{20}/\Gamma_4$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.33±0.61±0.74</b>	AAD	220	ATLS $p p$ at 13 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			

10.4 ±3.1 ±1.6 AAD 16H ATLS Repl. by AAD 220

$$\Gamma(J/\psi(1S)D_s^{*+})/\Gamma(J/\psi(1S)D_s^+) \quad \Gamma_{20}/\Gamma_{19}$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.00±0.23 OUR AVERAGE</b>			
1.93±0.24±0.09	<sup>1</sup> AAD	220	ATLS $p p$ at 13 TeV
2.37±0.56±0.10	AAIJ	13AS	LHCb $p p$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.8 +1.2 -0.8 ±0.3	AAD	16H	ATLS $p p$ at 7, 8 TeV

<sup>1</sup> Supersedes the measurement of AAD 16H.

$$\Gamma(J/\psi(1S)p\bar{p}\pi^+)/\Gamma(J/\psi(1S)\pi^+) \quad \Gamma_{21}/\Gamma_4$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.143 +0.041 -0.036</b>	AAIJ	14AQ	LHCb $p p$ at 7, 8 TeV

$$\Gamma(\chi_{c0}\pi^+)/\Gamma_{\text{total}} \quad \Gamma_{22}/\Gamma$$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>24.0 +8.6 -7.6 ±0.4</b>	<sup>1,2</sup> AAIJ	16AT	LHCb $p p$ at 7 and 8 TeV

<sup>1</sup> AAIJ 16AT reports  $[\Gamma(B_c^+ \rightarrow \chi_{c0}\pi^+)/\Gamma_{\text{total}}] \times [\Gamma(\bar{b} \rightarrow B^+)/\Gamma_{\text{total}}] = (9.8 +3.4 -3.0 \pm 0.8) \times 10^{-6}$  which we divide by our best value  $\Gamma(\bar{b} \rightarrow B^+)/\Gamma_{\text{total}} = 0.408 \pm 0.007$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> The significance of the observed signal is 4.0 standard deviations.

$\Gamma(p\bar{p}\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{23}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	<sup>1</sup> AAIJ	16K	LHCb $p\bar{p}$ at 7, 8 TeV
<sup>1</sup> Measures the ratio $(f_c/f_u) \times B(B_c^+ \rightarrow p\bar{p}\pi^+) < 3.6 \times 10^{-8}$ at 95% CL, in the region $m(p\bar{p}) < 2.85 \text{ GeV}/c^2$ , where $f_c$ ( $f_u$ ) represents the fragmentation fraction of the $b$ -quark into the $B_c^+$ ( $B_u^+$ ) meson.			

 $\Gamma(D^0 K^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$   $\Gamma_{24}/\Gamma \times B$ 

VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>3.8^{+1.2}_{-1.0} \pm 0.1</math></b>	<sup>1</sup> AAIJ	17AG	LHCb $p\bar{p}$ at 7, 8 TeV

<sup>1</sup> AAIJ 17AG reports  $[\Gamma(B_c^+ \rightarrow D^0 K^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] = (9.3^{+2.8}_{-2.5} \pm 0.6) \times 10^{-7}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = (40.8 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(D^0 \pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$   $\Gamma_{25}/\Gamma \times B$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;1.6 \times 10^{-7}</math></b>	95	<sup>1</sup> AAIJ	17AG	LHCb $p\bar{p}$ at 7, 8 TeV

<sup>1</sup> AAIJ 17AG reports  $[\Gamma(B_c^+ \rightarrow D^0 \pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 3.9 \times 10^{-7}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

 $\Gamma(D^{*0} \pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$   $\Gamma_{26}/\Gamma \times B$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;4 \times 10^{-7}</math></b>	95	<sup>1</sup> AAIJ	17AG	LHCb $p\bar{p}$ at 7, 8 TeV

<sup>1</sup> AAIJ 17AG reports  $[\Gamma(B_c^+ \rightarrow D^{*0} \pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 1.1 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

 $\Gamma(D^{*0} K^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$   $\Gamma_{27}/\Gamma \times B$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;4 \times 10^{-7}</math></b>	95	<sup>1</sup> AAIJ	17AG	LHCb $p\bar{p}$ at 7, 8 TeV

<sup>1</sup> AAIJ 17AG reports  $[\Gamma(B_c^+ \rightarrow D^{*0} K^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 1.1 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

 $\Gamma(D_s^+ \bar{D}^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$   $\Gamma_{28}/\Gamma \times B$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;1.4 \times 10^{-7}</math></b>	90	<sup>1</sup> AAIJ	18P	LHCb $p\bar{p}$ at 7, 8 TeV

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D_s^+ \bar{D}^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 0.9 \times 10^{-3}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

 $\Gamma(D_s^+ D^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$   $\Gamma_{29}/\Gamma \times B$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;6 \times 10^{-8}</math></b>	90	<sup>1</sup> AAIJ	18P	LHCb $p\bar{p}$ at 7, 8 TeV

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D_s^+ D^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 3.7 \times 10^{-4}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

### $\Gamma(D^+ \bar{D}^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{30}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-6}$	90	<sup>1</sup> AAIJ	18P LHCb	$p p$ at 7, 8 TeV
<sup>1</sup> AAIJ 18P reports $[\Gamma(B_c^+ \rightarrow D^+ \bar{D}^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 1.9 \times 10^{-2}$ which we multiply by our best values $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ , $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .				

### $\Gamma(D^+ D^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{31}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<1.9 \times 10^{-6}$	90	<sup>1</sup> AAIJ	18P LHCb	Repl. by AAIJ 21AF
<sup>1</sup> AAIJ 18P reports $[\Gamma(B_c^+ \rightarrow D^+ D^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 1.2 \times 10^{-2}$ which we multiply by our best values $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ , $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .				

### $[\Gamma(D_s^{*+} \bar{D}^0) + \Gamma(D_s^+ \bar{D}^*(2007)^0)]/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $(\Gamma_{32}+\Gamma_{33})/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<4 \times 10^{-7}$	90	<sup>1</sup> AAIJ	18P LHCb	Repl. by AAIJ 21AF
<sup>1</sup> AAIJ 18P reports $[\Gamma(B_c^+ \rightarrow D_s^{*+} \bar{D}^0) + \Gamma(B_c^+ \rightarrow D_s^+ \bar{D}^*(2007)^0)]/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 2.8 \times 10^{-3}$ which we multiply by our best values $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ , $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .				

### $[\Gamma(D_s^{*+} D^0) + \Gamma(D_s^+ D^*(2007)^0)]/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $(\Gamma_{34}+\Gamma_{35})/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<5 \times 10^{-7}$	90	<sup>1</sup> AAIJ	18P LHCb	Repl. by AAIJ 21AF
<sup>1</sup> AAIJ 18P reports $[\Gamma(B_c^+ \rightarrow D_s^{*+} D^0) + \Gamma(B_c^+ \rightarrow D_s^+ D^*(2007)^0)]/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 3.0 \times 10^{-3}$ which we multiply by our best values $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ , $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .				

### $\Gamma(D^*(2010)^+ \bar{D}^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{36}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.2 \times 10^{-3}$	90	<sup>1</sup> BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$
<sup>1</sup> BARATE 98Q reports $B(Z \rightarrow B_c X) \times B(B_c \rightarrow D^*(2010)^+ \bar{D}^0) < 1.9 \times 10^{-3}$ at 90%CL. We rescale to our PDG 98 values of $B(Z \rightarrow b\bar{b})$ .				

### $[\Gamma(D^*(2010)^+ \bar{D}^0, D^{*+} \rightarrow D^+ \pi^0/\gamma) + \Gamma(D^+ \bar{D}^*(2007)^0)]/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $(\Gamma_{37}+\Gamma_{38})/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<9 \times 10^{-6}$	90	<sup>1</sup> AAIJ	18P LHCb	Repl. by AAIJ 21AF

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D^*(2010)^+ \bar{D}^0, D^{*+} \rightarrow D^+ \pi^0/\gamma) + \Gamma(B_c^+ \rightarrow D^+ \bar{D}^*(2007)^0)] / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 5.5 \times 10^{-2}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

### $[\Gamma(D^*(2010)^+ D^0, D^{*+} \rightarrow D^+ \pi^0/\gamma) + \Gamma(D^+ D^*(2007)^0)] / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{40}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.4 \times 10^{-6}$  90 <sup>1</sup> AAIJ 18P LHCb Repl. by AAIJ 21AF

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D^*(2010)^+ D^0, D^{*+} \rightarrow D^+ \pi^0/\gamma) + \Gamma(B_c^+ \rightarrow D^+ D^*(2007)^0)] / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 2.2 \times 10^{-2}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

### $\Gamma(D_s^{*+} \bar{D}^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{42}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.7 \times 10^{-6}$  90 <sup>1</sup> AAIJ 18P LHCb Repl. by AAIJ 21AF

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D_s^{*+} \bar{D}^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 1.1 \times 10^{-2}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

### $\Gamma(D_s^{*+} D^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{43}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.1 \times 10^{-6}$  90 <sup>1</sup> AAIJ 18P LHCb Repl. by AAIJ 21AF

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D_s^{*+} D^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 2.0 \times 10^{-2}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

### $\Gamma(D^*(2010)^+ \bar{D}^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{44}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.0 \times 10^{-4}$  90 <sup>1</sup> AAIJ 18P LHCb Repl. by AAIJ 21AF

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D^*(2010)^+ \bar{D}^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 6.5 \times 10^{-1}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

### $\Gamma(D^*(2010)^+ D^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{45}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2.0 \times 10^{-5}$  90 <sup>1</sup> AAIJ 18P LHCb Repl. by AAIJ 21AF

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D^*(2010)^+ D^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+) / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 1.3 \times 10^{-1}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

### $\Gamma(D^+ K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{46}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-7}$	90	<sup>1</sup> AAIJ	13R LHCb	$p p$ at 7 TeV

<sup>1</sup> AAIJ 13R reports  $[\Gamma(B_c^+ \rightarrow D^+ K^{*0}) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+) < 0.5 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

### $\Gamma(D^+ \bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{47}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.6 \times 10^{-7}$	90	<sup>1</sup> AAIJ	13R LHCb	$p p$ at 7 TeV

<sup>1</sup> AAIJ 13R reports  $[\Gamma(B_c^+ \rightarrow D^+ \bar{K}^{*0}) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+) < 0.4 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

### $\Gamma(D_s^+ K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{48}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.9 \times 10^{-7}$	90	<sup>1</sup> AAIJ	13R LHCb	$p p$ at 7 TeV

<sup>1</sup> AAIJ 13R reports  $[\Gamma(B_c^+ \rightarrow D_s^+ K^{*0}) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+) < 0.7 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

### $\Gamma(D_s^+ \bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{49}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4 \times 10^{-7}$	90	<sup>1</sup> AAIJ	13R LHCb	$p p$ at 7 TeV

<sup>1</sup> AAIJ 13R reports  $[\Gamma(B_c^+ \rightarrow D_s^+ \bar{K}^{*0}) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+) < 1.1 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

### $\Gamma(D_s^+ \phi)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{50}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.3 \times 10^{-7}$	90	<sup>1</sup> AAIJ	13R LHCb	$p p$ at 7 TeV

<sup>1</sup> AAIJ 13R reports  $[\Gamma(B_c^+ \rightarrow D_s^+ \phi) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+) < 0.8 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

### $\Gamma(K^+ K^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{51}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.6 \times 10^{-7}$	90	<sup>1</sup> AAIJ	13BS LHCb	$p p$ at 7 TeV

<sup>1</sup> Derived from  $\Gamma(K^+ K^0)/\Gamma \times B(\bar{b} \rightarrow B_c) / (B(B^+ \rightarrow K^0 \pi^+) B(\bar{b} \rightarrow B^+) < 5.8\% \text{ at } 90\% \text{ CL using normalization mode } B(B^+ \rightarrow K^0 \pi^+) = (23.97 \pm 0.53 \pm 0.71) \times 10^{-6}$  and assuming a  $B$  production ratio  $f(\bar{b} \rightarrow B_u^+) = 0.33$ .

### $\Gamma(B_s^0 \pi^+ / B(\bar{b} \rightarrow B_s))/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{52}/\Gamma \times B$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
$2.37 \pm 0.31 \pm 0.11 \pm 0.17$	<sup>1</sup> AAIJ	13BU LHCb	$p p$ at 7, 8 TeV

<sup>1</sup> The last uncertainty is due to the uncertainty of the  $B_c^+$  lifetime measurement.

$\Gamma(D_s^+ \bar{D}^0)/\Gamma_{\text{total}}$				$\Gamma_{28}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.2 \times 10^{-4}$	90	1 AAIJ	21AF LHCb	$p p$ at 13 TeV

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

$\Gamma(D_s^+ D^0)/\Gamma_{\text{total}}$				$\Gamma_{29}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-4}$	90	1 AAIJ	21AF LHCb	$p p$ at 13 TeV

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

$\Gamma(D^+ \bar{D}^0)/\Gamma_{\text{total}}$				$\Gamma_{30}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.9 \times 10^{-4}$	90	1 AAIJ	21AF LHCb	$p p$ at 13 TeV

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

$\Gamma(D^+ D^0)/\Gamma_{\text{total}}$				$\Gamma_{31}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-4}$	90	1 AAIJ	21AF LHCb	$p p$ at 13 TeV

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

$\Gamma(D_s^{*+} \bar{D}^0)/\Gamma_{\text{total}}$				$\Gamma_{32}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.3 \times 10^{-4}$	90	1 AAIJ	21AF LHCb	$p p$ at 13 TeV

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

$\Gamma(D_s^{*+} \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$				$\Gamma_{33}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.6 \times 10^{-4}$	90	1 AAIJ	21AF LHCb	$p p$ at 13 TeV

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

$\Gamma(D_s^{*+} D^0)/\Gamma_{\text{total}}$				$\Gamma_{34}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.9 \times 10^{-3}$	90	1 AAIJ	21AF LHCb	$p p$ at 13 TeV

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

$\Gamma(D_s^{*+} D^*(2007)^0)/\Gamma_{\text{total}}$				$\Gamma_{35}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.6 \times 10^{-4}$	90	1 AAIJ	21AF LHCb	$p p$ at 13 TeV

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

$\Gamma(D^*(2010)^+ \bar{D}^0)/\Gamma_{\text{total}}$				$\Gamma_{36}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.8 \times 10^{-4}$	90	1 AAIJ	21AF LHCb	$p p$ at 13 TeV

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

$\Gamma(D^*(2007)^+ D^0)/\Gamma_{\text{total}}$	$\Gamma_{39}/\Gamma$		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$<2.0 \times 10^{-4}$	90	<sup>1</sup> AAIJ	21AF LHCb $p p$ at 13 TeV
<sup>1</sup> Uses $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$ determined by AAIJ 19AI.			
$\Gamma(D^+ \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$	$\Gamma_{38}/\Gamma$		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$<6.5 \times 10^{-4}$	90	<sup>1</sup> AAIJ	21AF LHCb $p p$ at 13 TeV
<sup>1</sup> Uses $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$ determined by AAIJ 19AI.			
$\Gamma(D^+ D^*(2007)^0)/\Gamma_{\text{total}}$	$\Gamma_{41}/\Gamma$		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$<3.7 \times 10^{-4}$	90	<sup>1</sup> AAIJ	21AF LHCb $p p$ at 13 TeV
<sup>1</sup> Uses $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$ determined by AAIJ 19AI.			
$\Gamma(D_s^{*+} \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$	$\Gamma_{42}/\Gamma$		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$<1.3 \times 10^{-3}$	90	<sup>1</sup> AAIJ	21AF LHCb $p p$ at 13 TeV
<sup>1</sup> Uses $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$ determined by AAIJ 19AI.			
$\Gamma(D_s^{*+} D^*(2007)^0)/\Gamma_{\text{total}}$	$\Gamma_{43}/\Gamma$		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$<1.3 \times 10^{-3}$	90	<sup>1</sup> AAIJ	21AF LHCb $p p$ at 13 TeV
<sup>1</sup> Uses $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$ determined by AAIJ 19AI.			
$\Gamma(D^*(2010)^+ \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$	$\Gamma_{44}/\Gamma$		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$<1.0 \times 10^{-3}$	90	<sup>1</sup> AAIJ	21AF LHCb $p p$ at 13 TeV
<sup>1</sup> Uses $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$ determined by AAIJ 19AI.			
$\Gamma(D^*(2010)^+ D^*(2007)^0)/\Gamma_{\text{total}}$	$\Gamma_{45}/\Gamma$		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$<7.7 \times 10^{-4}$	90	<sup>1</sup> AAIJ	21AF LHCb $p p$ at 13 TeV
<sup>1</sup> Uses $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$ determined by AAIJ 19AI.			
$\Gamma(B_s^0 \pi^+)/\Gamma(J/\psi(1S)\pi^+)$	$\Gamma_{53}/\Gamma_4$		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$91 \pm 10 \pm 8.5$		<sup>1</sup> AAIJ	23M LHCb $p p$ at 13 TeV
<sup>1</sup> The $B_s^0$ mesons are reconstructed via the decays $B_s^0 \rightarrow J/\psi \phi$ and $B_s^0 \rightarrow D_s^- \pi^+$ . The third uncertainty includes systematic ( $\pm 8$ ) and imprecise knowledge of the branching fractions ( $\pm 3$ ).			

## POLARIZATION IN $B_c^+$ DECAY

In decays involving two vector mesons, one can distinguish among the states in which meson polarizations are both longitudinal ( $L$ ) or both are transverse and parallel ( $\parallel$ ) or perpendicular ( $\perp$ ) to each other with the parameters  $\Gamma_L/\Gamma$ ,  $\Gamma_\perp/\Gamma$ , and the relative phases  $\phi_\parallel$  and  $\phi_\perp$ . See the definitions in the note on “Polarization in  $B$  Decays” review in the  $B^0$  Particle Listings.

### $\Gamma_L/\Gamma$ in $B_c^+ \rightarrow J/\psi D_s^{*+}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.34±0.09 OUR AVERAGE</b>			
0.30±0.10±0.04	1, <sup>2</sup> AAD	220 ATLAS	$p p$ at 13 TeV
0.48±0.20	<sup>3</sup> AAIJ	13AS LHCb	$p p$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.62±0.24	<sup>4</sup> AAD	16H ATLAS	$p p$ at 7, 8 TeV
<sup>1</sup> Supersedes the measurement of AAD 16H. <sup>2</sup> AAD 220 measures $1 - \Gamma_L/\Gamma = 0.70 \pm 0.10 \pm 0.04$ . <sup>3</sup> AAIJ 13AS measures $1 - \Gamma_L/\Gamma = 0.52 \pm 0.20$ . <sup>4</sup> AAD 16H measures $1 - \Gamma_L/\Gamma = 0.38 \pm 0.24$ .			

### $A_P(B_c^+)$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>-1.0±1.0 OUR AVERAGE</b>			
-2.5±2.1±0.5	<sup>1</sup> AAIJ	19AI LHCb	$p p$ at 7 TeV
-0.5±1.1±0.4	<sup>1</sup> AAIJ	19AI LHCb	$p p$ at 13 TeV
<sup>1</sup> Measured using $B_c^+$ semileptonic decays.			

## $B_c^+$ REFERENCES

AAIJ	23M	JHEP 2307 066	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAD	22O	JHEP 2208 087	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	22P	JHEP 2201 065	R. Aaij <i>et al.</i>	(LHCb Collab.)
AABOUD	21	PR D104 012010	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AAIJ	21AF	JHEP 2112 117	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	20R	JHEP 2007 123	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	19AI	PR D100 112006	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18C	PRL 120 121801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18P	NP B930 563	R. Aaij <i>et al.</i>	(LHCb Collab.)
SIRUNYAN	18BY	EPJ C78 457	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAIJ	17AG	PRL 118 111803	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17L	PR D95 032005	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAD	16H	EPJ C76 4	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	16AF	JHEP 1609 153	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16AT	PR D94 091102	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16K	PL B759 313	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	16A	PR D93 052001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAIJ	15AY	PR D92 072007	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15G	PL B742 29	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15M	PRL 114 132001	R. Aaij <i>et al.</i>	(LHCb Collab.)
KHACHATRY...	15AA	JHEP 1501 063	V. Khachatryan <i>et al.</i>	(CMS Collab.)
AAIJ	14AQ	PRL 113 152003	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14G	EPJ C74 2839	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14P	JHEP 1405 148	R. Aaij <i>et al.</i>	(LHCb Collab.)

AAIJ	14W	PR D90 032009	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AM	PR D87 071103	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AS	PR D87 112012	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D89 019901 (errat.)	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BS	PL B726 646	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BU	PRL 111 181801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BY	JHEP 1309 075	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13CA	JHEP 1311 094	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13R	JHEP 1302 043	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	13	PR D87 011101	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAIJ	12AV	PRL 109 232001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12Y	PRL 108 251802	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	09H	PRL 102 092001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
AALTONEN	08M	PRL 100 182002	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	08T	PRL 101 012001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABULENCIA	06C	PRL 96 082002	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABULENCIA	06O	PRL 97 012002	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABE	98M	PRL 81 2432	F. Abe <i>et al.</i>	(CDF Collab.)
Also		PR D58 112004	F. Abe <i>et al.</i>	(CDF Collab.)
ACKERSTAFF	98O	PL B420 157	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	98Q	EPJ C4 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
PDG	98	EPJ C3 1	C. Caso <i>et al.</i>	(PDG Collab.)
ABREU	97E	PL B398 207	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BARATE	97H	PL B402 213	R. Barate <i>et al.</i>	(ALEPH Collab.)
ABE	96R	PRL 77 5176	F. Abe <i>et al.</i>	(CDF Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	(PDG Collab.)

---