

$J/\psi(1S)$	$I^G(J^{PC}) = 0^-(1^{--})$
--------------------------------	-----------------------------

 $J/\psi(1S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3096.900 ± 0.006 OUR AVERAGE				
3096.900 $\pm 0.002 \pm 0.006$		¹ ANASHIN 15	KEDR	$e^+ e^- \rightarrow$ hadrons
3096.89 ± 0.09	502	² ARTAMONOV 00	OLYA	$e^+ e^- \rightarrow$ hadrons
3096.91 $\pm 0.03 \pm 0.01$		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+ e^-$
3096.95 $\pm 0.1 \pm 0.3$	193	BAGLIN	SPEC	$\bar{p}p \rightarrow e^+ e^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3096.66 $\pm 0.19 \pm 0.02$	6.1k	⁴ AAIJ 15BI	LHCb	$p\bar{p} \rightarrow J/\psi X$
3096.917 $\pm 0.010 \pm 0.007$		AULCHENKO 03	KEDR	$e^+ e^- \rightarrow$ hadrons
3097.5 ± 0.3		GRIBUSHIN 96	FMPS	$515 \pi^- Be \rightarrow 2\mu X$
3098.4 ± 2.0	38k	LEMOIGNE 82	GOLI	$185 \pi^- Be \rightarrow \gamma\mu^+\mu^- A$
3096.93 ± 0.09	502	⁵ ZHOLENTZ 80	REDE	$e^+ e^-$
3097.0 ± 1		⁶ BRANDELIK 79c	DASP	$e^+ e^-$

¹ Supersedes AULCHENKO 03.² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).³ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.⁴ From a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays. Systematic uncertainties not estimated.⁵ Superseded by ARTAMONOV 00.⁶ From a simultaneous fit to $e^+ e^-$, $\mu^+ \mu^-$ and hadronic channels assuming $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$. **$J/\psi(1S)$ WIDTH**

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
92.6 ± 1.7 OUR AVERAGE				
Error includes scale factor of 1.1.				
92.45 $\pm 1.40 \pm 1.48$		¹ ANASHIN 20	KEDR	$e^+ e^-$
96.1 ± 3.2	13k	² ADAMS 06A	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
84.4 ± 8.9		BAI 95B	BES	$e^+ e^-$
91 $\pm 11 \pm 6$		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+ e^-$
85.5 ± 6.1 ± 5.8		⁴ HSUEH 92	RVUE	See Υ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •				
92.94 ± 1.83		^{5,6} ANASHIN 18A	KEDR	$e^+ e^-$
94.1 ± 2.7		⁷ ANASHIN 10	KEDR	$3.097 e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
93.7 ± 3.5	7.8k	² AUBERT 04	BABR	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$

¹ Based on the same dataset as ANASHIN 18A and correlated to the values reported there² Calculated by us from the reported values of $\Gamma(e^+ e^-) \times B(\mu^+ \mu^-)$ using $B(e^+ e^-) = (5.94 \pm 0.06)\%$ and $B(\mu^+ \mu^-) = (5.93 \pm 0.06)\%$.

³ The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].⁴ Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79C.⁵ Using $\Gamma(e^+e^-)$ from ANASHIN 18A and $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ from PDG 16.⁶ Superseded by ANASHIN 20 that is based on the same dataset .⁷ Assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ and using $\Gamma(e^+e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$.

$J/\psi(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(87.7 \pm 0.5) %	
Γ_2 virtual $\gamma \rightarrow$ hadrons	(13.46 \pm 0.07) %	
Γ_3 ggg	(64.1 \pm 1.0) %	
Γ_4 γgg	(8.8 \pm 1.1) %	
Γ_5 e^+e^-	(5.971 \pm 0.032) %	
Γ_6 $e^+e^- \gamma$	[a] (8.8 \pm 1.4) $\times 10^{-3}$	
Γ_7 $\mu^+\mu^-$	(5.961 \pm 0.033) %	

Decays involving hadronic resonances

Γ_8 $\rho\pi$	(1.88 \pm 0.12) %	
Γ_9 $\rho^0\pi^0$	(6.2 \pm 0.6) $\times 10^{-3}$	
Γ_{10} $a_2(1320)^0\pi^+\pi^- \rightarrow 2(\pi^+\pi^-)\pi^0$	(2.8 \pm 0.6) $\times 10^{-3}$	
Γ_{11} $a_2(1320)^+\pi^-\pi^0 + \text{c.c} \rightarrow 2(\pi^+\pi^-)\pi^0$	(3.7 \pm 0.7) $\times 10^{-3}$	
Γ_{12} $a_2(1320)\rho$	(1.09 \pm 0.22) %	
Γ_{13} $\eta\pi^+\pi^-$	(3.8 \pm 0.7) $\times 10^{-4}$	
Γ_{14} $\eta\rho$	(1.93 \pm 0.23) $\times 10^{-4}$	
Γ_{15} $\eta\pi^+\pi^-\pi^0$	(1.17 \pm 0.20) %	
Γ_{16} $\eta\pi^+\pi^-3\pi^0$	(4.9 \pm 1.0) $\times 10^{-3}$	
Γ_{17} $\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-$	(1.2 \pm 0.4) $\times 10^{-4}$	
Γ_{18} $\eta\phi(2170) \rightarrow \eta K^*(892)^0\bar{K}^*(892)^0$	< 2.52 $\times 10^{-4}$	CL=90%
Γ_{19} ηK^+K^-	(8.6 \pm 3.0) $\times 10^{-4}$	
Γ_{20} $\eta K^\pm K_S^0\pi^\mp$	[b] (2.2 \pm 0.4) $\times 10^{-3}$	
Γ_{21} $\eta K^*(892)^0\bar{K}^*(892)^0$	(1.15 \pm 0.26) $\times 10^{-3}$	
Γ_{22} $\rho\eta'(958)$	(8.1 \pm 0.8) $\times 10^{-5}$	S=1.6
Γ_{23} $\rho^\pm\pi^\mp\pi^+\pi^-2\pi^0$	(2.8 \pm 0.8) %	
Γ_{24} $\rho^+\rho^-\pi^+\pi^-\pi^0$	(6 \pm 4) $\times 10^{-3}$	
Γ_{25} $\rho^+K^+K^-\pi^- + \text{c.c} \rightarrow K^+K^-\pi^+\pi^-\pi^0$	(3.5 \pm 0.8) $\times 10^{-3}$	
Γ_{26} $\rho^\mp K^\pm K_S^0$	(1.9 \pm 0.4) $\times 10^{-3}$	
Γ_{27} $h_1(1415)\eta' \rightarrow \gamma\eta\eta'$		

Γ_{28}	$h_1(1595)\eta' \rightarrow \gamma\eta\eta'$			
Γ_{29}	$\rho(1450)\pi$	seen		
Γ_{30}	$\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0$	(2.2 \pm 1.1) $\times 10^{-4}$		
Γ_{31}	$\rho(1450)^\pm\pi^\mp \rightarrow K_S^0 K^\pm\pi^\mp$	(3.3 \pm 0.6) $\times 10^{-4}$		
Γ_{32}	$\rho(1450)^0\pi^0 \rightarrow K^+K^-\pi^0$	(2.7 \pm 0.6) $\times 10^{-4}$		
Γ_{33}	$\rho(1450)\eta'(958) \rightarrow \pi^+\pi^-\eta'(958)$	(3.3 \pm 0.7) $\times 10^{-6}$		
Γ_{34}	$\rho(1700)\pi$	seen		
Γ_{35}	$\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0$	(1.6 \pm 1.1) $\times 10^{-4}$		
Γ_{36}	$\rho(2150)\pi$	seen		
Γ_{37}	$\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$	(10 \pm 40) $\times 10^{-6}$		
Γ_{38}	$\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0$			
Γ_{39}	$\omega\pi^0$	(4.5 \pm 0.5) $\times 10^{-4}$	$S=1.4$	
Γ_{40}	$\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0$	(1.6 \pm 0.7) $\times 10^{-5}$		
Γ_{41}	$\omega\pi^+\pi^-$	(8.5 \pm 1.0) $\times 10^{-3}$	$S=1.3$	
Γ_{42}	$\omega\pi^0\pi^0$	(3.4 \pm 0.8) $\times 10^{-3}$		
Γ_{43}	$\omega 3\pi^0$	(1.9 \pm 0.6) $\times 10^{-3}$		
Γ_{44}	$\omega f_2(1270)$	(4.3 \pm 0.6) $\times 10^{-3}$		
Γ_{45}	$\omega\eta$	(1.74 \pm 0.20) $\times 10^{-3}$	$S=1.6$	
Γ_{46}	$\omega\pi^+\pi^-\pi^0$	(4.0 \pm 0.7) $\times 10^{-3}$		
Γ_{47}	$\omega\pi^0\eta$	(3.4 \pm 1.7) $\times 10^{-4}$		
Γ_{48}	$\omega\pi^+\pi^+\pi^-\pi^-$	(8.5 \pm 3.4) $\times 10^{-3}$		
Γ_{49}	$\omega\pi^+\pi^-2\pi^0$	(3.3 \pm 0.5) %		
Γ_{50}	$\omega\eta'\pi^+\pi^-$	(1.12 \pm 0.13) $\times 10^{-3}$		
Γ_{51}	$\omega\eta'(958)$	(1.89 \pm 0.18) $\times 10^{-4}$		
Γ_{52}	$\omega f_0(980)$	(1.4 \pm 0.5) $\times 10^{-4}$		
Γ_{53}	$\omega f_0(1710) \rightarrow \omega K\bar{K}$	(4.8 \pm 1.1) $\times 10^{-4}$		
Γ_{54}	$\omega f_1(1420)$	(6.8 \pm 2.4) $\times 10^{-4}$		
Γ_{55}	$\omega f'_2(1525)$	< 2.2 $\times 10^{-4}$	$CL=90\%$	
Γ_{56}	$\omega X(1835) \rightarrow \omega p\bar{p}$	< 3.9 $\times 10^{-6}$	$CL=95\%$	
Γ_{57}	$\omega X(1835), X \rightarrow \eta'\pi^+\pi^-$	< 6.2 $\times 10^{-5}$		
Γ_{58}	ωK^+K^-	(1.52 \pm 0.31) $\times 10^{-3}$		
Γ_{59}	$\omega K^\pm K_S^0\pi^\mp$	[b] (3.4 \pm 0.5) $\times 10^{-3}$		
Γ_{60}	$\omega K\bar{K}$	(1.9 \pm 0.4) $\times 10^{-3}$		
Γ_{61}	$\omega K^*(892)\bar{K} + c.c.$	(6.1 \pm 0.9) $\times 10^{-3}$		
Γ_{62}	$\eta' K^{*\pm}\bar{K}^\mp$	(1.48 \pm 0.13) $\times 10^{-3}$		
Γ_{63}	$\eta' K^{*0}\bar{K}^0 + c.c.$	(1.66 \pm 0.21) $\times 10^{-3}$		
Γ_{64}	$\eta' h_1(1415) \rightarrow \eta' K^*\bar{K} + c.c.$	(2.16 \pm 0.31) $\times 10^{-4}$		
Γ_{65}	$\eta' h_1(1415) \rightarrow \eta' K^{*\pm}\bar{K}^\mp$	(1.51 \pm 0.23) $\times 10^{-4}$		
Γ_{66}	$\eta' h_1(1415) \rightarrow \gamma\eta'\eta'$	(4.7 \pm 1.1) $\times 10^{-7}$		
Γ_{67}	$\bar{K}K^*(892)+c.c.$	seen		
Γ_{68}	$\bar{K}K^*(892)+c.c. \rightarrow K_S^0 K^\pm\pi^\mp$	(4.8 \pm 0.5) $\times 10^{-3}$		

Γ_{69}	$K^+ K^*(892)^- + \text{c.c.}$	$(6.0 \pm 0.8) \times 10^{-3}$	S=2.9
Γ_{70}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(2.69 \pm 0.13) \times 10^{-3}$	
Γ_{71}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.0 \pm 0.4) \times 10^{-3}$	
Γ_{72}	$K^0 \bar{K}^*(892)^0 + \text{c.c.}$	$(4.2 \pm 0.4) \times 10^{-3}$	
Γ_{73}	$K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.2 \pm 0.4) \times 10^{-3}$	
Γ_{74}	$\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}$	$(5.7 \pm 0.8) \times 10^{-3}$	
Γ_{75}	$K^*(892)^\pm K^\mp \pi^0$	$(4.1 \pm 1.3) \times 10^{-3}$	
Γ_{76}	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$(2.0 \pm 0.5) \times 10^{-3}$	
Γ_{77}	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$	$(6.7 \pm 2.2) \times 10^{-4}$	
Γ_{78}	$K^*(892)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-$	$(3.8 \pm 0.5) \times 10^{-3}$	
Γ_{79}	$K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(6.3 \pm 0.6) \times 10^{-6}$	
Γ_{80}	$K^*(892)^0 K_S^0 \pi^0$	$(7 \pm 4) \times 10^{-4}$	
Γ_{81}	$K^*(892)^\pm K^*(700)^\mp$	$(1.1 \pm 1.0) \times 10^{-3}$	
Γ_{82}	$K^*(892)^0 \bar{K}^*(892)^0$	$(2.3 \pm 0.6) \times 10^{-4}$	
Γ_{83}	$K^*(892)^\pm K^*(892)^\mp$	$(1.00 \pm 0.22) \times 10^{-3}$	
Γ_{84}	$K_1(1400)^\pm K^\mp$	$(3.8 \pm 1.4) \times 10^{-3}$	
Γ_{85}	$K^*(1410) \bar{K} + \text{c.c.}$	seen	
Γ_{86}	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0$	$(7 \pm 4) \times 10^{-5}$	
Γ_{87}	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(8 \pm 5) \times 10^{-5}$	
Γ_{88}	$K_2^*(1430) \bar{K} + \text{c.c.}$	seen	
Γ_{89}	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0$	$(1.0 \pm 0.5) \times 10^{-4}$	
Γ_{90}	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(3.8 \pm 1.0) \times 10^{-4}$	
Γ_{91}	$\bar{K}_2^*(1430) K + \text{c.c.}$	$< 4.0 \times 10^{-3}$	CL=90%
Γ_{92}	$K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(2.69 \pm 0.25) \times 10^{-4}$	
Γ_{93}	$K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-$	$(2.6 \pm 0.9) \times 10^{-3}$	
Γ_{94}	$K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}$	$(3.6 \pm 1.8) \times 10^{-3}$	
Γ_{95}	$\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}$	$(4.67 \pm 0.29) \times 10^{-3}$	
Γ_{96}	$K_2^*(1430)^- K^*(892)^+ + \text{c.c.}$	$(3.4 \pm 2.9) \times 10^{-3}$	

Γ_{97}	$K_2^*(1430)^- K^*(892)^+ +$ c.c. $\rightarrow K^*(892)^+ K_S^0 \pi^- +$ c.c.	$(4 \pm 4) \times 10^{-4}$
Γ_{98}	$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$	$< 2.9 \times 10^{-3} \text{ CL}=90\%$
Γ_{99}	$\bar{K}_2(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow$ $K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(6.9 \pm 0.9) \times 10^{-4}$
Γ_{100}	$K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(1.10 \pm 0.60) \times 10^{-5}$
Γ_{101}	$K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(6.2 \pm 2.9) \times 10^{-6}$
Γ_{102}	$K_1(1270)^{\pm} K^{\mp}$	$< 3.0 \times 10^{-3} \text{ CL}=90\%$
Γ_{103}	$K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(8.5 \pm 2.5) \times 10^{-7}$
Γ_{104}	$a_2(1320)^{\pm} \pi^{\mp}$	$[b] < 4.3 \times 10^{-3} \text{ CL}=90\%$
Γ_{105}	$\phi \pi^0$	$3 \times 10^{-6} \text{ or } 1 \times 10^{-7}$
Γ_{106}	$\phi \pi^+ \pi^-$	$(9.4 \pm 1.5) \times 10^{-4} \quad S=1.7$
Γ_{107}	$\phi \pi^0 \pi^0$	$(5.0 \pm 1.0) \times 10^{-4}$
Γ_{108}	$\phi 2(\pi^+ \pi^-)$	$(1.60 \pm 0.32) \times 10^{-3}$
Γ_{109}	$\phi \eta$	$(7.4 \pm 0.6) \times 10^{-4} \quad S=1.2$
Γ_{110}	$\phi \eta'(958)$	$(4.6 \pm 0.5) \times 10^{-4} \quad S=2.2$
Γ_{111}	$\phi \eta \eta'$	$(2.32 \pm 0.17) \times 10^{-4}$
Γ_{112}	$\phi f_0(980)$	$(3.2 \pm 0.9) \times 10^{-4} \quad S=1.9$
Γ_{113}	$\phi f_0(980) \rightarrow \phi \pi^+ \pi^-$	$(2.60 \pm 0.34) \times 10^{-4}$
Γ_{114}	$\phi f_0(980) \rightarrow \phi \pi^0 \pi^0$	$(1.8 \pm 0.5) \times 10^{-4}$
Γ_{115}	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$	$(4.5 \pm 1.0) \times 10^{-6}$
Γ_{116}	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 p^0 \pi^0$	$(1.7 \pm 0.6) \times 10^{-6}$
Γ_{117}	$\phi f_0(980) \eta \rightarrow \eta \phi \pi^+ \pi^-$	$(3.2 \pm 1.0) \times 10^{-4}$
Γ_{118}	$\phi a_0(980)^0 \rightarrow \phi \eta \pi^0$	$(4.4 \pm 1.4) \times 10^{-6}$
Γ_{119}	$\phi f_2(1270)$	$(3.2 \pm 0.6) \times 10^{-4}$
Γ_{120}	$\phi f_1(1285)$	$(2.6 \pm 0.5) \times 10^{-4}$
Γ_{121}	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow$ $\phi \pi^0 \pi^+ \pi^-$	$(9.4 \pm 2.8) \times 10^{-7}$
Γ_{122}	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow$ $\phi 3\pi^0$	$(2.1 \pm 2.2) \times 10^{-7}$
Γ_{123}	$\phi \eta(1405) \rightarrow \phi \eta \pi^+ \pi^-$	$(2.0 \pm 1.0) \times 10^{-5}$
Γ_{124}	$\phi f'_2(1525)$	$(8 \pm 4) \times 10^{-4} \quad S=2.7$
Γ_{125}	$\phi X(1835) \rightarrow \phi p \bar{p}$	$< 2.1 \times 10^{-7} \text{ CL}=90\%$
Γ_{126}	$\phi X(1835) \rightarrow \phi \eta \pi^+ \pi^-$	$< 2.8 \times 10^{-4} \text{ CL}=90\%$
Γ_{127}	$\phi X(1870) \rightarrow \phi \eta \pi^+ \pi^-$	$< 6.13 \times 10^{-5} \text{ CL}=90\%$
Γ_{128}	$\phi K \bar{K}$	$(1.77 \pm 0.16) \times 10^{-3} \quad S=1.3$
Γ_{129}	$\phi f_0(1710) \rightarrow \phi K \bar{K}$	$(3.6 \pm 0.6) \times 10^{-4}$
Γ_{130}	$\phi K^+ K^-$	$(8.3 \pm 1.1) \times 10^{-4}$
Γ_{131}	$\phi K_S^0 K_S^0$	$(5.9 \pm 1.5) \times 10^{-4}$
Γ_{132}	$\phi K^{\pm} K_S^0 \pi^{\mp}$	$[b] (7.2 \pm 0.8) \times 10^{-4}$

Γ_{133}	$\phi K^*(892) \bar{K} + \text{c.c.}$	$(2.18 \pm 0.23) \times 10^{-3}$	
Γ_{134}	$b_1(1235)^\pm \pi^\mp$	$[b] (3.0 \pm 0.5) \times 10^{-3}$	
Γ_{135}	$b_1(1235)^0 \pi^0$	$(2.3 \pm 0.6) \times 10^{-3}$	
Γ_{136}	$f'_2(1525) K^+ K^-$	$(1.04 \pm 0.35) \times 10^{-3}$	
Γ_{137}	$\Delta(1232)^+ \bar{p}$	$< 1 \times 10^{-4}$	CL=90%
Γ_{138}	$\Delta(1232)^{++} \bar{p} \pi^-$	$(1.6 \pm 0.5) \times 10^{-3}$	
Γ_{139}	$\Delta(1232)^{++} \bar{\Delta}(1232)^{--}$	$(1.10 \pm 0.29) \times 10^{-3}$	
Γ_{140}	$\bar{\Sigma}(1385)^0 p K^-$	$(5.1 \pm 3.2) \times 10^{-4}$	
Γ_{141}	$\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.}$	$< 8.2 \times 10^{-6}$	CL=90%
Γ_{142}	$\Sigma(1385)^- \bar{\Sigma}^+ + \text{c.c.}$	$[b] (3.0 \pm 0.7) \times 10^{-4}$	
Γ_{143}	$\Sigma(1385)^+ \bar{\Sigma}^- + \text{c.c.}$	$(3.3 \pm 0.8) \times 10^{-4}$	
Γ_{144}	$\Sigma(1385)^- \bar{\Sigma}(1385)^+ + \text{c.c.}$	$[b] (1.08 \pm 0.06) \times 10^{-3}$	
Γ_{145}	$\Sigma(1385)^+ \bar{\Sigma}(1385)^- + \text{c.c.}$	$(1.25 \pm 0.07) \times 10^{-3}$	
Γ_{146}	$\Sigma(1385)^0 \bar{\Sigma}(1385)^0$	$(1.07 \pm 0.08) \times 10^{-3}$	
Γ_{147}	$\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda}$	$< 4.1 \times 10^{-6}$	CL=90%
Γ_{148}	$\bar{\Lambda}(1520) \Lambda + \text{c.c.}$	$< 1.80 \times 10^{-3}$	CL=90%
Γ_{149}	$\Xi^0 \bar{\Xi}^0$	$(1.17 \pm 0.04) \times 10^{-3}$	
Γ_{150}	$\Xi(1530)^- \bar{\Xi}^+ + \text{c.c.}$	$(3.18 \pm 0.08) \times 10^{-4}$	
Γ_{151}	$\Xi(1530)^0 \bar{\Xi}^0$	$(3.2 \pm 1.4) \times 10^{-4}$	
Γ_{152}	$\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}$	$[c] < 1.1 \times 10^{-5}$	CL=90%
Γ_{153}	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	$[c] < 2.1 \times 10^{-5}$	CL=90%
Γ_{154}	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	$[c] < 1.6 \times 10^{-5}$	CL=90%
Γ_{155}	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	$[c] < 5.6 \times 10^{-5}$	CL=90%
Γ_{156}	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	$[c] < 1.1 \times 10^{-5}$	CL=90%

Decays into stable hadrons

Γ_{157}	$2(\pi^+ \pi^-) \pi^0$	$(4.2 \pm 0.4) \%$	S=2.1
Γ_{158}	$3(\pi^+ \pi^-) \pi^0$	$(2.9 \pm 0.6) \%$	
Γ_{159}	$\pi^+ \pi^- 3\pi^0$	$(1.9 \pm 0.9) \%$	
Γ_{160}	$\rho^\pm \pi^\mp \pi^0 \pi^0$	$(1.41 \pm 0.22) \%$	
Γ_{161}	$\rho^+ \rho^- \pi^0$	$(6.0 \pm 1.1) \times 10^{-3}$	
Γ_{162}	$\pi^+ \pi^- 4\pi^0$	$(6.5 \pm 1.3) \times 10^{-3}$	
Γ_{163}	$\pi^+ \pi^- \pi^0$	$(2.00 \pm 0.07) \%$	S=2.0
Γ_{164}	$2(\pi^+ \pi^- \pi^0)$	$(1.61 \pm 0.20) \%$	
Γ_{165}	$\pi^+ \pi^- \pi^0 K^+ K^-$	$(1.52 \pm 0.27) \%$	S=1.4
Γ_{166}	$\pi^+ \pi^-$	$(1.47 \pm 0.14) \times 10^{-4}$	
Γ_{167}	$2(\pi^+ \pi^-)$	$(3.20 \pm 0.25) \times 10^{-3}$	S=1.2
Γ_{168}	$3(\pi^+ \pi^-)$	$(4.3 \pm 0.4) \times 10^{-3}$	
Γ_{169}	$2(\pi^+ \pi^-) 3\pi^0$	$(6.2 \pm 0.9) \%$	
Γ_{170}	$4(\pi^+ \pi^-) \pi^0$	$(9.0 \pm 3.0) \times 10^{-3}$	
Γ_{171}	$2(\pi^+ \pi^-) \eta$	$(2.29 \pm 0.28) \times 10^{-3}$	
Γ_{172}	$3(\pi^+ \pi^-) \eta$	$(7.2 \pm 1.5) \times 10^{-4}$	
Γ_{173}	$2(\pi^+ \pi^- \pi^0) \eta$	$(1.6 \pm 0.5) \times 10^{-3}$	

Γ_{174}	$\pi^+ \pi^- \pi^0 \pi^0 \eta$	$(2.4 \pm 0.5) \times 10^{-3}$
Γ_{175}	$\rho^\pm \pi^\mp \pi^0 \eta$	$(1.9 \pm 0.8) \times 10^{-3}$
Γ_{176}	$K^+ K^-$	$(2.86 \pm 0.21) \times 10^{-4}$
Γ_{177}	$K_S^0 K_L^0$	$(1.95 \pm 0.11) \times 10^{-4}$
Γ_{178}	$K_S^0 K_S^0$	$< 1.4 \times 10^{-8}$ CL=95%
Γ_{179}	$K \bar{K} \pi$	$(6.1 \pm 1.0) \times 10^{-3}$
Γ_{180}	$K^+ K^- \pi^0$	$(2.88 \pm 0.12) \times 10^{-3}$
Γ_{181}	$K_S^0 K^\pm \pi^\mp$	$(5.3 \pm 0.5) \times 10^{-3}$
Γ_{182}	$K_S^0 K_L^0 \pi^0$	$(2.06 \pm 0.26) \times 10^{-3}$
Γ_{183}	$K^*(892)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0$	$(1.21 \pm 0.18) \times 10^{-3}$
Γ_{184}	$K_2^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0$	$(4.3 \pm 1.3) \times 10^{-4}$
Γ_{185}	$K^+ K^- \pi^+ \pi^-$	$(7.0 \pm 1.0) \times 10^{-3}$
Γ_{186}	$K^+ K^- \pi^0 \pi^0$	$(2.13 \pm 0.22) \times 10^{-3}$
Γ_{187}	$K^+ K^- \pi^0 \pi^0 \pi^0$	$(1.61 \pm 0.29) \times 10^{-3}$
Γ_{188}	$K_S^0 K^\pm \pi^\mp \pi^0 \pi^0$	$(5.3 \pm 0.7) \times 10^{-3}$
Γ_{189}	$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(6.3 \pm 0.4) \times 10^{-3}$
Γ_{190}	$K_S^0 K^\pm \rho(770)^\pm \pi^0$	$(2.9 \pm 0.8) \times 10^{-3}$
Γ_{191}	$K_S^0 K_L^0 \pi^+ \pi^-$	$(3.8 \pm 0.6) \times 10^{-3}$
Γ_{192}	$K_S^0 K_L^0 \pi^0 \pi^0$	$(1.9 \pm 0.4) \times 10^{-3}$
Γ_{193}	$K_S^0 K_L^0 \eta$	$(1.45 \pm 0.33) \times 10^{-3}$
Γ_{194}	$K_S^0 K_S^0 \pi^+ \pi^-$	$(1.68 \pm 0.19) \times 10^{-3}$
Γ_{195}	$K^\mp K_S^0 \pi^\pm \pi^0$	$(5.7 \pm 0.5) \times 10^{-3}$
Γ_{196}	$K_S^0 K^\pm \pi^\mp \rho(770)^0$	$(3.1 \pm 0.5) \times 10^{-3}$
Γ_{197}	$K^+ K^- 2(\pi^+ \pi^-)$	$(3.1 \pm 1.3) \times 10^{-3}$
Γ_{198}	$K^+ K^- \pi^+ \pi^- \eta$	$(4.7 \pm 0.7) \times 10^{-3}$
Γ_{199}	$2(K^+ K^-)$	$(7.2 \pm 0.8) \times 10^{-4}$
Γ_{200}	$K^+ K^- K_S^0 K_S^0$	$(4.2 \pm 0.7) \times 10^{-4}$
Γ_{201}	$K_S^0 K^*(892)^0 \pi^+ \pi^-$	$(1.7 \pm 0.6) \times 10^{-3}$
Γ_{202}	$K_S^0 K^*(892)^0 \pi^0 \pi^0$	$(1.01 \pm 0.18) \times 10^{-3}$
Γ_{203}	$K^\mp K^*(892)^\pm \pi^+ \pi^-$	$(3.4 \pm 1.2) \times 10^{-3}$
Γ_{204}	$K^*(892)^\pm K^*(892)^0 \pi^\mp$	$(4.8 \pm 1.0) \times 10^{-3}$
Γ_{205}	$K^\mp K^*(892)^\pm \pi^0 \pi^0$	$(1.57 \pm 0.32) \times 10^{-3}$
Γ_{206}	$K^*(892)^+ K^*(892)^- \pi^0$	$(1.12 \pm 0.23) \%$
Γ_{207}	$p \bar{p}$	$(2.120 \pm 0.029) \times 10^{-3}$
Γ_{208}	$p \bar{p} \pi^0$	$(1.19 \pm 0.08) \times 10^{-3}$
Γ_{209}	$p \bar{p} \pi^+ \pi^-$	$(6.0 \pm 0.5) \times 10^{-3}$
Γ_{210}	$p \bar{p} \pi^+ \pi^- \pi^0$	$[d] (2.3 \pm 0.9) \times 10^{-3}$
Γ_{211}	$p \bar{p} \eta$	$(2.00 \pm 0.12) \times 10^{-3}$
Γ_{212}	$p \bar{p} \rho$	$< 3.1 \times 10^{-4}$ CL=90%
Γ_{213}	$p \bar{p} \omega$	$(9.8 \pm 1.0) \times 10^{-4}$ S=1.3

Γ_{214}	$p\bar{p}\eta'(958)$	$(1.29 \pm 0.14) \times 10^{-4}$	S=2.0
Γ_{215}	$p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta$	$(6.8 \pm 1.8) \times 10^{-5}$	
Γ_{216}	$p\bar{p}\phi$	$(5.19 \pm 0.33) \times 10^{-5}$	
Γ_{217}	$p\bar{n}\pi^-$	$(2.12 \pm 0.09) \times 10^{-3}$	
Γ_{218}	$n\bar{n}$	$(2.09 \pm 0.16) \times 10^{-3}$	
Γ_{219}	$n\bar{n}\pi^+\pi^-$	$(4 \pm 4) \times 10^{-3}$	
Γ_{220}	$nN(1440)$	seen	
Γ_{221}	$nN(1520)$	seen	
Γ_{222}	$nN(1535)$	seen	
Γ_{223}	$\Lambda\bar{\Lambda}$	$(1.88 \pm 0.08) \times 10^{-3}$	S=2.6
Γ_{224}	$\Lambda\bar{\Lambda}\pi^0$	$(3.8 \pm 0.4) \times 10^{-5}$	
Γ_{225}	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(4.3 \pm 1.0) \times 10^{-3}$	
Γ_{226}	$\Lambda\bar{\Lambda}\eta$	$(1.62 \pm 0.17) \times 10^{-4}$	
Γ_{227}	$\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.}$	[b] $(1.26 \pm 0.05) \times 10^{-3}$	S=1.2
Γ_{228}	$\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.}$	$(1.21 \pm 0.07) \times 10^{-3}$	S=1.8
Γ_{229}	$pK^-\bar{\Lambda} + \text{c.c.}$	$(8.6 \pm 1.1) \times 10^{-4}$	
Γ_{230}	$pK^-\bar{\Sigma}^0$	$(2.9 \pm 0.8) \times 10^{-4}$	
Γ_{231}	$\bar{\Lambda}nK_S^0 + \text{c.c.}$	$(6.5 \pm 1.1) \times 10^{-4}$	
Γ_{232}	$\Lambda\bar{\Sigma} + \text{c.c.}$	$(2.83 \pm 0.23) \times 10^{-5}$	
Γ_{233}	$\Sigma^+\bar{\Sigma}^-$	$(1.07 \pm 0.04) \times 10^{-3}$	
Γ_{234}	$\Sigma^0\bar{\Sigma}^0$	$(1.172 \pm 0.032) \times 10^{-3}$	S=1.4
Γ_{235}	$\Sigma^+\bar{\Sigma}^-\eta$	$(6.3 \pm 0.4) \times 10^{-5}$	
Γ_{236}	$\Xi^-\bar{\Xi}^+$	$(9.7 \pm 0.8) \times 10^{-4}$	S=1.4

Radiative decays

Γ_{237}	$\gamma\eta_c(1S)$	$(1.41 \pm 0.14) \%$	S=1.3
Γ_{238}	$\gamma\eta_c(1S) \rightarrow 3\gamma$	seen	
Γ_{239}	$\gamma\eta_c(1S) \rightarrow \gamma\eta\eta\eta'$	seen	
Γ_{240}	3γ	$(1.16 \pm 0.22) \times 10^{-5}$	
Γ_{241}	4γ	$< 9 \times 10^{-6}$	CL=90%
Γ_{242}	5γ	$< 1.5 \times 10^{-5}$	CL=90%
Γ_{243}	$\gamma\pi^0$	$(3.39 \pm 0.08) \times 10^{-5}$	
Γ_{244}	$\gamma\pi^0\pi^0$	$(1.15 \pm 0.05) \times 10^{-3}$	
Γ_{245}	$\gamma 2\pi^+ 2\pi^-$	$(2.8 \pm 0.5) \times 10^{-3}$	S=1.9
Γ_{246}	$\gamma f_2(1270)f_2(1270)$	$(9.5 \pm 1.7) \times 10^{-4}$	
Γ_{247}	$\gamma f_2(1270)f_2(1270)$ (non resonant)	$(8.2 \pm 1.9) \times 10^{-4}$	
Γ_{248}	$\gamma\pi^+\pi^- 2\pi^0$	$(8.3 \pm 3.1) \times 10^{-3}$	
Γ_{249}	$\gamma K_S^0 K_S^0$	$(8.1 \pm 0.4) \times 10^{-4}$	
Γ_{250}	$\gamma(K\bar{K}\pi) [JPC = 0 - +]$	$(7 \pm 4) \times 10^{-4}$	S=2.1
Γ_{251}	$\gamma K^+ K^- \pi^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$	
Γ_{252}	$\gamma K^*(892)\bar{K}^*(892)$	$(4.0 \pm 1.3) \times 10^{-3}$	
Γ_{253}	$\gamma\eta$	$(1.090 \pm 0.013) \times 10^{-3}$	
Γ_{254}	$\gamma\eta\pi^0$	$(2.14 \pm 0.31) \times 10^{-5}$	

Γ_{255}	$\gamma f_0(500) \rightarrow \gamma\pi\pi$				
Γ_{256}	$\gamma f_0(500) \rightarrow \gamma K\bar{K}$				
Γ_{257}	$\gamma f_0(500) \rightarrow \gamma\eta\eta$				
Γ_{258}	$\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0$	< 2.5	$\times 10^{-6}$	CL=95%	
Γ_{259}	$\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0$	< 6.6	$\times 10^{-6}$	CL=95%	
Γ_{260}	$\gamma\eta\pi\pi$	(6.1 \pm 1.0)	$\times 10^{-3}$		
Γ_{261}	$\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-$	(6.2 \pm 2.4)	$\times 10^{-4}$		
Γ_{262}	$\gamma\eta'(958)$	(5.28 \pm 0.06)	$\times 10^{-3}$	S=1.3	
Γ_{263}	$\gamma f_0(980) \rightarrow \gamma\pi\pi$				
Γ_{264}	$\gamma f_0(980) \rightarrow \gamma K\bar{K}$				
Γ_{265}	$\gamma\rho\rho$	(4.5 \pm 0.8)	$\times 10^{-3}$		
Γ_{266}	$\gamma\rho\omega$	< 5.4	$\times 10^{-4}$	CL=90%	
Γ_{267}	$\gamma\rho\phi$	< 8.8	$\times 10^{-5}$	CL=90%	
Γ_{268}	$\gamma\omega\omega$	(1.61 \pm 0.33)	$\times 10^{-3}$		
Γ_{269}	$\gamma\phi\phi$	(4.0 \pm 1.2)	$\times 10^{-4}$	S=2.1	
Γ_{270}	$\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi$	(2.8 \pm 0.6)	$\times 10^{-3}$	S=1.6	
Γ_{271}	$\gamma\eta(1405/1475) \rightarrow \gamma\eta\rho^0$	(7.8 \pm 2.0)	$\times 10^{-5}$	S=1.8	
Γ_{272}	$\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-$	(3.0 \pm 0.5)	$\times 10^{-4}$		
Γ_{273}	$\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0$	(1.7 \pm 0.4)	$\times 10^{-3}$	S=1.3	
Γ_{274}	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi$	< 8.2	$\times 10^{-5}$	CL=95%	
Γ_{275}	$\gamma\eta(1405) \rightarrow \gamma\gamma\gamma$	< 2.63	$\times 10^{-6}$	CL=90%	
Γ_{276}	$\gamma\eta(1475) \rightarrow \gamma\gamma\gamma$	< 1.86	$\times 10^{-6}$	CL=90%	
Γ_{277}	$\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0$	(1.3 \pm 0.9)	$\times 10^{-4}$		
Γ_{278}	$\gamma\eta(1760) \rightarrow \gamma\omega\omega$	(1.98 \pm 0.33)	$\times 10^{-3}$		
Γ_{279}	$\gamma\eta(1760) \rightarrow \gamma\gamma\gamma$	< 4.80	$\times 10^{-6}$	CL=90%	
Γ_{280}	$\gamma\eta(2225)$	(3.14 \pm 0.50)	$\times 10^{-4}$		
Γ_{281}	$\gamma f_2(1270)$	(1.63 \pm 0.12)	$\times 10^{-3}$	S=1.3	
Γ_{282}	$\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0$	(2.58 \pm 0.60)	$\times 10^{-5}$		
Γ_{283}	$\gamma f_1(1285)$	(6.1 \pm 0.8)	$\times 10^{-4}$		
Γ_{284}	$\gamma f_0(1370) \rightarrow \gamma\pi\pi$				
Γ_{285}	$\gamma f_0(1370) \rightarrow \gamma K\bar{K}$	(4.2 \pm 1.5)	$\times 10^{-4}$		
Γ_{286}	$\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0$	(1.1 \pm 0.4)	$\times 10^{-5}$		
Γ_{287}	$\gamma f_0(1370) \rightarrow \gamma\eta\eta$				
Γ_{288}	$\gamma f_0(1370) \rightarrow \gamma\eta\eta'$				
Γ_{289}	$\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi$	(7.9 \pm 1.3)	$\times 10^{-4}$		
Γ_{290}	$\gamma f_0(1500) \rightarrow \gamma\pi\pi$	(1.09 \pm 0.24)	$\times 10^{-4}$		
Γ_{291}	$\gamma f_0(1500) \rightarrow \gamma\eta\eta$	(1.7 \pm 0.6)	$\times 10^{-5}$		
Γ_{292}	$\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0$	(1.59 \pm 0.24)	$\times 10^{-5}$		
Γ_{293}	$\gamma f_0(1500) \rightarrow \gamma\eta\eta'$				
Γ_{294}	$\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-$	(4.5 \pm 1.2)	$\times 10^{-4}$		
Γ_{295}	$\gamma f'_2(1525)$	(5.7 \pm 0.8)	$\times 10^{-4}$	S=1.5	

Γ_{296}	$\gamma f'_2(1525) \rightarrow \gamma K_S^0 K_S^0$	$(8.0 \pm 0.7) \times 10^{-5}$
Γ_{297}	$\gamma f'_2(1525) \rightarrow \gamma \eta \eta$	$(3.4 \pm 1.4) \times 10^{-5}$
Γ_{298}	$\gamma f_2(1565) \rightarrow \gamma \eta \eta'$	
Γ_{299}	$\gamma f_2(1640) \rightarrow \gamma \omega \omega$	$(2.8 \pm 1.8) \times 10^{-4}$
Γ_{300}	$\gamma f_0(1710) \rightarrow \gamma \pi \pi$	$(3.8 \pm 0.5) \times 10^{-4}$
Γ_{301}	$\gamma f_0(1710) \rightarrow \gamma K \bar{K}$	$(9.5 \pm 1.0) \times 10^{-4}$ S=1.5
Γ_{302}	$\gamma f_0(1710) \rightarrow \gamma \omega \omega$	$(3.1 \pm 1.0) \times 10^{-4}$
Γ_{303}	$\gamma f_0(1710) \rightarrow \gamma \eta \eta$	$(2.4 \pm 1.2) \times 10^{-4}$
Γ_{304}	$\gamma f_0(1710) \rightarrow \gamma \eta \eta'$	
Γ_{305}	$\gamma f_0(1710) \rightarrow \gamma \omega \phi$	$(2.5 \pm 0.6) \times 10^{-4}$
Γ_{306}	$\gamma f_0(1770) \rightarrow \gamma K_S^0 K_S^0$	$(1.11 \pm 0.20) \times 10^{-5}$
Γ_{307}	$\gamma f_2(1810) \rightarrow \gamma \eta \eta$	$(5.4 \pm 3.5) \times 10^{-5}$
Γ_{308}	$\gamma \eta_1(1855) \rightarrow \gamma \eta \eta'$	$(2.7 \pm 0.4) \times 10^{-6}$
Γ_{309}	$\gamma f_0(1770) \rightarrow \gamma \eta \eta'$	
Γ_{310}	$\gamma f_2(1910) \rightarrow \gamma \omega \omega$	$(2.0 \pm 1.4) \times 10^{-4}$
Γ_{311}	$\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892)$	$(7.0 \pm 2.2) \times 10^{-4}$
Γ_{312}	$\gamma f_2(2010) \rightarrow \gamma \eta \eta'$	
Γ_{313}	$\gamma f_0(2020) \rightarrow \gamma \pi \pi$	
Γ_{314}	$\gamma f_0(2020) \rightarrow \gamma K \bar{K}$	
Γ_{315}	$\gamma f_0(2020) \rightarrow \gamma \eta \eta$	
Γ_{316}	$\gamma f_0(2020) \rightarrow \gamma \eta' \eta'$	$(2.63 \pm 0.32) \times 10^{-4}$
Γ_{317}	$\gamma f_0(2020) \rightarrow \gamma \eta \eta'$	
Γ_{318}	$\gamma f_4(2050)$	$(2.7 \pm 0.7) \times 10^{-3}$
Γ_{319}	$\gamma f_4(2050) \rightarrow \gamma \eta \eta'$	
Γ_{320}	$\gamma f_0(2100) \rightarrow \gamma \eta \eta$	$(1.13 \pm 0.30) \times 10^{-4}$
Γ_{321}	$\gamma f_0(2100) \rightarrow \gamma K \bar{K}$	
Γ_{322}	$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	$(6.2 \pm 1.0) \times 10^{-4}$
Γ_{323}	$\gamma f_0(2200)$	seen
Γ_{324}	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(5.9 \pm 1.3) \times 10^{-4}$
Γ_{325}	$\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0$	$(2.72 \pm 0.19) \times 10^{-4}$
Γ_{326}	$\gamma f_0(2200) \rightarrow \gamma \pi \pi$	
Γ_{327}	$\gamma f_0(2200) \rightarrow \gamma \eta \eta$	
Γ_{328}	$\gamma f_J(2220)$	seen
Γ_{329}	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	$< 3.9 \times 10^{-5}$ CL=90%
Γ_{330}	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 4.1 \times 10^{-5}$ CL=90%
Γ_{331}	$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	$(1.5 \pm 0.8) \times 10^{-5}$
Γ_{332}	$\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0$	$(4.9 \pm 0.7) \times 10^{-5}$

Γ_{333}	$\gamma f_0(2330) \rightarrow \gamma\pi\pi$				
Γ_{334}	$\gamma f_0(2330) \rightarrow \gamma\eta\eta$				
Γ_{335}	$\gamma f_0(2330) \rightarrow \gamma\eta'\eta'$	$(6.1 \pm 4.0) \times 10^{-6}$			
Γ_{336}	$\gamma f_0(2330) \rightarrow \gamma\eta\eta'$				
Γ_{337}	$\gamma f_2(2340) \rightarrow \gamma\eta\eta$	$(5.6 \pm 2.4) \times 10^{-5}$			
Γ_{338}	$\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0$	$(5.5 \pm 4.0) \times 10^{-5}$			
Γ_{339}	$\gamma f_2(2340) \rightarrow \gamma\eta'\eta'$	$(8.7 \pm 0.9) \times 10^{-6}$			
Γ_{340}	$\gamma f_0(2470) \rightarrow \gamma\eta'\eta'$	$(8.2 \pm 4.0) \times 10^{-7}$			
Γ_{341}	$\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta'$	$(2.7 \pm 0.6) \times 10^{-4}$	S=1.6		
Γ_{342}	$\gamma X(1835) \rightarrow \gamma p\bar{p}$	$(7.7 \pm 1.5) \times 10^{-5}$			
Γ_{343}	$\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta$	$(3.3 \pm 2.0) \times 10^{-5}$			
Γ_{344}	$\gamma X(1835) \rightarrow \gamma\gamma\phi(1020)$				
Γ_{345}	$\gamma X(1835) \rightarrow \gamma\gamma\gamma$	$< 3.56 \times 10^{-6}$	CL=90%		
Γ_{346}	$\gamma X(1835) \rightarrow \gamma 3(\pi^+\pi^-)$	$(2.4 \pm 0.7) \times 10^{-5}$			
Γ_{347}	$\gamma X(2370) \rightarrow \gamma K^+ K^- \eta'$	$(1.8 \pm 0.7) \times 10^{-5}$			
Γ_{348}	$\gamma X(2370) \rightarrow \gamma K_S^0 K_S^0 \eta'$	$(1.2 \pm 0.5) \times 10^{-5}$			
Γ_{349}	$\gamma X(2370) \rightarrow \gamma\eta\eta\eta'$	$< 9.2 \times 10^{-6}$	CL=90%		
Γ_{350}	$\gamma p\bar{p}$	$(3.8 \pm 1.0) \times 10^{-4}$			
Γ_{351}	$\gamma p\bar{p}\pi^+\pi^-$	$< 7.9 \times 10^{-4}$	CL=90%		
Γ_{352}	$\gamma\Lambda\bar{\Lambda}$	$< 1.3 \times 10^{-4}$	CL=90%		
Γ_{353}	$\gamma A^0 \rightarrow \gamma \text{invisible}$	$[e] < 1.7 \times 10^{-6}$	CL=90%		
Γ_{354}	$\gamma A^0 \rightarrow \gamma\mu^+\mu^-$	$[f] < 7.8 \times 10^{-7}$	CL=90%		

Dalitz decays

Γ_{355}	$\pi^0 e^+ e^-$	$(7.6 \pm 1.4) \times 10^{-7}$		
Γ_{356}	$\eta e^+ e^-$	$(1.42 \pm 0.08) \times 10^{-5}$		
Γ_{357}	$\eta'(958) e^+ e^-$	$(6.59 \pm 0.18) \times 10^{-5}$		
Γ_{358}	$X(1835) e^+ e^-, X \rightarrow \pi^+\pi^-\eta'$	$(3.58 \pm 0.25) \times 10^{-6}$		
Γ_{359}	$X(2120) e^+ e^-, X \rightarrow \pi^+\pi^-\eta'$	$(8.2 \pm 1.3) \times 10^{-7}$		
Γ_{360}	$X(2370) e^+ e^-, X \rightarrow \pi^+\pi^-\eta'$	$(1.08 \pm 0.17) \times 10^{-6}$		
Γ_{361}	$\eta U \rightarrow \eta e^+ e^-$	$[g] < 9.11 \times 10^{-7}$	CL=90%	
Γ_{362}	$\eta'(958) U \rightarrow \eta'(958) e^+ e^-$	$[g] < 2.0 \times 10^{-7}$	CL=90%	
Γ_{363}	$\phi e^+ e^-$	$< 1.2 \times 10^{-7}$	CL=90%	

Weak decays

Γ_{364}	$D^- e^+ \nu_e + \text{c.c.}$	$< 7.1 \times 10^{-8}$	CL=90%	
Γ_{365}	$\overline{D}^0 e^+ e^- + \text{c.c.}$	$< 8.5 \times 10^{-8}$	CL=90%	
Γ_{366}	$D_s^- e^+ \nu_e + \text{c.c.}$	$< 1.3 \times 10^{-6}$	CL=90%	
Γ_{367}	$D_s^{*-} e^+ \nu_e + \text{c.c.}$	$< 1.8 \times 10^{-6}$	CL=90%	

Γ_{368}	$D^- \pi^+ + \text{c.c.}$	< 7.5	$\times 10^{-5}$	CL=90%
Γ_{369}	$\overline{D}^0 \overline{K}^0 + \text{c.c.}$	< 1.7	$\times 10^{-4}$	CL=90%
Γ_{370}	$\overline{D}^0 \overline{K}^{*0} + \text{c.c.}$	< 2.5	$\times 10^{-6}$	CL=90%
Γ_{371}	$D_s^- \pi^+ + \text{c.c.}$	< 1.3	$\times 10^{-4}$	CL=90%
Γ_{372}	$D_s^- \rho^+ + \text{c.c.}$	< 1.3	$\times 10^{-5}$	CL=90%

Charge conjugation (C), Parity (P), Lepton Family number (LF) violating modes

Γ_{373}	$\gamma\gamma$	C	< 2.7	$\times 10^{-7}$	CL=90%
Γ_{374}	$\gamma\phi$	C	< 1.4	$\times 10^{-6}$	CL=90%
Γ_{375}	$e^\pm \mu^\mp$	LF	< 1.6	$\times 10^{-7}$	CL=90%
Γ_{376}	$e^\pm \tau^\mp$	LF	< 7.5	$\times 10^{-8}$	CL=90%
Γ_{377}	$\mu^\pm \tau^\mp$	LF	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{378}	$\Lambda_c^+ e^- + \text{c.c.}$		< 6.9	$\times 10^{-8}$	CL=90%

Other decays

Γ_{379}	invisible	< 7	$\times 10^{-4}$	CL=90%
----------------	-----------	-----	------------------	--------

[a] For $E_\gamma > 100$ MeV.

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

[c] $\Theta(1540)$ is a hypothetical pentaquark state of 1.54 GeV/c² mass and a width of less than 25 MeV/c².

[d] Includes $p\bar{p}\pi^+\pi^-\gamma$ and excludes $p\bar{p}\eta$, $p\bar{p}\omega$, $p\bar{p}\eta'$.

[e] For a narrow state A with mass less than 960 MeV.

[f] For a narrow scalar or pseudoscalar A^0 with mass 0.21–3.0 GeV.

[g] For a dark photon U with mass between 100 and 2100 MeV.

FIT INFORMATION

A multiparticle fit to $\eta_c(1S)$, $J/\psi(1S)$, $\psi(2S)$, $h_c(1P)$, and B^\pm with the total width, 10 combinations of partial widths obtained from integrated cross section, and 38 branching ratios uses 113 measurements to determine 19 parameters. The overall fit has a $\chi^2 = 184.6$ for 94 degrees of freedom.

$J/\psi(1S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$				Γ_1
VALUE (keV)		DOCUMENT ID	TECN	COMMENT
$81.37 \pm 1.36 \pm 1.30$		¹ ANASHIN	20	KEDR $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
74.1 \pm 8.1	BAI	95B	BES	$e^+ e^-$
59 \pm 24	BALDINI...	75	FRAG	$e^+ e^-$
59 \pm 14	BOYARSKI	75	MRK1	$e^+ e^-$

50 ± 25 ESPOSITO 75B FRAM $e^+ e^-$ ¹ Based on the same dataset as ANASHIN 18A and correlated to the values reported there $\Gamma(e^+ e^-)$ Γ_5

<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.53 ± 0.10 OUR AVERAGE				
5.550 $\pm 0.056 \pm 0.089$	1,2	ANASHIN	18A KEDR	$e^+ e^-$
5.36 ± 0.29 -0.28	3	HSUEH	92 RVUE	See γ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.58 $\pm 0.05 \pm 0.08$	4	ABLIKIM	16Q BES3	$3.773 e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
5.71 ± 0.16	13k	5 ADAMS	06A CLEO	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
5.57 ± 0.19	7.8k	5 AUBERT	04 BABR	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
5.14 ± 0.39	BAI	95B BES	$e^+ e^-$	
4.72 ± 0.35	ALEXANDER	89 RVUE	See γ mini-review	
4.4 ± 0.6	3 BRANDELIK	79c DASP	$e^+ e^-$	
4.6 ± 0.8	6 BALDINI-...	75 FRAG	$e^+ e^-$	
4.8 ± 0.6	BOYARSKI	75 MRK1	$e^+ e^-$	
4.6 ± 1.0	ESPOSITO	75B FRAM	$e^+ e^-$	

¹ From the cross sections of $e^+ e^- \rightarrow e^+ e^-$ and $e^+ e^- \rightarrow$ hadrons near the $J/\psi(1S)$ peak.² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.³ From a simultaneous fit to $e^+ e^-$, $\mu^+ \mu^-$, and hadronic channels assuming $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$.⁴ Using $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.973 \pm 0.007 \pm 0.037)\%$ from ABLIKIM 13R.⁵ Calculated by us from the reported values of $\Gamma(e^+ e^-) \times B(\mu^+ \mu^-)$ using $B(\mu^+ \mu^-) = (5.93 \pm 0.06)\%$.⁶ Assuming equal partial widths for $e^+ e^-$ and $\mu^+ \mu^-$. $\Gamma(\mu^+ \mu^-)$ Γ_7

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.13 ± 0.52	BAI	95B BES	$e^+ e^-$
4.8 ± 0.6	BOYARSKI	75 MRK1	$e^+ e^-$
5 ± 1	ESPOSITO	75B FRAM	$e^+ e^-$

 $\Gamma(\gamma\gamma)$ Γ_{373}

<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5.4	90	BRANDELIK	79c DASP	$e^+ e^-$

 $J/\psi(1S) \Gamma(i) \Gamma(e^+ e^-) / \Gamma(\text{total})$

This combination of a partial width with the partial width into $e^+ e^-$ and with the total width is obtained from the integrated cross section into channel(I) in the $e^+ e^-$ annihilation.

 $\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_1 \Gamma_5 / \Gamma$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4.884 $\pm 0.048 \pm 0.078$	1,2 ANASHIN	18A KEDR	$e^+ e^-$

4	± 0.8	³ BALDINI...	75	FRAG	$e^+ e^-$
3.9	± 0.8	³ ESPOSITO	75B	FRAM	$e^+ e^-$

¹ From the cross sections of $e^+ e^- \rightarrow e^+ e^-$ and $e^+ e^- \rightarrow$ hadrons near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

$\Gamma(e^+ e^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_5 \Gamma_5/\Gamma$

VALUE (eV)		DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
333.1 \pm 6.6 \pm 4.0	^{1,2} ANASHIN	18A	KEDR	$e^+ e^-$
332.3 \pm 6.4 \pm 4.8	ANASHIN	10	KEDR	$3.097 e^+ e^- \rightarrow e^+ e^-$
350 \pm 20	BRANDELIK	79C	DASP	$e^+ e^-$
320 \pm 70	³ BALDINI...	75	FRAG	$e^+ e^-$
340 \pm 90	³ ESPOSITO	75B	FRAM	$e^+ e^-$
360 \pm 100	³ FORD	75	SPEC	$e^+ e^-$

¹ From the cross sections of $e^+ e^- \rightarrow e^+ e^-$ and $e^+ e^- \rightarrow$ hadrons near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

$\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_7 \Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
333 \pm 4 OUR AVERAGE				
333.4 \pm 2.5 \pm 4.4	ABLIKIM	16Q	BES3	$3.773 e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
331.8 \pm 5.2 \pm 6.3	ANASHIN	10	KEDR	$3.097 e^+ e^- \rightarrow \mu^+ \mu^-$
338.4 \pm 5.8 \pm 7.1	13k	ADAMS	06A	CLEO $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
330.1 \pm 7.7 \pm 7.3	7.8k	AUBERT	04	BABR $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
510 \pm 90	DASP	75	DASP	$e^+ e^-$
380 \pm 50	¹ ESPOSITO	75B	FRAM	$e^+ e^-$

¹ Data redundant with branching ratios or partial widths above.

$\Gamma(\eta \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{13} \Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 \pm 0.4 OUR AVERAGE				
2.34 \pm 0.43 \pm 0.16	49	LEES	18	BABR $e^+ e^- \rightarrow \eta \pi^+ \pi^- \gamma$
2.22 \pm 0.96 \pm 0.02	9	¹ AUBERT	07AU	BABR $10.6 e^+ e^- \rightarrow \eta \pi^+ \pi^- \gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \eta \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+ \pi^- \pi^0)] = 0.51 \pm 0.22 \pm 0.03$ eV which we divide by our best value $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = (23.02 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta \pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{15} \Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
64.8 \pm 11.1 \pm 0.4	200	¹ LEES	21C	BABR $e^+ e^- \rightarrow \gamma_{ISR} (\pi^+ \pi^- 4\pi^0)$

¹ LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \eta \pi^+ \pi^- \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] = 21.1 \pm 1.7 \pm 3.2$ eV which we divide by our best value $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta\pi^+\pi^-3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{16}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
26.9±5.7±0.1	101	1 LEES	21C BABR	$e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-3\pi^0\gamma\gamma)$
¹ LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 10.6 \pm 1.6 \pm 1.6$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(\eta K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{19}\Gamma_5/\Gamma$			
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
4.76±1.64±0.03	1 LEES	23	BABR	$e^+e^- \rightarrow \gamma_{ISR}$ hadrons
¹ LEES 23 reports $[\Gamma(J/\psi(1S) \rightarrow \eta K^+K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] = 1.55 \pm 0.51 \pm 0.16$ eV which we divide by our best value $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(\eta K^\pm K_S^0 \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{20}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
7.3±1.4±0.4	44	LEES	17D BABR	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$\Gamma(K_S^0 K^\pm \pi^\mp \rho(770)^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{196}\Gamma_5/\Gamma$			
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
17.3±2.1±1.7	LEES	23	BABR	$e^+e^- \rightarrow \gamma_{ISR}$ hadrons

$\Gamma(\rho^\pm \pi^\mp \pi^+ \pi^- 2\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{23}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
155±26±36	14k	LEES	21	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$\Gamma(\rho^+ \rho^- \pi^+ \pi^- \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{24}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
32±13±15	14k	LEES	21	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$\Gamma(\rho^\mp K^\pm K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{26}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.4±1.0±1.9	130	LEES	17D BABR	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{41}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
53.6±5.0±0.4	788	1 AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 47.8 \pm 3.1 \pm 3.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(\omega\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{42}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
27.8±3.5±0.2	398	1 LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$
¹ LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 24.8 \pm 1.8 \pm 2.5$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(\omega 3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{43}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±3.1±0.1	89	1 LEES	21C BABR	$e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$
¹ LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \omega 3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 9.4 \pm 2.3 \pm 1.5$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(\omega\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{46}\Gamma_5/\Gamma$			
VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.2±0.3±0.2	170	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\pi^0\gamma$

$\Gamma(\omega\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{45}\Gamma_5/\Gamma$			
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
16.9±7.6±0.2	1 LEES	21C BABR	$e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$	
¹ Different final state as in AUBERT 06. LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \omega\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 4.9 \pm 2.1 \pm 0.7$ eV which we divide by our best values $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.21) \times 10^{-2}$, $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.				

$\Gamma(\omega\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{47}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.90±0.96±0.01	27	1 LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$
¹ LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 1.7 \pm 0.8 \pm 0.3$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(\omega\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{49}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
185±30±1	14k	1 LEES	21 BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$
¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 165 \pm 9 \pm 25$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$$\Gamma(\omega K\bar{K}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{60}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.70±1.98±0.03	24	1 AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow \omega K^+ K^- \gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega K\bar{K}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 3.3 \pm 1.3 \pm 1.2 \text{ eV}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^+ K^*(892)^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{69}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
29.0±1.7±1.3		AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K^+ K^*(892)^- \gamma$

$$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{70}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.96±0.85±0.70	155	AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^0 \gamma$

$$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{71}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
16.76±1.70±1.00	89	AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{72}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
26.6±2.5±1.5		AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K^0 \bar{K}^*(892)^0 \gamma$

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{73}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
17.70±1.70±1.00	94	AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

$$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{74}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
42.6±4.8±7.2	99	1 LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 1/6 to account for $B(K^*(892)^0 \rightarrow K_S^0 \pi^0) = 1/6$.

$$\Gamma(K^*(892)^\pm K^\mp \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{75}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
22.8±2.8±6.8	80	1 LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 1/4 to account for $B(K^*(892)^\pm \rightarrow K_S^0 \pi^\pm) = 1/4$.

$$\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{76}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.0±2.8 OUR AVERAGE				

9.2±1.2±3.2	64	1 LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$
14.8±4.8±1.2	53	2 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K^0 \gamma$

¹ Dividing by 1/2 to take into account $B(K^*(892)^\pm \rightarrow K^\pm \pi^\mp) = 1/2$.

² Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{77}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.7±1.2±0.3	53	LEES	14H	BABR $e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

$\Gamma(K^*(892)^0 K_S^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{80}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.60±0.75±2.25	34	1 LEES	17D	BABR $e^+ e^- \rightarrow K_S^0 K_S^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 2/3 to account for $B(K^*(892)^0 \rightarrow K^+ \pi^-) = 2/3$.

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{82}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.28±0.34±0.07	47±12	1 LEES	12F	BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.28 \pm 0.40 \pm 0.11 \quad 25 \pm 8 \quad ^{1,2} \text{ AUBERT } 07AK \text{ BABR } 10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Dividing by $(2/3)^2$ to take twice into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3 B(K^{*0} \rightarrow K\pi)$.

² Superseded by LEES 12F.

$\Gamma(K^*(892)^\pm K^*(892)^\mp) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{83}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.80±0.48±0.32	1±5	1 LEES	14H	BABR $e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by $(1/4)^2$ to take twice into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$\Gamma(K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{94}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.1±9.8±0.5	35	1,2 LEES	14H	BABR $e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by 1/4 to take into account $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K\pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 10.0 \pm 4.8 \pm 0.8 \text{ eV}$ which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$	Γ_{93}/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.65±0.80±0.44	1094	ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{95}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
25.8±1.4±0.6	710	1,2,3 LEES	12F	BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$33 \pm 4 \pm 1 \quad 317 \quad ^{2,4} \text{ AUBERT } 07AK \text{ BABR } 10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 12.89 \pm 0.54 \pm 0.41 \text{ eV}$ which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is

their experiment's error and our second error is the systematic error from using our best value.

² Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3 B(K^{*0} \rightarrow K\pi)$.

³ The $K_2^*(1430)$ cannot be distinguished from the $K_0^*(1430)$.

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 16.4 \pm 1.1 \pm 1.4$

eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\frac{\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}}{\Gamma_{96}\Gamma_5/\Gamma}$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
18.6±16.1±0.4	8 ± 8	1,2 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by $(1/4)^2$ to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ and $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4$ $B(K^*(1430) \rightarrow K\pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 9.28 \pm 8.0 \pm 0.32$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\frac{\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}}{\Gamma_{97}\Gamma_5/\Gamma}$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.32±2.00±0.08	8 ± 8	1 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$$\frac{\Gamma(\bar{K}_2(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}}{\Gamma_{99}\Gamma_5/\Gamma}$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.8±0.4±0.3	110 ± 14	1 AUBERT 07AK	BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3$.

$$\frac{\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}}{\Gamma_{106}\Gamma_5/\Gamma}$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.48±0.35 OUR AVERAGE				

$4.46 \pm 0.49 \pm 0.05$	181	1 LEES	12F BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
$4.51 \pm 0.48 \pm 0.05$	254 ± 23	2 SHEN	09 BELL	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.3 \pm 0.7 \pm 0.1$	103	3 AUBERT,BE 06D	BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
-----------------------	-----	-----------------	------	---

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.19 \pm 0.23 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² SHEN 09 reports $4.50 \pm 0.41 \pm 0.26$ eV from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)]$ assuming

$B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.6) \times 10^{-2}$, which we rescale to our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by LEES 12F. AUBERT,BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.61 \pm 0.30 \pm 0.18$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{107}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.77±0.57±0.03	45	¹ LEES	12F BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.13±0.88±0.03	23	² AUBERT,BE	06D BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$	
¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.36 \pm 0.27 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² Superseded by LEES 12F. AUBERT,BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.54 \pm 0.40 \pm 0.16$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(\phi 2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{108}\Gamma_5/\Gamma$
VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.96±0.19±0.01	35	¹ AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \phi 2(\pi^+\pi^-)\gamma$	
¹ AUBERT 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi 2(\pi^+\pi^-)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = (0.47 \pm 0.09 \pm 0.03) \times 10^{-2}$ keV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(\phi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{109}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
4.6±1.4 OUR AVERAGE					
4.1±1.6±0.4		¹ LEES	23 BABR	$e^+e^- \rightarrow \gamma_{ISR}$ hadrons	
6.1±2.7±0.4	6	² AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \phi\eta\gamma$	
¹ LEES 23 quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi^0) = 0.64 \pm 0.26 \pm 0.06$ eV.					
² AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi) = 0.84 \pm 0.37 \pm 0.05$ eV.					

$\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{113}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.44±0.19 OUR AVERAGE					
1.40±0.25±0.01	57 ± 9	¹ LEES	12F BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$	
1.48±0.27±0.09	60 ± 11	² SHEN	09 BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.02 \pm 0.24 \pm 0.01$ 20 ± 5 ³ AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.69 \pm 0.11 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Multiplied by 2/3 to take into account the $\phi \pi^+ \pi^-$ mode only. Using $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$.

³ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.50 \pm 0.11 \pm 0.04$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

$\Gamma_{114} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.98 ± 0.26 ± 0.01	16 ± 4	¹ LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.96 \pm 0.40 \pm 0.01$ 7.0 ± 2.8 ² AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.48 \pm 0.12 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.47 \pm 0.19 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f_2(1270)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

$\Gamma_{119} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.79 ± 0.32 ± 0.02	61	1,2,3 LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.08 \pm 0.73 \pm 0.05$ 44 ^{2,4} AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = 1.51 \pm 0.25 \pm 0.10$ eV which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.3 \pm 2.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

³ Using $\pi^+ \pi^-$ invariant mass between 1.1 and 1.5 GeV. May include other sources such as $f_0(1370)$.

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = 3.44 \pm 0.55 \pm 0.28$ eV which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.3 \pm 2.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f'_2(1525)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{124}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.1±3.2±0.2	11	1,2 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹ Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$ and using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f'_2(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 7.2 \pm 2.8 \pm 0.3$ eV which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{130}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.60±0.62±0.05	163	1 LEES	12F BABR	$10.6 e^+ e^- \rightarrow K^+ K^- K^+ K^- \gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.26 \pm 0.26 \pm 0.16$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi K_S^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{131}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.26±0.84±0.03	29	1 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹ LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi K_S^0 K_S^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.6 \pm 0.4 \pm 0.1$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

/

$\Gamma(f'_2(1525) K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{136}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.8±1.9±0.1	16	1,2 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹ Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow f'_2(1525) K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 5.12 \pm 1.68 \pm 0.20$ eV which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+ \pi^-)\pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{157}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
303±5±18	4990	AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-)\pi^0 \gamma$

$\Gamma(\pi^+\pi^-3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{159}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
100 ± 50 OUR AVERAGE				Error includes scale factor of 4.3.

55 ± 16 ± 1	14k	1 LEES	21 BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$
150.0 ± 4.0 ± 15.0	2.3k	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] = 19.2 \pm 4.5 \pm 3.2 \text{ eV}$ which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}} = 0.3469 \pm 0.0034$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\pi^+\pi^-4\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{162}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
35.8 ± 4.4 ± 5.4	340	LEES	21C BABR	$e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$

 $\Gamma(\rho^\pm\pi^\mp\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{160}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
78.0 ± 9.0 ± 8.0	1.2k	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

 $\Gamma(\rho^+\rho^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{161}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
33.0 ± 5.0 ± 3.3	529	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

 $\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{163}\Gamma_5/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.1248 ± 0.0019 ± 0.0026		LEES	21B BABR	$10.5 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.122 ± 0.005 ± 0.008	AUBERT,B	04N BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
-----------------------	----------	----------	---

 $\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{164}\Gamma_5/\Gamma$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.9 ± 0.5 ± 1.0	761	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$

 $\Gamma(\pi^+\pi^-\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{165}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
107.0 ± 4.3 ± 6.4	768	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\pi^0\gamma$

 $\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{167}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.4 ± 0.9 ± 0.4		LEES	12E BABR	$10.6 e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

19.5 ± 1.4 ± 1.3	270	¹ AUBERT	05D BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)\gamma$
------------------	-----	---------------------	----------	---

¹ Superseded by LEES 12E.

$$\Gamma(3(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{168}\Gamma_5/\Gamma$$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.37±0.16±0.14	496	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow 3(\pi^+\pi^-)\gamma$

$$\Gamma(2(\pi^+\pi^-)3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{169}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
345±10±50	14k	LEES	21	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$$\Gamma(2(\pi^+\pi^-)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{171}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1±2.4±0.1	85	¹ AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)\eta\gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 5.16 \pm 0.85 \pm 0.39$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(2(\pi^+\pi^-\pi^0)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{173}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.1±2.6±1.4	14k	LEES	21	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$$\Gamma(\pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{174}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1± 2.7 OUR AVERAGE				

26.1±17.9±0.3	14k	¹ LEES	21	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$
12.8± 1.8±2.0	203	LEES	18E	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+\pi^-\pi^0)] = 6 \pm 4 \pm 1$ eV which we divide by our best value $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.02 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\rho^\pm\pi^\mp\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{175}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±4.1±1.6	168	LEES	18E	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

$$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{176}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

1.78±0.11±0.05	462	¹ LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
1.94±0.11±0.05	462	² LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
1.42±0.23±0.08	51	³ LEES	13Q	BABR $e^+e^- \rightarrow K^+K^-\gamma$

¹ $\sin\phi > 0$.

² $\sin\phi < 0$.

³ Interference with non-resonant K^+K^- production not taken into account.

$$\Gamma(K_S^0 K_L^0 \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{182}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.4±1.3±0.6	182	LEES	17A	BABR $e^+e^- \rightarrow K_S^0 K_L^0 \pi^0\gamma$

$$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{183} \Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.7 ± 0.9 ± 0.4	106	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

$$\Gamma(K_2^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{184} \Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.4 ± 0.7 ± 0.1	37	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

$$\Gamma(K_S^0 K^*(892)^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{201} \Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
9.45 ± 3.15 ± 0.90	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons

$$\Gamma(K_S^0 K^*(892)^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{202} \Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
5.59 ± 0.79 ± 0.55	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons

$$\Gamma(K^\mp K^*(892)^\pm \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{203} \Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
18.6 ± 6.3 ± 1.8	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons

$$\Gamma(K^\mp K^*(892)^\pm \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{205} \Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
8.67 ± 1.56 ± 0.84	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons

$$\Gamma(K^*(892)^\pm K^*(892)^0 \pi^\mp) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{204} \Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
26.6 ± 4.5 ± 2.7	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons

$$\Gamma(K^*(892)^+ K^*(892)^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{206} \Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
62.1 ± 10.8 ± 6.30	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons

$$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{185} \Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
37.94 ± 0.81 ± 1.10	3.1k	LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

36.3 ± 1.3 ± 2.1	1.5k	¹ AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
33.6 ± 2.7 ± 2.7	233	² AUBERT	05D BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

¹ Superseded by LEES 12F.

² Superseded by AUBERT 07AK.

$$\Gamma(K^+ K^- \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{186} \Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.75 ± 0.81 ± 0.90	388	LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

13.6 ± 1.1 ± 1.3	203	¹ AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$
------------------	-----	---------------------	-----------	---

¹ Superseded by LEES 12F.

$\Gamma(K^+ K^- \pi^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{187} \Gamma_5/\Gamma$			
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
8.9±1.3±0.9	LEES	23	BABR	$e^+ e^- \rightarrow \gamma_{ISR}$ hadrons
$\Gamma(K_S^0 K^\pm \pi^\mp \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{188} \Gamma_5/\Gamma$			
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
29.3±2.6±2.9	LEES	23	BABR	$e^+ e^- \rightarrow \gamma_{ISR}$ hadrons
$\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{189} \Gamma_5/\Gamma$			
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
34.6±1.4±1.8	LEES	23	BABR	$e^+ e^- \rightarrow \gamma_{ISR}$ hadrons
$\Gamma(K_S^0 K_L^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{191} \Gamma_5/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
20.8±2.3±2.1	248	LEES	14H	BABR
				$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_L^0 \gamma$
$\Gamma(K_S^0 K_L^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{192} \Gamma_5/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.3±2.3±0.5	47	LEES	17A	BABR
				$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \pi^0 \gamma$
$\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{193} \Gamma_5/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.0±1.8±0.4	45	LEES	17A	BABR
				$e^+ e^- \rightarrow K_S^0 K_L^0 \eta \gamma$
$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{194} \Gamma_5/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.3±0.9±0.5	133	LEES	14H	BABR
				$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$
$\Gamma(K^\mp K_S^0 \pi^\pm \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{195} \Gamma_5/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
31.7±1.9±1.8	393	LEES	17D	BABR
				$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$
$\Gamma(K_S^0 K^\pm \rho(770)^\pm \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{190} \Gamma_5/\Gamma$			
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
16.0±4.1±1.6	LEES	23	BABR	
				$e^+ e^- \rightarrow \gamma_{ISR}$ hadrons
$\Gamma(K^+ K^- 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{197} \Gamma_5/\Gamma$			
<u>VALUE (10⁻² keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.75±0.23±0.17	205	AUBERT	06D	BABR
				$10.6 e^+ e^- \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$
$\Gamma(K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{198} \Gamma_5/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
25.9±3.9±0.1	73	¹ AUBERT	07AU	BABR
				$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 10.2 \pm 1.3 \pm 0.8$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(K^+ K^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{199}\Gamma_5/\Gamma$


VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.00±0.33±0.29	287 ± 24	LEES	12F BABR	10.6 $e^+ e^- \rightarrow 2(K^+ K^-)\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.11±0.39±0.30	156 ± 15	¹ AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow 2(K^+ K^-)\gamma$
4.0 ± 0.7 ± 0.6	38	² AUBERT	05D BABR	10.6 $e^+ e^- \rightarrow 2(K^+ K^-)\gamma$

¹ Superseded by LEES 12F.² Superseded by AUBERT 07AK.
 $\Gamma(K^+ K^- K_S^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{200}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3±0.4±0.1	29	LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

 $\Gamma(p\bar{p}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{207}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.9±0.6 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.		

11.3±0.4±0.3 821 ¹ LEES 130 BABR $e^+ e^- \rightarrow p\bar{p}\gamma$ 12.9±0.4±0.4 918 ² LEES 13Y BABR $e^+ e^- \rightarrow p\bar{p}\gamma$ 9.7±1.7 ³ ARMSTRONG 93B E760 $\bar{p}p \rightarrow e^+ e^-$ **• • • We do not use the following data for averages, fits, limits, etc. • • •**12.0±0.6±0.5 438 ⁴ AUBERT 06B BABR $e^+ e^- \rightarrow p\bar{p}\gamma$ ¹ ISR photon reconstructed in the detector² ISR photon undetected³ Using $\Gamma_{\text{total}} = 85.5^{+6.1}_{-5.8}$ MeV.⁴ Superseded by LEES 130WEIGHTED AVERAGE
11.9±0.6 (Error scaled by 1.8)

$\Gamma(p\bar{p}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} (\text{eV})$

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$				$\Gamma_{223}\Gamma_5/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
10.7±0.9±0.7	AUBERT	07BD BABR	$10.6 e^+ e^- \rightarrow \Lambda\bar{\Lambda}\gamma$	
$\Gamma(\Sigma^+\bar{\Sigma}^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$				$\Gamma_{233}\Gamma_5/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$6.8 \pm 1.5 \pm 0.8$	GONG	23	BELL $e^+ e^- \rightarrow \Sigma^+\bar{\Sigma}^-$	
$\Gamma(\Sigma^0\bar{\Sigma}^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$				$\Gamma_{234}\Gamma_5/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$5.2 \pm 1.5 \pm 0.6$	GONG	23	BELL $e^+ e^- \rightarrow \Sigma^0\bar{\Sigma}^0$	
$6.4 \pm 1.2 \pm 0.6$	AUBERT	07BD BABR	$10.6 e^+ e^- \rightarrow \Sigma^0\bar{\Sigma}^0\gamma$	

J/ $\psi(1S)$ BRANCHING RATIOS

For the first four branching ratios, see also the partial widths, and (partial widths) $\times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ above.

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.877±0.005 OUR AVERAGE				
0.878 ± 0.005	BAI	95B	BES $e^+ e^-$	
0.86 ± 0.02	BOYARSKI	75	MRK1 $e^+ e^-$	

$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$				Γ_2/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.1346±0.0007	¹ LIAO	23	RVUE $e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.135 ± 0.003	^{2,3} SETH	04	RVUE $e^+ e^-$	
0.17 ± 0.02	² BOYARSKI	75	MRK1 $e^+ e^-$	

¹ Using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (5.967 \pm 0.023)\%$ and $R = 2.26 \pm 0.01$ determined by a fit to data from Mark-I, DM2, BESII, KEDR, and BESIII.

² Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.

³ Using $B(J/\psi \rightarrow \ell^+ \ell^-) = (5.90 \pm 0.09)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C. Superseded by LIAO 23.

$\Gamma(ggg)/\Gamma_{\text{total}}$				Γ_3/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
64.1±1.0	6 M	¹ BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- + \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the PDG 08 values of $B(\ell^+ \ell^-)$, $B(\text{virtual } \gamma \rightarrow \text{hadrons})$, and $B(\gamma \eta_c)$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 08.

$\Gamma(\gamma gg)/\Gamma_{\text{total}}$	Γ_4/Γ				
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
8.79±1.05	200 k	1 BESSON	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma + \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the value of $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ measurement of BESSON 08.

$\Gamma(\gamma gg)/\Gamma(ggg)$	Γ_4/Γ_3				
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
13.7±0.1±0.7	6 M	BESSON	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$	Γ_5/Γ				
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
5.971±0.032 OUR AVERAGE					
5.983±0.007±0.037	720k	ABLIKIM	13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.945±0.067±0.042	15k	LI	05C	CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.90 ± 0.05 ± 0.10		BAI	98D	BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.09 ± 0.33		BAI	95B	BES	$e^+ e^-$
5.92 ± 0.15 ± 0.20		COFFMAN	92	MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ± 0.9		BOYARSKI	75	MRK1	$e^+ e^-$

$\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$	Γ_6/Γ				
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
8.8±1.3±0.4		1 ARMSTRONG	96	E760	$\bar{p}p \rightarrow e^+ e^- \gamma$

¹ For $E_\gamma > 100$ MeV.

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$	Γ_7/Γ				
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
5.961±0.033 OUR AVERAGE					
5.973±0.007±0.038	770k	ABLIKIM	13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.960±0.065±0.050	17k	LI	05C	CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.84 ± 0.06 ± 0.10		BAI	98D	BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.08 ± 0.33		BAI	95B	BES	$e^+ e^-$
5.90 ± 0.15 ± 0.19		COFFMAN	92	MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ± 0.9		BOYARSKI	75	MRK1	$e^+ e^-$

$\Gamma(e^+ e^-)/\Gamma(\mu^+ \mu^-)$	Γ_5/Γ_7			
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.0016±0.0031 OUR AVERAGE				
1.0022±0.0044±0.0048	¹ AULCHENKO 14	KEDR	3.097	$e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
1.0017±0.0017±0.0033	² ABLIKIM 13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
1.002 ± 0.021 ± 0.013	³ ANASHIN 10	KEDR	3.097	$e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
0.997 ± 0.012 ± 0.006	LI 05C	CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.011 ± 0.013 ± 0.016	BAI 98D	BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
1.00 ± 0.07	BAI 95B	BES	$e^+ e^-$	

1.00 ± 0.05	BOYARSKI	75	MRK1	$e^+ e^-$
0.91 ± 0.15	ESPOSITO	75B	FRAM	$e^+ e^-$
0.93 ± 0.10	FORD	75	SPEC	$e^+ e^-$

¹ From 235.3k $J/\psi \rightarrow e^+ e^-$ and 156.6k $J/\psi \rightarrow \mu^+ \mu^-$ observed events.

² Not independent of the corresponding measurements of $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$.


³ Not independent of the corresponding measurements of $\Gamma(e^+ e^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+ \mu^-) \times \Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$.

———— HADRONIC DECAYS ——

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ
1.88 ± 0.12 OUR AVERAGE				Error includes scale factor of 2.6. See the ideogram below.	
2.072 $\pm 0.017 \pm 0.062$	19.8k	¹ ANASHIN	23	KEDR $e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$	
2.18 ± 0.19		^{2,3} AUBERT,B	04N	BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$	
2.184 $\pm 0.005 \pm 0.201$	220k	^{3,4} BAI	04H	BES $e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$	
2.091 $\pm 0.021 \pm 0.116$		^{3,5} BAI	04H	BES $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	
1.21 ± 0.20		BAI	96D	BES $e^+ e^- \rightarrow \rho\pi$	
1.42 $\pm 0.01 \pm 0.19$		COFFMAN	88	MRK3 $e^+ e^-$	
1.3 ± 0.3	150	FRANKLIN	83	MRK2 $e^+ e^-$	
1.6 ± 0.4	183	ALEXANDER	78	PLUT $e^+ e^-$	
1.33 ± 0.21		BRANDELIK	78B	DASP $e^+ e^-$	
1.0 ± 0.2	543	BARTEL	76	CNTR $e^+ e^-$	
1.3 ± 0.3	153	JEAN-MARIE	76	MRK1 $e^+ e^-$	

WEIGHTED AVERAGE
1.88 ± 0.12 (Error scaled by 2.6)



$\Gamma(\rho\pi)/\Gamma_{\text{total}}$ (units 10^{-2})

¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0 \pi^0$, $J/\psi \rightarrow \rho^+ \pi^-$, $J/\psi \rightarrow \rho^- \pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered.

² From the ratio of $\Gamma(e^+ e^-) B(\pi^+ \pi^- \pi^0)$ and $\Gamma(e^+ e^-) B(\mu^+ \mu^-)$ (AUBERT 04).

³ Not independent of their $B(\pi^+ \pi^- \pi^0)$.

⁴ From $J/\psi \rightarrow \pi^+ \pi^- \pi^0$ events directly.

⁵ Obtained comparing the rates for $\pi^+ \pi^- \pi^0$ and $\mu^+ \mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+ \mu^-) = 5.88 \pm 0.10\%$.

$\Gamma(\rho\pi)/\Gamma(\pi^+ \pi^- \pi^0)$

Γ_8/Γ_{163}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.142 ± 0.011 ± 0.026	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.331 ± 0.033	20k	² LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho^0 \pi^0)/\Gamma(\rho\pi)$

Γ_9/Γ_8

VALUE	DOCUMENT ID	TECN	COMMENT
0.328 ± 0.005 ± 0.027	COFFMAN	88	MRK3 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.35 ± 0.08	ALEXANDER	78	PLUT $e^+ e^-$
0.32 ± 0.08	BRANDELIK	78B	DASP $e^+ e^-$
0.39 ± 0.11	BARTEL	76	CNTR $e^+ e^-$
0.37 ± 0.09	JEAN-MARIE	76	MRK1 $e^+ e^-$

$\Gamma(\rho^+ K^+ K^- \pi^- + c.c \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

Γ_{25}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.53 ± 0.16 ± 0.81	485	ANASHIN	22	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(a_2(1320)^0 \pi^+ \pi^- \rightarrow 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$

Γ_{10}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.84 ± 0.08 ± 0.60	1317	ANASHIN	22	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(a_2(1320)^+ \pi^- \pi^0 + c.c \rightarrow 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$

Γ_{11}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.67 ± 0.09 ± 0.73	1628	ANASHIN	22	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(a_2(1320)\rho)/\Gamma_{\text{total}}$

Γ_{12}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
10.9 ± 2.2 OUR AVERAGE				
11.7 ± 0.7 ± 2.5	7584	AUGUSTIN	89 DM2	$J/\psi \rightarrow \rho^0 \rho^\pm \pi^\mp$
8.4 ± 4.5	36	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(\eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{13}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.78 ± 0.68	471	¹ ABLIKIM	19Q BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \eta \pi^+ \pi^-$

¹ From an energy scan of $e^+ e^- \rightarrow J/\psi \rightarrow \eta \pi^+ \pi^-$ assuming PDG 16 values for $\Gamma(e^+ e^-)$, $\Gamma(\mu^+ \mu^-)$, and $\Gamma(\text{total})$.

$\Gamma(\eta\rho)/\Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.193 ± 0.023 OUR AVERAGE				
$0.194 \pm 0.017 \pm 0.029$	299	JOUSSET	90	DM2 $J/\psi \rightarrow$ hadrons
$0.193 \pm 0.013 \pm 0.029$		COFFMAN	88	MRK3 $e^+ e^- \rightarrow \pi^+ \pi^- \eta$

 $\Gamma(\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{17}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.20 \pm 0.14 \pm 0.37$	471	ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

 $\Gamma(\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.52 \times 10^{-4}$	90	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$

 $\Gamma(\eta K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ Γ_{20}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$21.8 \pm 2.2 \pm 3.4$	232 ± 23	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$

 $\Gamma(\eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{21}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.15 \pm 0.13 \pm 0.22$	209	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$

 $\Gamma(\rho\eta'(958))/\Gamma_{\text{total}}$ Γ_{22}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.1 ± 0.8 OUR AVERAGE		Error includes scale factor of 1.6.		
$7.90 \pm 0.19 \pm 0.49$	3476	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$
$8.3 \pm 3.0 \pm 1.2$	19	JOUSSET	90 DM2	$J/\psi \rightarrow$ hadrons
$11.4 \pm 1.4 \pm 1.6$		COFFMAN	88 MRK3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+ \pi^- \eta'$.

 $\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{30}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.2 \pm 0.2 \pm 1.1$	19.8k	¹ ANASHIN	23 KEDR	$e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$

¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0\pi^0$, $J/\psi \rightarrow \rho^+\pi^-$, $J/\psi \rightarrow \rho^-\pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered.

 $\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{30}/Γ_{163}

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				

$10.9 \pm 1.7 \pm 2.7$	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
0.80 ± 0.27	20k	² LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho(1450)^{\pm}\pi^{\mp} \rightarrow K_S^0 K^{\pm}\pi^{\mp})/\Gamma(K_S^0 K^{\pm}\pi^{\mp})$	Γ_{31}/Γ_{181}			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
6.3±0.8±0.6	4k	1 LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^{\pm}\pi^{\mp}$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(\rho(1450)^0\pi^0 \rightarrow K^+ K^- \pi^0)/\Gamma(K^+ K^- \pi^0)$	Γ_{32}/Γ_{180}			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
9.3±2.0±0.6	2k	1 LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(\rho(1450)\eta'(958) \rightarrow \pi^+\pi^-\eta'(958))/\Gamma_{\text{total}}$	Γ_{33}/Γ			
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
3.28±0.55±0.44	119	1 ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.

$\Gamma(\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$	Γ_{35}/Γ_{163}			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8±2±5	20k	1 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

22±6	20k	2 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
------	-----	--------	----------	--------------------------------------

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$	Γ_{37}/Γ_{163}			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4±1±20	20k	1 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

600±250	20k	2 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
---------	-----	--------	----------	--------------------------------------

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.


$\Gamma(\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$	Γ_{38}/Γ_{163}			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

4.0±0.8	20k	1 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
---------	-----	--------	----------	--------------------------------------

¹ From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$	Γ_{39}/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.45 ±0.05 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.		
0.538±0.012±0.065	2090	1 ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\pi^0$
0.360±0.028±0.054	222	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.482±0.019±0.064		COFFMAN	88 MRK3	$e^+e^- \rightarrow \pi^0\pi^+\pi^-\pi^0$

¹ Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.



$$\Gamma(\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8±3±2	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$


¹ From a Dalitz plot analysis in an isobar model and significance 4.9σ .

$$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.5±1.0 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
10.6±1.2±0.1	3531	¹ ANASHIN	22 KEDR	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.0±1.6	18058	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.8±1.6	215	BURMESTER	77D PLUT	e^+e^-
6.8±1.9	348	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

¹ ANASHIN 22 reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (0.946 \pm 0.016 \pm 0.108) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma_{40}/\Gamma_{163}$$

 $\Gamma(\omega\pi^0\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.4 \pm 0.3 \pm 0.7$	509	AUGUSTIN	89	DM2 $J/\psi \rightarrow \pi^+\pi^-3\pi^0$

 Γ_{42}/Γ $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.3 ± 0.6 OUR AVERAGE				
$4.3 \pm 0.2 \pm 0.6$	5860	AUGUSTIN	89	DM2 e^+e^-
4.0 ± 1.6	70	BURMESTER	77D	PLUT e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.9 ± 0.8	81	VANNUCCI	77	MRK1 $e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$


 Γ_{44}/Γ $\Gamma(\omega\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.74 ± 0.20 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
2.352 ± 0.273	5k	¹ ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\eta$
$1.44 \pm 0.40 \pm 0.14$	13	² AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \omega\eta\gamma$
$1.43 \pm 0.10 \pm 0.21$	378	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$1.71 \pm 0.08 \pm 0.20$		COFFMAN	88 MRK3	$e^+e^- \rightarrow 3\pi\eta$

 Γ_{45}/Γ

¹ Using $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$, $B(\eta \rightarrow \pi^+\pi^-\pi^0) = 22.6 \pm 0.4\%$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = 4.68 \pm 0.11\%$, and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

² Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.



$\Gamma(\omega\pi^+\pi^-\pi^-\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
85±34	140	VANNUCCI	77	$e^+e^- \rightarrow 3(\pi^+\pi^-)\pi^0$

Γ_{48}/Γ

$\Gamma(\omega\eta'\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.12±0.02±0.13	14k	1 ABLIKIM	19AC BES3	$J/\psi \rightarrow \omega\eta'\pi^+\pi^-$

Γ_{50}/Γ

$\Gamma(\omega\eta'(958))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.89±0.18 OUR AVERAGE				
2.08±0.30±0.14	137	1 ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$
2.26±0.43	218	2 ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\eta'$
1.8 ± 1.0 ± 0.3	6	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
1.66±0.17±0.19		COFFMAN	88 MRK3	$e^+e^- \rightarrow 3\pi\eta'$

Γ_{51}/Γ

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.

² Using $B(\eta' \rightarrow \pi^+\pi^-\eta) = (44.3 \pm 1.5)\%$, $B(\eta' \rightarrow \pi^+\pi^-\gamma) = 29.5 \pm 1.0\%$, $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$, and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

$\Gamma(\omega f_0(980))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.41±0.27±0.47		1 AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

Γ_{52}/Γ

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

$\Gamma(\omega f_0(1710) \rightarrow \omega K\bar{K})/\Gamma_{\text{total}}$ Γ_{53}/Γ

<u>VALUE</u> (units 10^{-4})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.8±1.1±0.3	1,2 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

¹ Includes unknown branching fraction $f_0(1710) \rightarrow K\bar{K}$.² Addition of $f_0(1710) \rightarrow K^+K^-$ and $f_0(1710) \rightarrow K^0\bar{K}^0$ branching ratios. $\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$ Γ_{54}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.8^{+1.9}_{-1.6}^{±1.7}	111 ⁺³¹ ₋₂₆	BECKER	87 MRK3	$e^+e^- \rightarrow \text{hadrons}$

 $\Gamma(\omega f'_2(1525))/\Gamma_{\text{total}}$ Γ_{55}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.2 × 10⁻⁴	90	1 VANNUCCI	77 MRK1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.8 \times 10^{-4}$	90	1 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
-----------------------	----	-----------	--------	-------------------------------------

¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$. $\Gamma(\omega X(1835) \rightarrow \omega p\bar{p})/\Gamma_{\text{total}}$ Γ_{56}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.9 × 10⁻⁶	95	ABLIKIM	13P BES3	$J/\psi \rightarrow \gamma\pi^0p\bar{p}$

 $\Gamma(\omega X(1835), X \rightarrow \eta'\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{57}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6.2 × 10⁻⁵	1 ABLIKIM	19AC BES3	$J/\psi \rightarrow \omega\eta'\pi^+\pi^-$

¹ Using the decays $\omega \rightarrow \pi^+\pi^-\pi^0$ and $\eta' \rightarrow \eta\pi^+\pi^-$. $\Gamma(\omega K^+K^-)/\Gamma_{\text{total}}$ Γ_{58}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.52±0.30±0.01	276	1 ANASHIN	22 KEDR	$J/\psi \rightarrow K^+K^-\pi^+\pi^-\pi^0$

¹ ANASHIN 22 reports $[\Gamma(J/\psi(1S) \rightarrow \omega K^+K^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (0.136 \pm 0.008 \pm 0.026) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\omega K^\pm K_S^0\pi^\mp)/\Gamma_{\text{total}}$ Γ_{59}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
34 ± 5 OUR AVERAGE				
37.7 \pm 0.8 \pm 5.8	1972 \pm 41	ABLIKIM	08E BES2	$e^+e^- \rightarrow J/\psi$
29.5 \pm 1.4 \pm 7.0	879 \pm 41	BECKER	87 MRK3	$e^+e^- \rightarrow \text{hadrons}$

 $\Gamma(\omega K\bar{K})/\Gamma_{\text{total}}$ Γ_{60}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
19 ± 4 OUR AVERAGE				
19.8 \pm 2.1 \pm 3.9		1 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
16 ± 10	22	FELDMAN	77 MRK1	e^+e^-

¹ Addition of ωK^+K^- and $\omega K^0\bar{K}^0$ branching ratios.

$\Gamma(\omega K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**61 ± 9 OUR AVERAGE**

62.0 ± 6.8 ± 10.6 899 ± 98

65.3 ± 10.2 ± 13.5 176 ± 28

53 ± 14 ± 14 530 ± 140

DOCUMENT ID TECN COMMENTABLIKIM 08E BES2 $J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$ ABLIKIM 08E BES2 $J/\psi \rightarrow \omega K^+ K^- \pi^0$ BECKER 87 MRK3 $e^+ e^- \rightarrow \text{hadrons}$ Γ_{61}/Γ $\Gamma(\eta' K^{*\pm}\bar{K}^\mp)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})**1.48 ± 0.13 OUR AVERAGE**

1.50 ± 0.02 ± 0.19

1.47 ± 0.03 ± 0.17

¹ From $\eta' K^+ K^- \pi^0$.² From $\eta' K_S^0 K^\pm \pi^\mp$. Γ_{62}/Γ DOCUMENT ID TECN COMMENT¹ ABLIKIM 18AB BES3 $J/\psi \rightarrow \eta' K^* \bar{K}$ ² ABLIKIM 18AB BES3 $J/\psi \rightarrow \eta' K^* \bar{K}$ $\Gamma(\eta' K^{*0}\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units 10^{-3})**1.66 ± 0.03 ± 0.21**¹ From $\eta' K_S^0 K^\pm \pi^\mp$. Γ_{63}/Γ DOCUMENT ID TECN COMMENT¹ ABLIKIM 18AB BES3 $J/\psi \rightarrow \eta' K^* \bar{K}$ $\Gamma(\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units 10^{-4})**2.16 ± 0.12 ± 0.29**¹ From $\eta' K_S^0 K^\pm \pi^\mp$. Γ_{64}/Γ DOCUMENT ID TECN COMMENT¹ ABLIKIM 18AB BES3 $J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$ $\Gamma(\eta' h_1(1415) \rightarrow \eta' K^{*\pm} \bar{K}^\mp)/\Gamma_{\text{total}}$ VALUE (units 10^{-4})**1.51 ± 0.09 ± 0.21**¹ From $\eta' K^+ K^- \pi^0$. Γ_{65}/Γ DOCUMENT ID TECN COMMENT¹ ABLIKIM 18AB BES3 $J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$ $\Gamma(\eta' h_1(1415) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$ VALUE (units 10^{-7})**4.69 ± 0.80 ± 0.74**¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta' \eta'$, and ($\eta' X$), with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner. Γ_{66}/Γ DOCUMENT ID TECN COMMENT¹ ABLIKIM 22C BES3 $J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$ $\Gamma(h_1(1415) \eta' \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$ VALUE (units 10^{-5})

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.08 ± 0.01 ^{+0.01} _{-0.02}DOCUMENT ID TECN COMMENT¹ ABLIKIM 22AS BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$ ¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta' P$ -wave.

$\Gamma(h_1(1595)\eta' \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.16 \pm 0.02^{+0.03}_{-0.01}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \eta\eta\eta'$

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta\eta'$ P -wave.

 $\Gamma(\bar{K}K^*(892)+\text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$ Γ_{68}/Γ_{181}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$90.5 \pm 0.9 \pm 3.8$	4k	¹ LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.


 $\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.0^{+0.8}_{-1.0}$ OUR AVERAGE				Error includes scale factor of 2.9. See the ideogram below.

$8.07 \pm 0.04^{+0.38}_{-0.61}$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
$4.57 \pm 0.17 \pm 0.70$	2285	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$5.26 \pm 0.13 \pm 0.53$		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp, K^+ K^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.6 ± 0.6	24	FRANKLIN	83 MRK2	$J/\psi \rightarrow K^+ K^- \pi^0$
3.2 ± 0.6	48	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
4.1 ± 1.2	39	BRAUNSCH...	76 DASP	$J/\psi \rightarrow K^\pm X$

WEIGHTED AVERAGE

6.0+0.8-1.0 (Error scaled by 2.9)


 $\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}} (\text{units } 10^{-3})$

$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{70}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.69 \pm 0.01^{+0.13}_{-0.20}$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

 $\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma(K^+ K^- \pi^0)$ Γ_{70}/Γ_{180}

<u>VALUE</u> (%)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$92.4 \pm 1.5 \pm 3.4$	2k	¹ LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

 $\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{72}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.2 ± 0.4 OUR AVERAGE				
$3.96 \pm 0.15 \pm 0.60$	1192	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$4.33 \pm 0.12 \pm 0.45$		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
2.7 ± 0.6	45	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

 $\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{74}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.73 \pm 0.14 \pm 0.82$		¹ ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				

¹ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

² Seen ABLIKIM 06C BES2 $J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$

¹ Obtained from $J/\psi \rightarrow K^*(892) K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-$ taking the value $2/3$ for the probability of the $K^*(892)^0 \rightarrow K^+ \pi^-$ decay.

² A $K_0^*(700)$ is observed by ABLIKIM 06C in the $K^+ \pi^-$ mass spectrum of the $\bar{K}^*(892)^0 K^+ \pi^-$ final state against the $\bar{K}^*(892)$. A corresponding branching fraction of the $J/\psi(1S)$ is not presented.

 $\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{78}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.81 \pm 0.10 \pm 0.54$	1559	ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{79}/Γ

<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$6.28^{+0.16+0.59}_{-0.17-0.52}$		ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

 $\Gamma(K^*(892)^\pm K^*(700)^\mp)/\Gamma_{\text{total}}$ Γ_{81}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.09 \pm 0.18^{+0.94}_{-0.54}$	655	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{82}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<5	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

$\Gamma(K^*(892)^\pm K^*(892)\mp)/\Gamma_{\text{total}}$					Γ_{83}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.00 \pm 0.19^{+0.11}_{-0.32}$	323	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$	

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$					Γ_{84}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$3.8 \pm 0.8 \pm 1.2$	1 BAI	99C	BES	$e^+ e^-$	

¹ Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

$\Gamma(K^*(1410)\bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0)$					Γ_{86}/Γ_{180}
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$2.3 \pm 1.1 \pm 0.7$	2k	1 LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$	

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K^*(1410)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$					Γ_{87}/Γ_{181}
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.5 \pm 0.5 \pm 0.9$	4k	1 LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$	

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0)$					Γ_{89}/Γ_{180}
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$3.5 \pm 1.3 \pm 0.9$	2k	1 LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$	

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$					Γ_{90}/Γ_{181}
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$7.1 \pm 1.3 \pm 1.2$	4k	1 LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$	

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(\bar{K}_2^*(1430)K + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{91}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<40 \times 10^{-4}$	90	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow K^0 \bar{K}_2^{*0}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<66 \times 10^{-4}$	90	BRAUNSCH... 76	DASP	$e^+ e^- \rightarrow K^\pm \bar{K}_2^{*\mp}$
----------------------	----	----------------	------	--

$\Gamma(K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$					Γ_{92}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$2.69 \pm 0.04^{+0.25}_{-0.19}$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$	

$\Gamma(\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{95}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
6.7 ± 2.6	40	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$	

$\Gamma(K_2^*(1430)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$					Γ_{98}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<29 \times 10^{-4}$	90	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$	

$\Gamma(K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$					Γ_{100}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.1 \pm 0.1^{+0.6}_{-0.1}$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$	

$\Gamma(K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$					Γ_{101}/Γ
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	
$6.2 \pm 0.7^{+2.8}_{-1.4}$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$	

$\Gamma(K_1(1270)^{\pm} K^{\mp})/\Gamma_{\text{total}}$					Γ_{102}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.0 \times 10^{-3}$	90	¹ BAI	99C BES	$e^+ e^-$	

¹ Assuming $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$

$\Gamma(K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$					Γ_{103}/Γ
VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT	
$8.54^{+1.07+2.35}_{-1.20-2.13}$		ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$	

$\Gamma(a_2(1320)^{\pm} \pi^{\mp})/\Gamma_{\text{total}}$					Γ_{104}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<43 \times 10^{-4}$	90	BRAUNSCH... 76	DASP	$e^+ e^-$	

$\Gamma(\phi \pi^0)/\Gamma_{\text{total}}$					Γ_{105}/Γ
The two different fit values of ABLIKIM 15K below have the same statistical significance of 6.4σ and cannot be distinguished at this moment.					
2.94 ± 0.16 ± 0.16	0.8k	¹ ABLIKIM	15K BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma\gamma$	
0.124 $\pm 0.033 \pm 0.030$	35 ± 9	² ABLIKIM	15K BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.4	90	³ ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \gamma\gamma$
<6.8	90	COFFMAN	88 MRK3	$e^+ e^- \rightarrow K^+ K^- \pi^0$

¹ Corresponding to one of the two fit solutions with $\delta = (-95.9 \pm 1.5)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi \pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

² Corresponding to one of the two fit solutions with $\delta = (-152.1 \pm 7.7)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi \pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

³ Superseded by ABLIKIM 15K.

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{106}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.94±0.15 OUR AVERAGE	Error includes scale factor of 1.7.			
1.09±0.02±0.13	ABLIKIM	05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
0.78±0.03±0.12	FALVARD	88	DM2	$J/\psi \rightarrow \text{hadrons}$
2.1 ± 0.9	23	FELDMAN	77	MRK1 e^+e^-

 $\Gamma(\phi 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{108}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
16.0±1.0±3.0	FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$

 $\Gamma(\phi\eta)/\Gamma_{\text{total}}$ Γ_{109}/Γ


<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.74 ±0.06 OUR AVERAGE	Error includes scale factor of 1.2.			
0.71 ± 0.10 ± 0.05	99±14	1 ZHU	23	BELL $e^+e^- \rightarrow \gamma(nS) \rightarrow \phi\eta\gamma$
0.898±0.024±0.089		ABLIKIM	05B	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
0.64 ± 0.04 ± 0.11	346	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
0.661±0.045±0.078		COFFMAN	88	MRK3 $e^+e^- \rightarrow K^+K^-\eta'$

¹ From a fit to the combined $\phi\eta$ invariant mass spectrum with a Gaussian function for the J/ψ signals and a second-order polynomial function for the backgrounds.

 $\Gamma(\phi\eta'(958))/\Gamma_{\text{total}}$ Γ_{110}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.6 ±0.5 OUR AVERAGE	Error includes scale factor of 2.2. See the ideogram below.				
5.10±0.03±0.32	31k	ABLIKIM	16M	BES3	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
5.46±0.31±0.56		ABLIKIM	05B	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
4.1 ± 0.3 ± 0.8	167	JOUSSET	90	DM2	$J/\psi \rightarrow \text{hadrons}$
3.08±0.34±0.36		COFFMAN	88	MRK3	$e^+e^- \rightarrow K^+K^-\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 13	90	VANNUCCI	77	MRK1	e^+e^-

WEIGHTED AVERAGE
4.6±0.5 (Error scaled by 2.2)



$\Gamma(\phi\eta'(958))/\Gamma_{\text{total}} (\text{units } 10^{-4})$

$\Gamma(\phi\eta\eta')/\Gamma_{\text{total}}$	Γ_{111}/Γ				
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2.32±0.06±0.16	2.2k	1 ABLIKIM	19AN BES3	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$	

¹ Including contributions from intermediate resonances. Evidence for an intermediate resonance at $M \approx 2$ GeV and $\Gamma \approx 150$ MeV decaying to $\phi\eta'$ with $J^P = 1^+$ or $J^P = 1^-$, and $B(J/\psi \rightarrow \eta X) \times B(X \rightarrow \phi\eta') \approx 10^{-4}$.

$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$	Γ_{112}/Γ				
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
3.2±0.9 OUR AVERAGE		Error includes scale factor of 1.9.			
4.6±0.4±0.8		1 FALVAR	88 DM2	$J/\psi \rightarrow \text{hadrons}$	
2.6±0.6	50	1 GIDAL	81 MRK2	$J/\psi \rightarrow K^+K^-K^+K^-$	

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

$\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{115}/Γ				
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
4.50±0.80±0.61	355	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+K^-3\pi$	

$\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\rho^0\pi^0)/\Gamma_{\text{total}}$	Γ_{116}/Γ				
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.67±0.50±0.24	70	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+K^-3\pi$	

$\Gamma(\phi f_0(980)\eta \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{117}/Γ				
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
3.23±0.75±0.73	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$	

$\Gamma(\phi a_0(980)^0 \rightarrow \phi\eta\pi^0)/\Gamma_{\text{total}}$	Γ_{118}/Γ				
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
4.37±1.35		1 ABLIKIM	18D BES3	$J/\psi \rightarrow \phi\eta\pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
5.0 ± 2.7 ± 2.5		2 ABLIKIM	11D BES3	$J/\psi \rightarrow \phi\eta\pi^0$	
1 Assuming constructive interference between $a_0(980) - f_0(980)$ mixing and electromagnetic decay. Destructive interference gives a value of $(4.93 \pm 1.77) \times 10^{-6}$ for this branching fraction.					
2 Assuming $a_0(980) - f_0(980)$ mixing and isospin breaking via γ^* and K^*K loops.					

$\Gamma(\phi f_2(1270))/\Gamma_{\text{total}}$	Γ_{119}/Γ				
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.45	90	FALVAR	88 DM2	$J/\psi \rightarrow \text{hadrons}$	
< 0.37	90	VANNUCCI	77 MRK1	$e^+e^- \rightarrow \pi^+\pi^-K^+K^-$	

$\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$	Γ_{120}/Γ				
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2.6±0.5 OUR AVERAGE					
3.4±1.8±1.5	1.1k	1 ABLIKIM	15H BES3	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$	
3.2±0.6±0.4		JOUSSET	90 DM2	$J/\psi \rightarrow \phi 2(\pi^+\pi^-)$	
2.1±0.5±0.4	25	2 JOUSSET	90 DM2	$J/\psi \rightarrow \phi\eta\pi^+\pi^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.6 \pm 0.2 \pm 0.1$ 16 BECKER 87 MRK3 $J/\psi \rightarrow \phi K\bar{K}\pi$

¹ ABLIKIM 15H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_1(1285)) / \Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta\pi^+\pi^-)] = (1.20 \pm 0.6 \pm 0.14) \times 10^{-4}$ which we divide by our best value $B(f_1(1285) \rightarrow \eta\pi^+\pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² We attribute to the $f_1(1285)$ the signal observed in the $\pi^+\pi^-\eta$ invariant mass distribution at 1297 MeV.

$\Gamma(\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-) / \Gamma_{\text{total}}$ Γ_{121}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$9.36 \pm 2.31 \pm 1.54$	78	ABLIKIM	15P	$J/\psi \rightarrow K^+K^-3\pi$

$\Gamma(\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi 3\pi^0) / \Gamma_{\text{total}}$ Γ_{122}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.08 \pm 1.63 \pm 1.47$	9	ABLIKIM	15P	$J/\psi \rightarrow K^+K^-3\pi$

$\Gamma(\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-) / \Gamma_{\text{total}}$ Γ_{123}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.01 \pm 0.58 \pm 0.82$		172	¹ ABLIKIM	15H	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 17 90 ² FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$

¹ With 3.6 σ significance.

² Includes unknown branching fraction $\eta(1405) \rightarrow \eta\pi\pi$.

$\Gamma(\phi f'_2(1525)) / \Gamma_{\text{total}}$ Γ_{124}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8 ± 4 OUR AVERAGE		Error includes scale factor of 2.7.		
$12.3 \pm 0.6 \pm 2.0$		^{1,2} FALVARD	88	$J/\psi \rightarrow \text{hadrons}$
4.8 ± 1.8	46	¹ GIDAL	81	$J/\psi \rightarrow K^+K^-K^+K^-$

¹ Re-evaluated using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.

² Including interference with $f_0(1710)$.

$\Gamma(\phi X(1835) \rightarrow \phi p\bar{p}) / \Gamma_{\text{total}}$ Γ_{125}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.1 \times 10^{-7}$	90	¹ ABLIKIM	16K	$J/\psi \rightarrow p\bar{p}K_S^0K_L^0, p\bar{p}K^+K^-$

¹ Upper limit applies to any $p\bar{p}$ mass enhancement near threshold.

$\Gamma(\phi X(1835) \rightarrow \phi\eta\pi^+\pi^-) / \Gamma_{\text{total}}$ Γ_{126}/Γ


VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.8 \times 10^{-4}$	90	ABLIKIM	15H	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

$\Gamma(\phi X(1870) \rightarrow \phi\eta\pi^+\pi^-) / \Gamma_{\text{total}}$ Γ_{127}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.13 \times 10^{-5}$	90	ABLIKIM	15H	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

$\Gamma(\phi K\bar{K})/\Gamma_{\text{total}}$ Γ_{128}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
17.7 ± 1.6 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
16.6 \pm 1.9 \pm 1.2	163 \pm 19	LEES	12F BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-)\gamma$
21.4 \pm 0.4 \pm 2.2		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
48 \pm 20 \pm 6	9.0 \pm 3.7 \pm 3.0	1,2 HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$
14.6 \pm 0.8 \pm 2.1		3 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
18 \pm 8	14	FELDMAN	77 MRK1	$e^+ e^-$

¹ We have multiplied $K^+ K^-$ measurement by 2 to obtain $K\bar{K}$.² Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.³ Addition of $\phi K^+ K^-$ and $\phi K^0 \bar{K}^0$ branching ratios.WEIGHTED AVERAGE
 17.7 ± 1.6 (Error scaled by 1.3) $\Gamma(\phi f_0(1710) \rightarrow \phi K\bar{K})/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$3.6 \pm 0.2 \pm 0.6$	1,2 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

¹ Including interference with $f'_2(1525)$.² Includes unknown branching fraction $f_0(1710) \rightarrow K\bar{K}$. $\Gamma(\phi K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ Γ_{132}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.2 ± 0.8 OUR AVERAGE				
7.4 \pm 0.6 \pm 1.4	227 \pm 19	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
7.4 \pm 0.9 \pm 1.1		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
7 \pm 0.6 \pm 1.0	163 \pm 15	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\phi K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS
--------------------------	------

 21.8 ± 2.3 OUR AVERAGE

$20.8 \pm 2.7 \pm 3.9$	195 ± 25
$29.6 \pm 3.7 \pm 4.7$	238 ± 30
$20.7 \pm 2.4 \pm 3.0$	
$20 \pm 3 \pm 3$	155 ± 20

 Γ_{133}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$
ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K^+ K^- \pi^0$
FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

 $\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS
--------------------------	------

 30 ± 5 OUR AVERAGE

31 ± 6	4600
29 ± 7	87

 Γ_{134}/Γ

DOCUMENT ID	TECN	COMMENT
AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
BURMESTER	77D PLUT	$e^+ e^-$

 $\Gamma(b_1(1235)^0 \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS
--------------------------	------

 $23 \pm 3 \pm 5$

$<0.1 \times 10^{-3}$	90
-----------------------	----

 Γ_{135}/Γ

DOCUMENT ID	TECN	COMMENT
AUGUSTIN	89 DM2	$e^+ e^-$

 $\Gamma(\Delta(1232)^+ \bar{p})/\Gamma_{\text{total}}$

VALUE	CL%
-------	-----

$<0.1 \times 10^{-3}$	90
-----------------------	----

 Γ_{137}/Γ

DOCUMENT ID	TECN	COMMENT
HENRARD	87 DM2	$e^+ e^-$

 $\Gamma(\Delta(1232)^{++} \bar{p}\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS
--------------------------	------

$1.58 \pm 0.23 \pm 0.40$	332
--------------------------	-------

 Γ_{138}/Γ

DOCUMENT ID	TECN	COMMENT
EATON	84 MRK2	$e^+ e^-$

 $\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS
--------------------------	------

$1.10 \pm 0.09 \pm 0.28$	233
--------------------------	-------

 Γ_{139}/Γ

DOCUMENT ID	TECN	COMMENT
EATON	84 MRK2	$e^+ e^-$

 $\Gamma(\Sigma(1385)^0 p K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS
--------------------------	------

$0.51 \pm 0.26 \pm 0.18$	89
--------------------------	------

 Γ_{140}/Γ

DOCUMENT ID	TECN	COMMENT
EATON	84 MRK2	$e^+ e^-$

 $\Gamma(\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
-------	-----

$<0.82 \times 10^{-5}$	90
------------------------	----

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.2 \times 10^{-3}$	90
-----------------------	----

 Γ_{141}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
HENRARD	87 DM2	$e^+ e^-$

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS
--------------------------	------

 0.30 ± 0.07 OUR AVERAGE

$0.30 \pm 0.03 \pm 0.08$	74 ± 8
$0.29 \pm 0.11 \pm 0.10$	26

 Γ_{142}/Γ

DOCUMENT ID	TECN	COMMENT
HENRARD	87 DM2	$e^+ e^-$
EATON	84 MRK2	$e^+ e^-$

$\Gamma(\Sigma(1385)^+ \bar{\Sigma}^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{143}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.33 ± 0.08 OUR AVERAGE				
$0.34 \pm 0.04 \pm 0.08$	77	HENRARD	87	DM2 $e^+ e^-$
$0.31 \pm 0.11 \pm 0.11$	28	EATON	84	MRK2 $e^+ e^-$

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{144}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.08 ± 0.06 OUR AVERAGE				
$1.096 \pm 0.012 \pm 0.071$	43k	ABLIKIM	16L	BES3 $e^+ e^-$
$1.23 \pm 0.07 \pm 0.30$	0.8k	ABLIKIM	12P	BES2 $e^+ e^-$
$1.00 \pm 0.04 \pm 0.21$	0.6k	HENRARD	87	DM2 $e^+ e^-$
$0.86 \pm 0.18 \pm 0.22$	56	EATON	84	MRK2 $e^+ e^-$

 $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{145}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.25 ± 0.07 OUR AVERAGE				
$1.258 \pm 0.014 \pm 0.078$	53k	ABLIKIM	16L	BES3 $e^+ e^-$
$1.50 \pm 0.08 \pm 0.38$	1k	ABLIKIM	12P	BES2 $e^+ e^-$
$1.19 \pm 0.04 \pm 0.25$	0.7k	HENRARD	87	DM2 $e^+ e^-$
$1.03 \pm 0.24 \pm 0.25$	68	EATON	84	MRK2 $e^+ e^-$

 $\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ Γ_{146}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.071 \pm 0.009 \pm 0.082$	103k	ABLIKIM	17E	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$

 $\Gamma(\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{147}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 4.1 \times 10^{-6}$	90	ABLIKIM	12B	BES3 $J/\psi \rightarrow \Lambda \bar{\Lambda} \gamma$

 $\Gamma(\bar{\Lambda}(1520) \Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{148}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.80 \times 10^{-3}$	90	LU	19	BELL $B^+ \rightarrow \bar{p} \Lambda K^+ K^+$

 $\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$ Γ_{149}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.17 ± 0.04 OUR AVERAGE				
$1.165 \pm 0.004 \pm 0.043$	135k	ABLIKIM	17E	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
$1.20 \pm 0.12 \pm 0.21$	206	ABLIKIM	080	BES2 $e^+ e^- \rightarrow J/\psi$

 $\Gamma(\Xi(1530)^- \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{150}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.318 ± 0.008 OUR AVERAGE				
$0.317 \pm 0.002 \pm 0.008$	70k	ABLIKIM	20	BES3 $e^+ e^- \rightarrow J/\psi$
$0.59 \pm 0.09 \pm 0.12$	75	HENRARD	87	DM2 $e^+ e^-$


$\Gamma(\Xi(1530)^0 \Xi^0)/\Gamma_{\text{total}}$					Γ_{151}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$0.32 \pm 0.12 \pm 0.07$	24 ± 9	HENRARD	87	DM2	$e^+ e^-$
$\Gamma(\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{152}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$< 1.1 \times 10^{-5}$	90	BAI	04G	BES2	$e^+ e^-$
$\Gamma(\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$					Γ_{153}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$< 2.1 \times 10^{-5}$	90	BAI	04G	BES2	$e^+ e^-$
$\Gamma(\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$					Γ_{154}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$< 1.6 \times 10^{-5}$	90	BAI	04G	BES2	$e^+ e^-$
$\Gamma(\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$					Γ_{155}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$< 5.6 \times 10^{-5}$	90	BAI	04G	BES2	$e^+ e^-$
$\Gamma(\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$					Γ_{156}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$< 1.1 \times 10^{-5}$	90	BAI	04G	BES2	$e^+ e^-$

———— STABLE HADRONS ——

$\Gamma(2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$					Γ_{157}/Γ
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
4.2 ± 0.4 OUR AVERAGE	Error includes scale factor of 2.1. See the ideogram below.				
$5.44 \pm 0.07 \pm 0.33$	23K	ANASHIN	22	KEDR	$J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$
$3.5 \pm 0.8 \pm 0.1$	14k	¹ LEES	21	BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-)3\pi^0\gamma$
4.73 ± 0.44	228k	² ABLIKIM	19Q	BES3	$J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$
3.25 ± 0.49	46055	AUGUSTIN	89	DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$
3.17 ± 0.42	147	FRANKLIN	83	MRK2	$e^+ e^- \rightarrow \text{hadrons}$
3.64 ± 0.52	1500	BURMESTER	77D	PLUT	$e^+ e^-$
4 ± 1	675	JEAN-MARIE	76	MRK1	$e^+ e^-$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] \times [\mathcal{B}(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0)] = (14.8 \pm 2.6 \pm 2.2) \times 10^{-3}$ keV which we divide by our best values $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV, $\mathcal{B}(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0) = (18.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² From an energy scan of $e^+ e^- \rightarrow J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$, assuming PDG 16 values for $\Gamma(e^+ e^-)$, $\Gamma(\mu^+ \mu^-)$, and $\Gamma(\text{total})$, and for a phase difference between strong and electromagnetic amplitudes of $(84.9 \pm 3.6)^\circ$. An alternative solution is $(4.85 \pm 0.45)\%$ with a phase of $(-84.7 \pm 3.1)^\circ$.

 $\Gamma(3(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

VALUE	EVTS
0.029±0.006 OUR AVERAGE	
0.028±0.009	11
0.029±0.007	181

 Γ_{158}/Γ

DOCUMENT ID TECN COMMENT

FRANKLIN	83	MRK2	$e^+e^- \rightarrow \text{hadrons}$
JEAN-MARIE	76	MRK1	e^+e^-

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{163}/Γ

VALUE (units 10^{-3})	EVTS
20.0 ±0.7 OUR AVERAGE	
18.78±0.13±0.51	19.8k
21.37±0.04 ^{+0.64} _{-0.62}	1.8M
23.0 ±2.0 ±0.4	256
21.84±0.05±2.01	220k
20.91±0.21±1.16	
15 ±2	168

DOCUMENT ID TECN COMMENT

1 ANASHIN	23	KEDR	$e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$
2 ABLIKIM	12H	BES3	$e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$
3 AUBERT	07AU	BABR	$10.6 e^+e^- \rightarrow J/\psi \pi^+\pi^-\gamma$
4,5 BAI		04H BES	e^+e^-
5,6 BAI		04H BES	e^+e^-
FRANKLIN	83	MRK2	e^+e^-

¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0\pi^0$, $J/\psi \rightarrow \rho^+\pi^-$, $J/\psi \rightarrow \rho^-\pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered.


² The quoted systematic error includes a contribution of 1.23% (added in quadrature) from the uncertainty on the number of J/ψ events.

³ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}} = 0.808 \pm 0.014$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ From $J/\psi \rightarrow \pi^+ \pi^- \pi^0$ events directly.

⁵ Mostly $\rho\pi$, see also $\rho\pi$ subsection.

⁶ Obtained comparing the rates for $\pi^+ \pi^- \pi^0$ and $\mu^+ \mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+ \mu^-) = 5.88 \pm 0.10\%$.



$\Gamma(\pi^+ \pi^- \pi^0 K^+ K^-)/\Gamma_{\text{total}}$

Γ_{165}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.52±0.27 OUR AVERAGE Error includes scale factor of 1.4.				
1.74±0.08±0.24	2616	ANASHIN	22	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.2 ± 0.3	309	VANNUCCI	77	MRK1 $e^+ e^-$

$\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{166}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.47±0.14 OUR AVERAGE				
1.47±0.13±0.13	140	1 METREVELI	12	$\psi(2S) \rightarrow 2(\pi^+ \pi^-)$
1.58±0.20±0.15	84	BALTRUSAIT..85D	MRK3	$e^+ e^-$
1.0 ± 0.5	5	BRANDELIK	78B	DASP $e^+ e^-$
1.6 ± 1.6	1	VANNUCCI	77	MRK1 $e^+ e^-$

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{167}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.20±0.25 OUR AVERAGE				Error includes scale factor of 1.2.
2.88±0.14±0.24	2654	ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+\pi^-)$
3.53±0.12±0.29	1107	¹ ABLIKIM	05H	BES2 $e^+e^- \rightarrow \psi(2S) \rightarrow J/\psi\pi^+\pi^-, J/\psi \rightarrow 2(\pi^+\pi^-)$
4.0 ± 1.0	76	JEAN-MARIE	76	MRK1 e^+e^-

¹ Computed using $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

 $\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{168}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
40±20	32	JEAN-MARIE	76	MRK1 e^+e^-

 $\Gamma(4(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{170}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
90±30	13	JEAN-MARIE	76	MRK1 e^+e^-

 $\Gamma(2(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}$ Γ_{171}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.29±0.28 OUR AVERAGE				
3.1 ± 1.5 ± 0.1	14k	¹ LEES	21	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$
2.26±0.08±0.27	4.8k	ABLIKIM	05C	BES2 $e^+e^- \rightarrow 2(\pi^+\pi^-)\eta$
¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] \times [B(\eta \rightarrow 3\pi^0)] = (5.6 \pm 2.6 \pm 0.8) \times 10^{-3}$ keV which we divide by our best values $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.53 \pm 0.10$ keV, $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.				

 $\Gamma(3(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}$ Γ_{172}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.24±0.96±1.11	616	ABLIKIM	05C	BES2 $e^+e^- \rightarrow 3(\pi^+\pi^-)\eta$


 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{176}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.86±0.09±0.19	1k	¹ METREVELI	12	$\psi(2S) \rightarrow \pi^+\pi^- K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.39±0.24±0.22	107	² BALTRUSAIT..85D	MRK3	e^+e^-
2.2 ± 0.9	6	² BRANDELIK	79C	DASP e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Interference with non-resonant K^+K^- production not taken into account.

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$	Γ_{177}/Γ
<i>VALUE (units 10^{-4})</i>	<i>EVTS</i>
1.95 ± 0.11 OUR AVERAGE	Error includes scale factor of 2.4. See the ideogram below.
1.93 $\pm 0.01 \pm 0.05$	110k
ABLIKIM	17AH BES3
$J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$	
2.62 $\pm 0.15 \pm 0.14$	0.3k
¹ METREVELI	12
$\psi(2S) \rightarrow \pi^+ \pi^- K_S^0 K_L^0$	
1.82 $\pm 0.04 \pm 0.13$	2.1k
² BAI	04A BES2
$J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
1.18 $\pm 0.12 \pm 0.18$	JOUSSET 90 DM2 $J/\psi \rightarrow \text{hadrons}$
1.01 $\pm 0.16 \pm 0.09$	74 BALTRUSAIT..85D MRK3 $e^+ e^-$

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.² Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6868 \pm 0.0027$.

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$	Γ_{178}/Γ
<i>VALUE</i>	<i>CL%</i>
$<1.4 \times 10^{-8}$	95
¹ ABLIKIM	17AH BES3
$J/\psi \rightarrow K_S^0 K_S^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$<1 \times 10^{-6}$	95
¹ BAI	04D BES $e^+ e^-$
$<5.2 \times 10^{-6}$	90
¹ BALTRUSAIT..85C	MRK3 $e^+ e^-$

¹ Forbidden by CP.

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>
61 ± 10 OUR AVERAGE	
55.2 ± 12.0	25
78.0 ± 21.0	126

 Γ_{179}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
FRANKLIN	83	MRK2 $e^+ e^- \rightarrow K^+ K^- \pi^0$
VANNUCCI	77	MRK1 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>
2.88 ± 0.01 ± 0.12	183k

 Γ_{180}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

 $\Gamma(K^+ K^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$

<u>VALUE (%)</u>	<u>EVTS</u>
12.0 ± 0.3 ± 0.9	23k

 $\Gamma_{180}/\Gamma_{163}$

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
LEES	17C BABR	$J/\psi \rightarrow h^+ h^- \pi^0$

 $\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma(\pi^+ \pi^- \pi^0)$

<u>VALUE (%)</u>	<u>EVTS</u>
26.5 ± 0.5 ± 2.1	24k

 $\Gamma_{181}/\Gamma_{163}$

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
LEES	17C BABR	$J/\psi \rightarrow h^0 h^+ h^-$

 $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>
7.04 ± 0.26 ± 0.92	2671

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.2 ± 2.3	205	VANNUCCI	77	MRK1	$e^+ e^-$
-----------	-----	----------	----	------	-----------

 Γ_{185}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>
31 ± 13	30

 Γ_{197}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
VANNUCCI	77	MRK1 $e^+ e^-$

 $\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.4 ± 0.5 -0.4 ± 0.2	11.0 ± 4.3 -3.5	1 HUANG	03 BELL	$B^+ \rightarrow 2(K^+ K^-) K^+$

0.7 ± 0.3	VANNUCCI	77	MRK1	$e^+ e^-$
-----------	----------	----	------	-----------

¹ Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>
2.120 ± 0.029 OUR AVERAGE	

 Γ_{207}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
ABLIKIM	12C BES3	$e^+ e^-$
¹ WU	06 BELL	$B^+ \rightarrow p\bar{p} K^+$
BAI	04E BES2	$e^+ e^- \rightarrow J/\psi$
BALDINI	98 FENI	$e^+ e^-$
PALLIN	87 DM2	$e^+ e^-$
EATON	84 MRK2	$e^+ e^-$
BRANDELIK	79C DASP	$e^+ e^-$
BESCH	78 BONA	$e^+ e^-$
² PERUZZI	78 MRK1	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.3 48 ANTONELLI 93 SPEC $e^+ e^-$

¹ WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assuming angular distribution $(1+\cos^2\theta)$.

$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$

Γ_{208}/Γ


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.19 ± 0.08 OUR AVERAGE	Error includes scale factor of 1.1.			
1.33 $\pm 0.02 \pm 0.11$	11k	ABLIKIM	09B	BES2 $e^+ e^-$
1.13 $\pm 0.09 \pm 0.09$	685	EATON	84	MRK2 $e^+ e^-$
1.4 ± 0.4		BRANDELIK	79C	DASP $e^+ e^-$
1.00 ± 0.15	109	PERUZZI	78	MRK1 $e^+ e^-$

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{209}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ± 0.5 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
6.46 $\pm 0.17 \pm 0.43$	1435	EATON	84	MRK2 $e^+ e^-$
3.8 ± 1.6	48	BESCH	81	BONA $e^+ e^-$
5.5 ± 0.6	533	PERUZZI	78	MRK1 $e^+ e^-$

WEIGHTED AVERAGE
6.0 ± 0.5 (Error scaled by 1.3)



$\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{210}/Γ

Including $p\bar{p}\pi^+\pi^-\gamma$ and excluding ω, η, η'

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.9 OUR AVERAGE	Error includes scale factor of 1.9.			
$3.36 \pm 0.65 \pm 0.28$	364	EATON	84	MRK2 e^+e^-
1.6 ± 0.6	39	PERUZZI	78	MRK1 e^+e^-

$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{211}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00 ± 0.12 OUR AVERAGE				
$1.91 \pm 0.02 \pm 0.17$	13k	¹ ABLIKIM	09	BES2 e^+e^-
$2.03 \pm 0.13 \pm 0.15$	826	EATON	84	MRK2 e^+e^-
2.5 ± 1.2		BRANDELIK	79c	DASP e^+e^-
2.3 ± 0.4	197	PERUZZI	78	MRK1 e^+e^-


¹ From the combination of $p\bar{p}\eta \rightarrow p\bar{p}\gamma\gamma$ and $p\bar{p}\eta \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ channels.

$\Gamma(p\bar{p}\rho)/\Gamma_{\text{total}}$ Γ_{212}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.31 \times 10^{-3}$	90	EATON	84	MRK2 $e^+e^- \rightarrow \text{hadrons}\gamma$

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ Γ_{213}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.98 ± 0.10 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
$0.90 \pm 0.02 \pm 0.09$	2670	ABLIKIM	13P	BES3 e^+e^-
$0.98 \pm 0.03 \pm 0.14$	2449	ABLIKIM	08	BES2 e^+e^-
$1.10 \pm 0.17 \pm 0.18$	486	EATON	84	MRK2 e^+e^-
1.6 ± 0.3	77	PERUZZI	78	MRK1 e^+e^-



$\Gamma(p\bar{p}\eta'(958))/\Gamma_{\text{total}}$ Γ_{214}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.129±0.014 OUR AVERAGE		Error includes scale factor of 2.0.		
0.126±0.002±0.007	16k	¹ ABLIKIM	19N	BES3 $e^+ e^-$
0.200±0.023±0.028	265 ± 31	² ABLIKIM	09	BES2 $e^+ e^-$
0.68 ± 0.23 ± 0.17	19	EATON	84	MRK2 $e^+ e^-$
1.8 ± 0.6	19	PERUZZI	78	MRK1 $e^+ e^-$

¹ From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$ channels.² From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\gamma\rho^0$ channels. $\Gamma(p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta)/\Gamma_{\text{total}}$ Γ_{215}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.8±1.2±1.3	ABLIKIM	14N	$e^+ e^- \rightarrow J/\psi$

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{216}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.519±0.033 OUR AVERAGE				
0.523±0.006±0.033	14k	ABLIKIM	16K	BES3 $J/\psi \rightarrow p\bar{p}K_S^0 K_L^0$, $p\bar{p}K^+ K^-$
0.45 ± 0.13 ± 0.07		FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$

 $\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{217}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.12±0.09 OUR AVERAGE				
2.36±0.02±0.21	59k	ABLIKIM	06K	$J/\psi \rightarrow p\pi^-\bar{n}$
2.47±0.02±0.24	55k	ABLIKIM	06K	$J/\psi \rightarrow \bar{p}\pi^+ n$
2.02±0.07±0.16	1288	EATON	84	MRK2 $e^+ e^- \rightarrow p\pi^-$
1.93±0.07±0.16	1191	EATON	84	MRK2 $e^+ e^- \rightarrow \bar{p}\pi^+$
1.7 ± 0.7	32	BESCH	81	BONA $e^+ e^- \rightarrow p\pi^-$
1.6 ± 1.2	5	BESCH	81	BONA $e^+ e^- \rightarrow \bar{p}\pi^+$
2.16±0.29	194	PERUZZI	78	MRK1 $e^+ e^- \rightarrow p\pi^-$
2.04±0.27	204	PERUZZI	78	MRK1 $e^+ e^- \rightarrow \bar{p}\pi^+$

 $\Gamma(n\bar{n})/\Gamma_{\text{total}}$ Γ_{218}/Γ


<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.09±0.16 OUR AVERAGE				
2.07±0.01±0.17	36k	ABLIKIM	12C	$e^+ e^-$
2.31±0.49	79	BALDINI	98	FENI $e^+ e^-$
1.8 ± 0.9		BESCH	78	BONA $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.90±0.55	40	ANTONELLI	93	SPEC $e^+ e^-$

 $\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{219}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.8±3.6	5	BESCH	81	BONA $e^+ e^-$

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$	Γ_{223}/Γ			
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.88 ± 0.08 OUR AVERAGE		Error includes scale factor of 2.6. See the ideogram below.		
1.64 ± 0.12 ± 0.09		ABLIKIM 23S	BES3	$e^+ e^- \rightarrow \gamma \Lambda\bar{\Lambda}$
1.943 ± 0.003 ± 0.033	441k	ABLIKIM 17L	BES3	$e^+ e^-$
2.03 ± 0.03 ± 0.15	8887	ABLIKIM 06	BES2	$J/\psi \rightarrow \Lambda\bar{\Lambda}$
1.96 $^{+0.47}_{-0.44}$ ± 0.04	46	¹ WU	06	$B^+ \rightarrow \Lambda\bar{\Lambda} K^+$
1.08 ± 0.06 ± 0.24	631	BAI	98G	BES $e^+ e^-$
1.38 ± 0.05 ± 0.20	1847	PALLIN	87	DM2 $e^+ e^-$
1.58 ± 0.08 ± 0.19	365	EATON	84	MRK2 $e^+ e^-$
2.6 ± 1.6	5	BESCH	81	BONA $e^+ e^-$
1.1 ± 0.2	196	PERUZZI	78	MRK1 $e^+ e^-$

¹ WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] = (2.00^{+0.34}_{-0.29} \pm 0.34) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$	Γ_{224}/Γ				
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.78 ± 0.27 ± 0.30		323	¹ ABLIKIM	13F	BES3 $J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.4	90	² ABLIKIM	07H	BES2	$e^+ e^- \rightarrow \psi(2S)$
23 ± 7	± 8	11	BAI	98G	$e^+ e^-$
22 ± 5	± 5	19	HENRARD	87	DM2 $e^+ e^-$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

$\Gamma(\Lambda \bar{\Lambda} \pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{225}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.30 $\pm 0.13 \pm 0.99$	2.4k	ABLIKIM	12P	BES2 J/ψ

$\Gamma(\Lambda \bar{\Lambda} \eta)/\Gamma_{\text{total}}$

Γ_{226}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
16.2 ± 1.7 OUR AVERAGE				
15.7 $\pm 0.80 \pm 1.54$	454	¹ ABLIKIM	13F	BES3 $J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

26.2 $\pm 6.0 \pm 4.4$ 44 ² ABLIKIM 07H BES2 $e^+ e^- \rightarrow \psi(2S)$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

$\Gamma(\Lambda \bar{\Sigma}^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{227}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.26 ± 0.05 OUR AVERAGE		Error includes scale factor of 1.2.		
1.244 $\pm 0.002 \pm 0.045$	2.6M	ABLIKIM	23BUBES3	$e^+ e^-$
1.52 $\pm 0.08 \pm 0.16$	589	¹ ABLIKIM	07H BES2	$e^+ e^-$
1.11 $\pm 0.06 \pm 0.20$	342 ± 18	HENRARD	87 DM2	$e^+ e^-$
1.38 $\pm 0.21 \pm 0.35$	118	EATON	84 MRK2	$e^+ e^-$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$.

$\Gamma(\Lambda \bar{\Sigma}^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{228}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.21 ± 0.07 OUR AVERAGE		Error includes scale factor of 1.8.		
1.221 $\pm 0.002 \pm 0.038$	2.7M	ABLIKIM	23BU BES3	$e^+ e^-$
0.90 $\pm 0.06 \pm 0.16$	225	HENRARD	87 DM2	$e^+ e^-$

1.53 $\pm 0.17 \pm 0.38$ 135 EATON 84 MRK2 $e^+ e^-$

$\Gamma(p K^- \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{229}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.86 ± 0.11 OUR AVERAGE				
0.84 $^{+0.17}_{-0.15} \pm 0.02$	45	¹ LU	19 BELL	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$
0.89 $\pm 0.07 \pm 0.14$	307	EATON	84 MRK2	$e^+ e^-$

¹ LU 19 reports $(8.32^{+1.63}_{-1.45} \pm 0.49) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow p K^- \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)]$ assuming $B(B^+ \rightarrow J/\psi(1S) K^+) = (1.026 \pm 0.031) \times 10^{-3}$, which we rescale to our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(pK^-\bar{\Sigma}^0)/\Gamma_{\text{total}}$		Γ_{230}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.29 \pm 0.06 \pm 0.05$	90	EATON	84	MRK2 $e^+ e^-$

$\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$		Γ_{231}/Γ		
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$6.46 \pm 0.20 \pm 1.07$	1058	¹ ABLIKIM	08C	BES2 $e^+ e^- \rightarrow J/\psi$

¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

$\Gamma(\Lambda\bar{\Sigma} + \text{c.c.})/\Gamma_{\text{total}}$		Γ_{232}/Γ			
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.83 ± 0.23 OUR AVERAGE					
2.74 $\pm 0.24 \pm 0.22$		234 \pm 21	¹ ABLIKIM	12B	BES3 $J/\psi \rightarrow \Lambda\bar{\Sigma}^0$
2.92 $\pm 0.22 \pm 0.24$		308 \pm 24	² ABLIKIM	12B	BES3 $J/\psi \rightarrow \bar{\Lambda}\Sigma^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<18		² HENRARD	87	DM2	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$
<15	90	PERUZZI	78	MRK1	$e^+ e^- \rightarrow \Lambda X$

¹ ABLIKIM 12B quotes $B(J/\psi \rightarrow \Lambda\bar{\Sigma}^0)$ which we multiply by 2.

² ABLIKIM 12B and HENRARD 87 quote results for $B(J/\psi \rightarrow \bar{\Lambda}\Sigma^0)$ which we multiply by 2.

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$		Γ_{233}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.07 ± 0.04 OUR AVERAGE				
1.061 $\pm 0.004 \pm 0.036$	87k	ABLIKIM	21AT	BES3 $J/\psi \rightarrow p\pi^0\bar{p}\pi^0$
1.50 $\pm 0.10 \pm 0.22$	399	ABLIKIM	080	BES2 $e^+ e^- \rightarrow J/\psi$


$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$		Γ_{234}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.172 ± 0.032 OUR AVERAGE				Error includes scale factor of 1.4.
1.164 $\pm 0.004 \pm 0.023$	111k	ABLIKIM	17L	BES3 $J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.33 $\pm 0.04 \pm 0.11$	1.7k	ABLIKIM	06	BES2 $J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.06 $\pm 0.04 \pm 0.23$	884	PALLIN	87	DM2 $e^+ e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.58 $\pm 0.16 \pm 0.25$	90	EATON	84	MRK2 $e^+ e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.3 ± 0.4	52	PERUZZI	78	MRK1 $e^+ e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.4 ± 2.6	3	BESCH	81	BONA $e^+ e^- \rightarrow \Sigma^+\bar{\Sigma}^-$

$\Gamma(\Sigma^+\bar{\Sigma}^-\eta)/\Gamma_{\text{total}}$		Γ_{235}/Γ		
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$6.34 \pm 0.21 \pm 0.37$	1821	ABLIKIM	22AY	BES3 $J/\psi \rightarrow \Sigma^+\bar{\Sigma}^-\eta$

$$\Gamma(\Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}$$

$$\Gamma_{236}/\Gamma$$

<i>VALUE</i> (units 10^{-3})	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
0.97 \pm 0.08 OUR AVERAGE		Error includes scale factor of 1.4.		See the ideogram below.
1.040 \pm 0.006 \pm 0.074	43k	ABLIKIM	16L	BES3 $J/\psi \rightarrow \Xi^- \bar{\Xi}^+$
0.90 \pm 0.03 \pm 0.18	961	ABLIKIM	12P	BES2 $J/\psi \rightarrow \Xi^- \bar{\Xi}^+$
0.70 \pm 0.06 \pm 0.12	132	HENRARD	87	DM2 $e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$
1.14 \pm 0.08 \pm 0.20	194	EATON	84	MRK2 $e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$
1.4 \pm 0.5	51	PERUZZI	78	MRK1 $e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$



$$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$$

Γ₂₃₇/Γ

<u>VALUE</u> (units 10^{-2})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.41 ± 0.14 OUR FIT	Error includes scale factor of 1.3.		
1.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 1.5.		
2.00 ± 0.31 ± 0.02	¹ MITCHELL	09	CLEO $e^+ e^- \rightarrow \gamma X$
1.27 ± 0.36	GAISER	86	CBAL $J/\psi \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3.40 ± 0.33	² ANASHIN	14	KEDR $J/\psi \rightarrow \gamma \eta_c$

¹ MITCHELL 09 reports $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Statistical uncertainty only.

$\Gamma(3\gamma)/\Gamma_{\text{total}}$ Γ_{240}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
11.6±2.2 OUR AVERAGE					
11.3±1.8±2.0		113 ± 18	ABLIKIM	13I	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
12 ± 3 ± 2		24.2 ^{+7.2} _{-6.0}	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<55		90	PARTRIDGE	80	CBAL $e^+ e^-$

 $\Gamma(4\gamma)/\Gamma_{\text{total}}$ Γ_{241}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<9 \times 10^{-6}$	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

 $\Gamma(5\gamma)/\Gamma_{\text{total}}$ Γ_{242}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<15 \times 10^{-6}$	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

 $\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ Γ_{243}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.39±0.08 OUR AVERAGE				
3.34±0.02±0.09	176k	ABLIKIM	23BD	BES3 $J/\psi \rightarrow \pi^0 \gamma$
3.59±0.20±0.04	1.6k	ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$
3.63±0.36±0.13		PEDLAR	09	CLE3 $J/\psi \rightarrow \pi^0 \gamma$
3.13 ^{+0.65} _{-0.47}	586	ABLIKIM	06E	BES2 $J/\psi \rightarrow \pi^0 \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.6 ± 1.1 ± 0.7	BLOOM	83	CBAL	$e^+ e^-$
7.3 ± 4.7	BRANDELIK	79C	DASP	$e^+ e^-$

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] = (3.57 \pm 0.12 \pm 0.16) \times 10^{-5}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\pi^0 \rightarrow 2\gamma) = (98.823 \pm 0.034) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

 $\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{244}/Γ

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.15±0.05	1 ABLIKIM	15AE	BES3 $J/\psi \rightarrow \gamma\pi^0\pi^0$


¹ The uncertainty is systematic as statistical is negligible.

 $\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_{245}/Γ

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.8 ± 0.5 OUR AVERAGE Error includes scale factor of 1.9. See the ideogram below.			
4.32±0.14±0.73	1 BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
2.08±0.13±0.35	2 BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
3.05±0.08±0.45	2 BALTRUSAIT..	86B	MRK3 $J/\psi \rightarrow 4\pi\gamma$
4.85±0.45±1.20	3 BURKE	82	MRK2 $e^+ e^-$

¹ 4π mass less than 3.0 GeV.

² 4π mass less than 2.0 GeV.
³ 4π mass less than 2.5 GeV.



$\Gamma(\gamma f_2(1270) f_2(1270))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$9.5 \pm 0.7 \pm 1.6$	646 ± 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

Γ_{246}/Γ

$\Gamma(\gamma f_2(1270) f_2(1270) (\text{non resonant}))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$8.2 \pm 0.8 \pm 1.7$	¹ ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

Γ_{247}/Γ

¹ Subtracting contribution from intermediate $\eta_c(1S)$ decays.

$\Gamma(\gamma \pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$8.3 \pm 0.2 \pm 3.1$	¹ BALTRUSAIT...86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$

Γ_{248}/Γ

¹ 4π mass less than 2.0 GeV.

$\Gamma(\gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.1 ± 0.4	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

Γ_{249}/Γ

$\Gamma(\gamma(K\bar{K}\pi)[J^P C = 0^- +])/\Gamma_{\text{total}}$ Γ_{250}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 2.1.		
0.58 ± 0.03 ± 0.20	¹ BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
2.1 ± 0.1 ± 0.7	² BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

¹ For a broad structure around 1800 MeV.² For a broad structure around 2040 MeV.
 $\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{251}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.1 ± 0.1 ± 0.6	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

 $\Gamma(\gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{252}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.0 ± 0.3 ± 1.3	320	¹ BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

¹ Summed over all charges.
 $\Gamma(\gamma \eta)/\Gamma_{\text{total}}$ Γ_{253}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.090 ± 0.013 OUR AVERAGE				

1.096 ± 0.001 ± 0.019	2.2M	ABLIKIM	23BD BES3	$J/\psi \rightarrow \eta \gamma$
1.067 ± 0.005 ± 0.023	87.9k	ABLIKIM	21AM BES3	$e^+ e^- \rightarrow J/\psi$
1.12 ± 0.05 ± 0.01	18.6k	¹ ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$
1.101 ± 0.029 ± 0.022		PEDLAR	09 CLE3	$J/\psi \rightarrow \eta \gamma$
1.123 ± 0.089	11k	ABLIKIM	06E BES2	$J/\psi \rightarrow \eta \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.88 ± 0.08 ± 0.11	BLOOM	83 CBAL	$e^+ e^-$
0.82 ± 0.10	BRANDELIK	79C DASP	$e^+ e^-$
1.3 ± 0.4	BARTEL	77 CNTR	$e^+ e^-$

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma \eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = (4.42 \pm 0.04 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma \eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

 $\Gamma(\gamma \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{254}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
21.4 ± 1.8 ± 2.5	596	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$

 $\Gamma(\gamma a_0(980)^0 \rightarrow \gamma \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{258}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2.5 \times 10^{-6}$	95	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$

 $\Gamma(\gamma a_2(1320)^0 \rightarrow \gamma \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{259}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 6.6 \times 10^{-6}$	95	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$

$\Gamma(\gamma\eta\pi\pi)/\Gamma_{\text{total}}$ Γ_{260}/Γ VALUE (units 10^{-3})**6.1 ± 1.0 OUR AVERAGE** $5.85 \pm 0.3 \pm 1.05$ $7.8 \pm 1.2 \pm 2.4$

1 Broad enhancement at 1700 MeV.

DOCUMENT ID

TECN

COMMENT

¹	EDWARDS	83B	CBAL	$J/\psi \rightarrow \eta\pi^+\pi^-$
¹	EDWARDS	83B	CBAL	$J/\psi \rightarrow \eta 2\pi^0$

 $\Gamma(\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{261}/Γ VALUE (units 10^{-4})**6.2 ± 2.2 ± 0.9**

DOCUMENT ID

TECN

COMMENT

BAI	99	BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-----	----	-----	---

 $\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ Γ_{262}/Γ VALUE (units 10^{-3})**5.28 ± 0.06 OUR AVERAGE**

EVTS

DOCUMENT ID

TECN

COMMENT

Error includes scale factor of 1.3. See the ideogram below.

 $5.40 \pm 0.01 \pm 0.11$

638k

ABLIKIM

23BD BES3

 $J/\psi \rightarrow \gamma\eta'$ $5.27 \pm 0.03 \pm 0.05$

36k

ABLIKIM

19T BES

 $J/\psi \rightarrow \gamma\eta'$ $5.43 \pm 0.23 \pm 0.09$

5.0k

¹ ABLIKIM

180 BES3

 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$ $4.77 \pm 0.22 \pm 0.06$ ² ABLIKIM

11 BES3

 $J/\psi \rightarrow \eta'\gamma$ $5.24 \pm 0.12 \pm 0.11$

PEDLAR

09 CLE3

 $J/\psi \rightarrow \eta'\gamma$ 5.55 ± 0.44

35k

ABLIKIM

06E BES2

 $J/\psi \rightarrow \eta'\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $4.50 \pm 0.14 \pm 0.53$

BOLTON

92B MRK3

 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \gamma\gamma$ $4.30 \pm 0.31 \pm 0.71$

BOLTON

92B MRK3

 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \pi^+\pi^-\pi^0$ $4.04 \pm 0.16 \pm 0.85$

622

AUGUSTIN

90 DM2

 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$ $4.39 \pm 0.09 \pm 0.66$

2420

AUGUSTIN

90 DM2

 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$ $4.1 \pm 0.3 \pm 0.6$

BLOOM

83 CBAL

 $e^+e^- \rightarrow 3\gamma + \text{hadrons}$ 2.9 ± 1.1

6

BRANDELIK


79C DASP

 $e^+e^- \rightarrow 3\gamma$ 2.4 ± 0.7

57

BARTEL

76 CNTR

 $e^+e^- \rightarrow 2\gamma\rho$ WEIGHTED AVERAGE
5.28±0.06 (Error scaled by 1.3) $\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}} \text{ (units } 10^{-3})$

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [\mathcal{B}(\eta'(958) \rightarrow \gamma\gamma)] = (1.26 \pm 0.02 \pm 0.05) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [\mathcal{B}(\eta'(958) \rightarrow \gamma\gamma)] \times [\mathcal{B}(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $\mathcal{B}(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $\mathcal{B}(\eta'(958) \rightarrow \gamma\gamma) = (2.307 \pm 0.033) \times 10^{-2}$, $\mathcal{B}(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² ABLIKIM 11 reports $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] / [\mathcal{B}(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [\mathcal{B}(\eta \rightarrow 2\gamma)]$ assuming $\mathcal{B}(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (43.2 \pm 0.7) \times 10^{-2}$, $\mathcal{B}(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$, which we rescale to our best values $\mathcal{B}(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (42.5 \pm 0.5) \times 10^{-2}$, $\mathcal{B}(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\gamma f_0(500) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{255}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10.5 \pm 2.0	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(500) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{256}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5 \pm 5	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(500) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$

Γ_{257}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4 \pm 3	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(980) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{263}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.3 \pm 0.2	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(980) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{264}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.8 \pm 0.3	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma\rho\rho)/\Gamma_{\text{total}}$

Γ_{265}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.5 \pm 0.8 OUR AVERAGE				
4.7 \pm 0.3 \pm 0.9		1 BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
3.75 \pm 1.05 \pm 1.20		2 BURKE 82	MRK2	$J/\psi \rightarrow 4\pi\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.09	90	3 BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$

¹ 4π mass less than 2.0 GeV.

² 4π mass less than 2.0 GeV. We have multiplied $2\rho^0$ measurement by 3 to obtain 2ρ .

³ 4π mass in the range 2.0–25 GeV.


$\Gamma(\gamma\rho\omega)/\Gamma_{\text{total}}$	Γ_{266}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.4 \times 10^{-4}$	90	ABLIKIM	08A BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\gamma\rho\phi)/\Gamma_{\text{total}}$	Γ_{267}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<8.8 \times 10^{-5}$	90	ABLIKIM	08A BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\gamma\omega\omega)/\Gamma_{\text{total}}$	Γ_{268}/Γ			
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.61 ± 0.33 OUR AVERAGE				
6.0 ± 4.8 ± 1.8		ABLIKIM	08A BES2	$J/\psi \rightarrow \gamma\omega\pi^+\pi^-$
1.41 ± 0.2 ± 0.42	120 \pm 17	BISELLO	87 SPEC	$e^+ e^-$, hadrons γ
1.76 ± 0.09 ± 0.45		BALTRUSAIT..85C	MRK3	$e^+ e^- \rightarrow$ hadrons γ

$\Gamma(\gamma\phi\phi)/\Gamma_{\text{total}}$	Γ_{269}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.0 ± 1.2 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.
7.5 ± 0.6 ± 1.2	168	BAI	90B MRK3	$J/\psi \rightarrow \gamma 4K$
3.4 ± 0.8 ± 0.6	33 \pm 7	¹ BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
3.1 ± 0.7 ± 0.4		¹ BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

¹ $\phi\phi$ mass less than 2.9 GeV, η_C excluded.



$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{270}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.6 OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.		
1.66 ± 0.1	1,2 BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
3.8 ± 0.3	3 AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
4.0 ± 0.7	3 EDWARDS	82E CBAL	$J/\psi \rightarrow K^+ K^- \pi^0 \gamma$
4.3 ± 1.7	3,4 SCHARRE	80 MRK2	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.78 ± 0.21 ± 0.33	3,5,6 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
0.83 ± 0.13 ± 0.18	3,7,8 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
0.66 ± 0.17 ± 0.24	3,6,9 BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1.03 ± 0.21 ± 0.26	3,8,10 BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹ Interference with the $J/\psi(1S)$ radiative transition to the broad $K\bar{K}\pi$ pseudoscalar state around 1800 is $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$.

² Interference with $J/\psi \rightarrow \gamma f_1(1420)$ is $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$.

³ Includes unknown branching fraction $\eta(1405) \rightarrow K\bar{K}\pi$.

⁴ Corrected for spin-zero hypothesis for $\eta(1405)$.

⁵ From fit to the $a_0(980)\pi^- \pi^+$ partial wave.

⁶ $a_0(980)\pi$ mode.


⁷ From fit to the $K^*(892)K^- \pi^+$ partial wave.

⁸ $K^* K$ mode.

⁹ From $a_0(980)\pi$ final state.

¹⁰ From $K^*(890)K$ final state.

WEIGHTED AVERAGE
2.8±0.6 (Error scaled by 1.6)



$$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}} \text{ (units } 10^{-3})$$

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0)/\Gamma_{\text{total}}$ Γ_{271}/Γ

<u>VALUE</u> (units 10^{-4})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.78 ± 0.20 OUR AVERAGE	Error includes scale factor of 1.8.		
1.07 ± 0.17 ± 0.11	¹ BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
0.64 ± 0.12 ± 0.07	¹ COFFMAN	90	MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

¹ Includes unknown branching fraction $\eta(1405) \rightarrow \gamma\rho^0$.

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{272}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.0 ± 0.5 OUR AVERAGE				
2.6 ± 0.7 ± 0.4		BAI	99	BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
3.38 ± 0.33 ± 0.64		¹ BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
7.0 ± 0.6 ± 1.1	261	² AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

¹ Via $a_0(980)\pi$.

² Includes unknown branching fraction to $\eta\pi^+\pi^-$.

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$ Γ_{273}/Γ

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 1.3.		
2.1 ± 0.4	BUGG	95	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
1.36 ± 0.38	^{1,2} BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.

² Includes unknown branching fraction to $\rho^0\rho^0$.

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi)/\Gamma_{\text{total}}$ Γ_{274}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<82	95		BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
7.03 ± 0.92 ± 0.91	1.3k	¹ ABLIKIM	18I	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
10.36 ± 1.51 ± 1.54	1.9k	² ABLIKIM	18I	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$

¹ Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

² Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{275}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.63 \times 10^{-6}$	90	ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

 $\Gamma(\gamma\eta(1475) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{276}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.86 \times 10^{-6}$	90	ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$ Γ_{277}/Γ

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.13 ± 0.09	^{1,2} BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.

² Includes unknown branching fraction to $\rho^0\rho^0$.

$\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$				Γ_{278}/Γ
<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.98 \pm 0.08 \pm 0.32$	1045	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

$\Gamma(\gamma\eta(1760) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$				Γ_{279}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<4.80 \times 10^{-6}$	90	ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$

$\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$				Γ_{280}/Γ
<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.14^{+0.50}_{-0.19} \text{ OUR AVERAGE}$				

$2.40 \pm 0.10^{+2.47}_{-0.18}$	1,2	ABLIKIM	16N BES3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$4.4 \pm 0.4 \pm 0.8$	196	2 ABLIKIM	08I BES	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$3.3 \pm 0.8 \pm 0.5$		2 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$2.7 \pm 0.6 \pm 0.6$		2 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$2.4^{+1.5}_{-1.0}$	3,4	BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$ that also finds significant signals for $\eta(2100)$, $0^- +$ phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen $0^- +$ state $X(2500)$ ($M = 2470^{+15+101}_{-19-23}$ MeV, $\Gamma = 230^{+64+56}_{-35-33}$ MeV).

² Includes unknown branching fraction to $\phi\phi$.

³ Estimated by us from various fits.

⁴ Includes unknown branching fraction to $\rho^0\rho^0$.

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$				Γ_{281}/Γ
<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.63 \pm 0.12 \text{ OUR AVERAGE}$	Error includes scale factor of 1.3. See the ideogram below.			

$2.07 \pm 0.16^{+0.02}_{-0.07}$	2.4k	1,2 DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
$1.63 \pm 0.26^{+0.02}_{-0.05}$		3 ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
$1.42 \pm 0.21^{+0.02}_{-0.05}$		4 ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$
$1.33 \pm 0.05 \pm 0.20$		5 AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$
$1.36 \pm 0.09 \pm 0.23$		5 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$
$1.48 \pm 0.25 \pm 0.30$	178	EDWARDS	82B CBAL	$e^+ e^- \rightarrow 2\pi^0\gamma$
2.0 ± 0.7	35	ALEXANDER	78 PLUT	$e^+ e^-$
1.2 ± 0.6	30	6 BRANDELIK	78B DASP	$e^+ e^- \rightarrow \pi^+\pi^-\gamma$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.


² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.744 \pm 0.052 \pm 0.122) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270)) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ Estimated using $B(f_2(1270) \rightarrow \pi\pi) = 0.843 \pm 0.012$. The errors do not contain the uncertainty in the $f_2(1270)$ decay.

⁶ Restated by us to take account of spread of E1, M2, E3 transitions.



$\Gamma(\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$

VALUE (units 10^{-5})

$2.58^{+0.08+0.59}_{-0.09-0.20}$

DOCUMENT ID

TECN

COMMENT

Γ_{282}/Γ

ABLIKIM 18AA BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_1(1285)) / \Gamma_{\text{total}}$

VALUE (units 10^{-3})

0.61 ± 0.08 OUR AVERAGE

DOCUMENT ID

TECN

COMMENT

Γ_{283}/Γ

$0.69 \pm 0.16 \pm 0.20$

¹ BAI 04J BES2 $J/\psi \rightarrow \gamma\gamma\rho^0$

$0.61 \pm 0.04 \pm 0.21$

² BAI 00D BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

$0.45 \pm 0.09 \pm 0.17$

³ BAI 99 BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

$0.625 \pm 0.063 \pm 0.103$

⁴ BOLTON 92 MRK3 $J/\psi \rightarrow \gamma f_1(1285)$

$0.70 \pm 0.08 \pm 0.16$

⁵ BOLTON 92B MRK3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

¹ Assuming $B(f_1(1285) \rightarrow \rho^0\gamma) = 0.055 \pm 0.013$.

² Assuming $\Gamma(f_1(1285) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}} = 0.090 \pm 0.004$.

³ Assuming $\Gamma(f_1(1285) \rightarrow \eta\pi\pi)/\Gamma_{\text{total}} = 0.5 \pm 0.18$.

⁴ Obtained summing the sequential decay channels

$$\begin{aligned} \mathcal{B}(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi\pi\pi\pi) &= (1.44 \pm 0.39 \pm 0.27) \times 10^{-4}; \\ \mathcal{B}(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980)\pi, a_0(980) \rightarrow \eta\pi) &= (3.90 \pm 0.42 \pm 0.87) \times 10^{-4}; \\ \mathcal{B}(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980)\pi, a_0(980) \rightarrow K\bar{K}) &= (0.66 \pm 0.26 \pm 0.29) \times 10^{-4}; \\ \mathcal{B}(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma\rho^0) &= (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}. \end{aligned}$$

⁵ Using $\mathcal{B}(f_1(1285) \rightarrow a_0(980)\pi) = 0.37$, and including unknown branching ratio for $a_0(980) \rightarrow \eta\pi$.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{284}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
38 \pm 10	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{285}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.19 \pm 0.73 \pm 1.34	478	1 DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
1.3 \pm 0.4	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$	

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

Γ_{286}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
1.07 \pm 0.08 \pm 0.36 $-0.07 -0.34$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$

Γ_{287}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
3.5 \pm 1.0	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$

Γ_{288}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.9 \pm 0.3	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$

Γ_{289}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.79 \pm 0.13 OUR AVERAGE			
0.68 \pm 0.04 \pm 0.24	BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
0.76 \pm 0.15 \pm 0.21	1,2 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
0.87 \pm 0.14 \pm 0.11	¹ BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹ Included unknown branching fraction $f_1(1420) \rightarrow K\bar{K}\pi$.

² From fit to the $K^*(892)K$ 1^{++} partial wave.

$\Gamma(\gamma f_0(1500) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{290}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.09 ± 0.24 OUR AVERAGE				
1.21 $\pm 0.29 \pm 0.24$	174	1 DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
1.00 $\pm 0.03 \pm 0.45$		2 ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1.02 $\pm 0.09 \pm 0.45$		2 ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.90 ± 0.17		SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
5.7 ± 0.8	3,4 BUGG		95 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi\pi$.

³ Including unknown branching ratio for $f_0(1500) \rightarrow \pi^+\pi^-\pi^+\pi^-$.

⁴ Assuming that $f_0(1500)$ decays only to two S -wave dipons.

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{291}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.65 \pm 0.26 \pm 0.51$	5.5k	1 ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.1 ± 0.4		SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{292}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.59 \pm 0.16 \pm 0.18$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.7 ± 0.3	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$ Γ_{293}/Γ

<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
18.1 $\pm 1.1 \pm 1.9$	1 ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
12 ± 5	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P -wave.

 $\Gamma(\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{294}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.5 \pm 1.0 \pm 0.7$	BAI	99 BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$	Γ_{295}/Γ
---	-----------------------

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

5.7 ± 0.8 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

$8.0 \pm 0.9 \pm 0.2$	750	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
$3.85 \pm 0.17 \pm 0.73$		³ BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
$3.6 \pm 0.4 \pm 0.4$		³ BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
$5.6 \pm 1.4 \pm 0.9$		³ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$4.5 \pm 0.4 \pm 0.9$		³ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.8 \pm 1.6 \pm 1.4$		³ BALTRUSAIT...87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •


<3.4	90	4	⁴ BRANDELIK	79C DASP	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
<2.3	90	3	ALEXANDER	78 PLUT	$e^+ e^- \rightarrow K^+ K^- \gamma$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = (7.09 \pm 0.46 \pm 0.67) \times 10^{-4}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.888$.

⁴ Assuming isotropic production and decay of the $f'_2(1525)$ and isospin.



$\Gamma(\gamma f_2(1565) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$ Γ_{298}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.32 \pm 0.05^{+0.12}_{-0.02}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta' P$ -wave.

 $\Gamma(\gamma f'_2(1525) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{296}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	--------------------	-------------	----------------

$7.99^{+0.03+0.69}_{-0.04-0.50}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
----------------------------------	---------	-----------	---

 $\Gamma(\gamma f'_2(1525) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{297}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	-------------	--------------------	-------------	----------------

$3.42^{+0.43+1.37}_{-0.51-1.30}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$
----------------------------------	------	----------------------	----------	-------------------------------------

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

 $\Gamma(\gamma f_2(1640) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$ Γ_{299}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	-------------	--------------------	-------------	----------------

$0.28 \pm 0.05 \pm 0.17$	141	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$
--------------------------	-----	---------	---------	---

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{300}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	-------------	--------------------	-------------	----------------

3.8 ± 0.5 OUR AVERAGE

$3.72 \pm 0.30 \pm 0.43$	483	¹ DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
$3.96 \pm 0.06 \pm 1.12$		² ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
$3.99 \pm 0.15 \pm 2.64$		² ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.2	³ SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-----------	------------------------	---------	---

2.5 ± 1.6 ± 0.8	BAI	98H BES	$J/\psi \rightarrow \gamma\pi^0\pi^0$
-----------------	-----	---------	---------------------------------------

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi\pi$.

³ There is a further $(2.4 \pm 0.8) \times 10^{-4}$ scalar contribution at 1765 MeV.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{301}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	-------------	--------------------	-------------	----------------

9.5 + 1.0 - 0.5 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

$8.00^{+0.12+1.24}_{-0.08-0.40}$		¹ ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$11.76 \pm 0.54 \pm 0.94$	1.2k	² DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
$9.62 \pm 0.29^{+3.51}_{-1.86}$		³ BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
$5.0 \pm 0.8^{+1.8}_{-0.4}$		1,4 BAI	96C BES	$J/\psi \rightarrow \gamma K^+K^-$

$9.2 \pm 1.4 \pm 1.4$	¹ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$10.4 \pm 1.2 \pm 1.6$	¹ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$9.6 \pm 1.2 \pm 1.8$	¹ BALTRUSAIT..87	MRK3		$J/\psi \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.3 ± 0.8	⁵ SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$1.6 \pm 0.2 \pm 0.6$	^{1,6} BAI		BES	$J/\psi \rightarrow \gamma K^+ K^-$
< 0.8	90			$J/\psi \rightarrow 4\pi\gamma$
$1.6 \pm 0.4 \pm 0.3$			89B	$J/\psi \rightarrow \gamma\pi^+\pi^-$
3.8 ± 1.6			MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$
			82D	$e^+e^- \rightarrow \eta\eta\gamma$

¹ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied $K^+ K^-$ measurement by 2, and $K_S^0 K_S^0$ by 4 to obtain $K\bar{K}$ result.

² Using CLEO-c data but not authored by the CLEO Collaboration.

³ Includes unknown branching ratio to $K^+ K^-$ or $K_S^0 K_S^0$.

⁴ Assuming $J^P = 2^+$ for $f_0(1710)$.


⁵ There is a further $(6 \pm 2) \times 10^{-4}$ scalar contribution at 1765 MeV.

⁶ Assuming $J^P = 0^+$ for $f_0(1710)$.

⁷ Includes unknown branching fraction to $\rho^0 \rho^0$.

⁸ Includes unknown branching fraction to $\pi^+ \pi^-$.

⁹ Includes unknown branching fraction to $\eta\eta$.



$\Gamma(\gamma f_0(1710) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$				Γ_{302}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.31 \pm 0.06 \pm 0.08$	180	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$				Γ_{303}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.35^{+0.13+1.24}_{-0.11-0.74}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.2 ± 0.4	² SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	--------------------------------	---

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² There is a further $(0.7 \pm 0.1) \times 10^{-4}$ scalar contribution at 1765 MeV.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$				Γ_{304}/Γ
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.5 ± 2.5	¹ SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	--------------------------------	---

¹ There is a further $(2.5 \pm 1.1) \times 10^{-5}$ scalar contribution at 1765 MeV.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma\omega\phi)/\Gamma_{\text{total}}$				Γ_{305}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>

2.5 ± 0.6 OUR AVERAGE

$2.00 \pm 0.08^{+1.38}_{-1.64}$	1.3k	ABLIKIM	13J BES3	$J/\psi \rightarrow \gamma\omega\phi$
---------------------------------	------	---------	----------	---------------------------------------

$2.61 \pm 0.27 \pm 0.65$	95	ABLIKIM	06J BES2	$J/\psi \rightarrow \gamma\omega\phi$
--------------------------	----	---------	----------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1 ± 0.1	¹ SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	--------------------------------	---

¹ There is a further $(2.2 \pm 0.4) \times 10^{-4}$ scalar contribution at 1765 MeV.

$\Gamma(\gamma f_0(1770) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$				Γ_{306}/Γ
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

$1.11 \pm 0.06^{+0.19}_{-0.32}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
---	---------	-----------	---

$\Gamma(\gamma f_0(1770) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$				Γ_{309}/Γ
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.11 $\pm 0.01^{+0.04}_{-0.03}$	¹ ABLIKIM 22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
---------------------------------	--------------------------------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta' P$ -wave.

$\Gamma(\gamma f_2(1810) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$				Γ_{307}/Γ
<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>	
$5.40^{+0.60+3.42}_{-0.67-2.35}$	5.5k	¹ ABLIKIM	13N $J/\psi \rightarrow \gamma\eta\eta$	

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma\eta_1(1855) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$				Γ_{308}/Γ
<u>VALUE</u> (units 10^{-6})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$2.70^{+0.41+0.16}_{-0.35}$	¹ ABLIKIM	22AI BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$	

¹ From a Breit-Wigner fit involving 9 resonances and the resonating exotic $\eta_1(1855) \rightarrow \eta\eta' P$ -wave. For analysis details see ABLIKIM 22AS.

$\Gamma(\gamma f_2(1910) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$				Γ_{310}/Γ
<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.20^{+0.04+0.13}_{-0.13}$	151	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

$\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$				Γ_{311}/Γ
<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$0.7^{+0.1+0.2}_{-0.2}$	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$	

$\Gamma(\gamma f_2(2010) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$				Γ_{312}/Γ
<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.71^{+0.06+0.10}_{-0.06}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$	

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta' P$ -wave.

$\Gamma(\gamma f_0(2020) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$				Γ_{313}/Γ
<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
42 \pm 10	SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$		

$\Gamma(\gamma f_0(2020) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$				Γ_{314}/Γ
<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
55 \pm 25	SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$		

$\Gamma(\gamma f_0(2020) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$				Γ_{315}/Γ
<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10 \pm 10	SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$		

$\Gamma(\gamma f_0(2020) \rightarrow \gamma\eta'\eta')/\Gamma_{\text{total}}$	Γ_{316}/Γ		
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.63 \pm 0.06^{+0.31}_{-0.46}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta'\eta'$, and ($\eta' X$), with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

$\Gamma(\gamma f_0(2020) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$	Γ_{317}/Γ		
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$2.28 \pm 0.12^{+0.29}_{-0.20}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P -wave.

$\Gamma(\gamma f_4(2050))/\Gamma_{\text{total}}$	Γ_{318}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.7 \pm 0.5 \pm 0.5$	¹ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$

¹ Assuming branching fraction $f_4(2050) \rightarrow \pi\pi/\text{total} = 0.167$.

$\Gamma(\gamma f_4(2050) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$	Γ_{319}/Γ		
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.06 \pm 0.01^{+0.03}_{-0.01}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P -wave.

$\Gamma(\gamma f_0(2100) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$	Γ_{320}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.13^{+0.09+0.64}_{-0.10-0.28}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ± 1.5	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	----	------	---

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_0(2100) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$	Γ_{322}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$6.24 \pm 0.48 \pm 0.87$	744	¹ DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.8	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	----	------	---

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2100) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{321}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
32 \pm 20	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 $\Gamma(\gamma f_0(2200))/\Gamma_{\text{total}}$ Γ_{323}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
1.5	¹ AUGUSTIN 88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
1 Includes unknown branching fraction to $K_S^0 K_S^0$.			

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{326}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
5 \pm 2	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{324}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.86 \pm 0.49 \pm 1.20	490	¹ DOBBS 15		$J/\psi \rightarrow \gamma K\bar{K}$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.5 \pm 0.5	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$	

¹ Using CLEO-c data but not authored by the CLEO Collaboration. $\Gamma(\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{325}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.72 \pm 0.08 \pm 0.17	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{327}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.7 \pm 0.4	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 $\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$ Γ_{328}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
>300			¹ BAI 96B	BES	$e^+ e^- \rightarrow \gamma\bar{p}p, K\bar{K}$
>250	99.9		² HASAN 96	SPEC	$\bar{p}p \rightarrow \pi^+\pi^-$
< 2.3	95		³ AUGUSTIN 88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$
< 1.6	95		³ AUGUSTIN 88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$12.4^{+6.4}_{-5.2} \pm 2.8$		23	³ BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$8.4^{+3.4}_{-2.8} \pm 1.6$		93	³ BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

¹ Using BARNES 93.² Using BAI 96B.³ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{329}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 3.9	90	1,2 DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

14 ± 8 ± 4	BAI	98H BES	$J/\psi \rightarrow \gamma\pi^0\pi^0$
8.4 ± 2.6 ± 3.0	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+\pi^-$ and $\pi^0\pi^0$ are $2.6/5.2 \times 10^{-5}$ and $1.3/1.9 \times 10^{-5}$, respectively.

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{330}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 4.1	90	1,2 DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.6	³ DEL-AMO-SA..100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+K^-$
< 2.9	³ DEL-AMO-SA..100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
6.6 ± 2.9 ± 2.4	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+K^-$
10.8 ± 4.0 ± 3.2	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for K^+K^- and $K_S^0K_S^0$ are $1.7/3.1 \times 10^{-5}$ and $1.2/2.0 \times 10^{-5}$, respectively.

³ For spin 2 and helicity 0; other combinations lead to more stringent upper limits.

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{331}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.5±0.6±0.5	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{332}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.95±0.21^{+0.66}_{-0.72}	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.1	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-----------	-----------	---------	---

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{333}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4 ± 2	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------	-----------	---------	---

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{334}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5 ± 0.4	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-----------	-----------	---------	---

$\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$	Γ_{335}/Γ			
<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$6.09 \pm 0.64^{+4.00}_{-1.68}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$	

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta' \eta'$, and ($\eta' X$), with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

$\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$	Γ_{336}/Γ			
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$0.10 \pm 0.02^{+0.01}_{-0.02}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$	

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta' P$ -wave.

$\Gamma(\gamma f_0(2470) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$	Γ_{340}/Γ			
<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$8.18 \pm 1.77^{+3.73}_{-2.23}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$	

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta' \eta'$, and ($\eta' X$), with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

$\Gamma(\gamma f_2(2340) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$	Γ_{337}/Γ				
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$5.60^{+0.62+2.37}_{-0.65-2.07}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$	

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_2(2340) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$	Γ_{339}/Γ			
<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$8.67 \pm 0.70^{+0.61}_{-1.67}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$	

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta' \eta'$, and ($\eta' X$), with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

$\Gamma(\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$	Γ_{338}/Γ			
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$5.54^{+0.34+3.82}_{-0.40-1.49}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$	

$\Gamma(\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{341}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.7 $^{+0.6}_{-0.8}$ OUR AVERAGE				Error includes scale factor of 1.6.
$3.93 \pm 0.38^{+0.31}_{-0.84}$		¹ ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
$2.2 \pm 0.4 \pm 0.4$	264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2.87 \pm 0.09^{+0.49}_{-0.52}$	4265	² ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

- ¹ From a fit of the measured $\pi^+\pi^-\eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold with a Flatte formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the $X(1835)$; fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.
- ² From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two states $\gamma X(2120)$ and $\gamma X(2370)$, for $M(\pi^+\pi^-\eta') < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$.

 $\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{342}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.77 $^{+0.15}_{-0.09}$ OUR AVERAGE				
$0.90^{+0.04+0.27}_{-0.11-0.55}$		¹ ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$
$1.14^{+0.43+0.42}_{-0.30-0.26}$	231	² ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p\bar{p}$
$0.70 \pm 0.04^{+0.19}_{-0.08}$		BAI	03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$

- ¹ From the fit including final state interaction effects in isospin 0 *S*-wave according to SIBIRTSEV 05A.
- ² From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.

 $\Gamma(\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta)/\Gamma_{\text{total}}$ Γ_{343}/Γ

<u>VALUE</u> (units 10^{-5})		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.31 $^{+0.33+1.96}_{-0.30-1.29}$		ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$

 $\Gamma(\gamma X(1835) \rightarrow \gamma\gamma\phi(1020))/\Gamma_{\text{total}}$ Γ_{344}/Γ

<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.77 \pm 0.35 \pm 0.25$	305	¹ ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
$8.09 \pm 1.99 \pm 1.36$	1.3k	² ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$

- ¹ Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.
- ² Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

$\Gamma(\gamma X(1835) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{345}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.56 \times 10^{-6}$	90	ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$

 $\Gamma(\gamma X(1835) \rightarrow \gamma 3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{346}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.44 \pm 0.36^{+0.60}_{-0.74}$	0.6k	ABLIKIM	13U	BES3 $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$

 $\Gamma(\gamma X(2370) \rightarrow \gamma K^+ K^- \eta')/\Gamma_{\text{total}}$ Γ_{347}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.79 \pm 0.23 \pm 0.65$	ABLIKIM	20Q	BES3 $J/\psi \rightarrow \gamma K^+ K^- \eta'$

 $\Gamma(\gamma X(2370) \rightarrow \gamma K_S^0 K_S^0 \eta')/\Gamma_{\text{total}}$ Γ_{348}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.18 \pm 0.32 \pm 0.39$	ABLIKIM	20Q	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$

 $\Gamma(\gamma X(2370) \rightarrow \gamma \eta \eta \eta')/\Gamma_{\text{total}}$ Γ_{349}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<9.2 \times 10^{-6}$	90	ABLIKIM	21C	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta \eta'$

 $\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{350}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.38 \pm 0.07 \pm 0.07$	49	EATON	84	MRK2 $e^+ e^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.11	90	PERUZZI	78	MRK1 $e^+ e^-$
---------	----	---------	----	----------------

 $\Gamma(\gamma p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{351}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<0.79 \times 10^{-3}$	90	EATON	84	MRK2 $e^+ e^-$

 $\Gamma(\gamma A\bar{A})/\Gamma_{\text{total}}$ Γ_{352}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<0.13 \times 10^{-3}$	90	HENRARD	87	DM2 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.16 \times 10^{-3}$	90	BAI	98G	BES $e^+ e^-$
------------------------	----	-----	-----	---------------

 $\Gamma(\gamma A^0 \rightarrow \gamma \text{invisible})/\Gamma_{\text{total}}$ Γ_{353}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.7 \times 10^{-6}$	90	88M	¹ ABLIKIM	20K	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<6.3 \times 10^{-6}$	90	3.7M	² INSLER	10	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
-----------------------	----	------	---------------------	----	--

¹ For a narrow state, A^0 , with mass $m_{A^0} < 1.2$ GeV. The limit varies with m_{A^0} , reaching its largest value of 1.7×10^{-6} at 1.2 GeV and being 7.0×10^{-7} for $m_{A^0} = 0$.

² The limit varies with mass m_{A^0} of a narrow state A^0 and is 4.3×10^{-6} for $m_{A^0} = 0$, reaches its largest value of 6.3×10^{-6} at $m_{A^0} = 500$ MeV, and is 3.6×10^{-6} at $m_{A^0} = 960$ MeV.

$\Gamma(\gamma A^0 \rightarrow \gamma\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{354}/Γ
(narrow state A^0 with 0.2 GeV $< m_{A^0} < 3$ GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.8 \times 10^{-7}$	90	¹ ABLIKIM	22H	BES3 $J/\psi \rightarrow \gamma\mu^+\mu^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<0.5 \times 10^{-5}$	90	² ABLIKIM	16E	BES3 $J/\psi \rightarrow \gamma\mu^+\mu^-$
$<2.1 \times 10^{-5}$	90	³ ABLIKIM	12	BES3 $J/\psi \rightarrow \gamma\mu^+\mu^-$

¹ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3.0 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(1.2\text{--}778.0) \times 10^{-9}$.

² For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(2.8\text{--}495.3) \times 10^{-8}$.

³ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.21–3.00 GeV. The measured 90% CL limit as a function of m_{A^0} ranges from 4×10^{-7} to 2.1×10^{-5} .

DALITZ DECAYS

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{355}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.56 \pm 1.32 \pm 0.50$	39	ABLIKIM	14I	BES3 $J/\psi \rightarrow \pi^0 e^+ e^-$

 $\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_{356}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.42 \pm 0.04 \pm 0.07$	2.47k	^{1,2} ABLIKIM	19A	BES3 $J/\psi \rightarrow \eta e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

$1.16 \pm 0.07 \pm 0.06$	320	¹ ABLIKIM	14I	BES3 $J/\psi \rightarrow \eta e^+ e^-$
--------------------------	-----	----------------------	-----	--

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decays.

² Approximation of the transition form factor squared as an incoherent sum of the ρ -meson and one-pole non-resonant amplitudes gives the pole mass $m(\Lambda) = 2.56 \pm 0.04 \pm 0.03$ GeV. Supersedes ABLIKIM 14I.

 $\Gamma(\eta'(958) e^+ e^-)/\Gamma_{\text{total}}$ Γ_{357}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.59 \pm 0.07 \pm 0.17$	8.9k	¹ ABLIKIM	19H	BES3 $J/\psi \rightarrow \eta'(958) e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

$5.81 \pm 0.16 \pm 0.31$	1.4k	^{1,2} ABLIKIM	14I	BES3 $J/\psi \rightarrow \eta'(958) e^+ e^-$
--------------------------	------	------------------------	-----	--

¹ Using both $\eta' \rightarrow \gamma\pi^+\pi^-$ and $\eta' \rightarrow \pi^+\pi^-\eta$ decays.

² Superseded by ABLIKIM 19H.

 $\Gamma(X(1835) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}$ Γ_{358}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.58 \pm 0.19 \pm 0.16$	1364	¹ ABLIKIM	22B	BES3 $J/\psi \rightarrow \pi^+ \pi^- \eta' e^+ e^-$

¹ Assuming constructive interference. Destructive interference gives a value of $(4.43 \pm 0.23 \pm 0.19) \times 10^{-6}$ for this branching fraction.

 $\Gamma(X(2120) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}$ Γ_{359}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.82 \pm 0.12 \pm 0.06$	310	ABLIKIM	22B	BES3 $J/\psi \rightarrow \pi^+ \pi^- \eta' e^+ e^-$

$\Gamma(X(2370)e^+e^-)$	$X \rightarrow \pi^+\pi^-\eta')$	Γ_{total}	Γ_{360}/Γ	
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.08 \pm 0.14 \pm 0.10$	397	ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'e^+e^-$

$\Gamma(\eta U \rightarrow \eta e^+e^-)$	Γ_{total}	Γ_{361}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.11 \times 10^{-7}$	90	¹ ABLIKIM	19A BES3	$J/\psi \rightarrow \eta e^+e^-$

¹ For a dark photon U with mass between 10 and 2400 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.9×10^{-8} to 91.1×10^{-8} .

$\Gamma(\eta'(958)U \rightarrow \eta'(958)e^+e^-)$	Γ_{total}	Γ_{362}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-7}$	90	¹ ABLIKIM	19H BES3	$J/\psi \rightarrow \eta'(958)e^+e^-$

¹ For a dark photon U with mass between 100 and 2100 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.8×10^{-8} to 2.0×10^{-7} . The corresponding limits on the branching fraction $J/\psi \rightarrow \eta' U$ range from 5.7×10^{-8} to 7.4×10^{-7} .

$\Gamma(\phi e^+e^-)$	Γ_{total}	Γ_{363}/Γ		
VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	¹ ABLIKIM	19AB BES3	$J/\psi \rightarrow \phi e^+e^-$

¹ Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ and $B(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) = (34.49 \pm 0.30)\%$.

WEAK DECAYS

SetUpHeader10

$\Gamma(D^-e^+\nu_e + \text{c.c.})$	Γ_{total}	Γ_{364}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.1 \times 10^{-8}$	90	ABLIKIM	21Q BES3	$e^+e^- \rightarrow J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.2 \times 10^{-5}$	90	ABLIKIM	06M BES2	$e^+e^- \rightarrow J/\psi$
-----------------------	----	---------	----------	-----------------------------

$\Gamma(\bar{D}^0e^+e^- + \text{c.c.})$	Γ_{total}	Γ_{365}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.5 \times 10^{-8}$	90	¹ ABLIKIM	17AF BES3	$e^+e^- \rightarrow J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.1 \times 10^{-5}$	90	ABLIKIM	06M BES2	$e^+e^- \rightarrow J/\psi$
-----------------------	----	---------	----------	-----------------------------

¹ Using D^0 decays to $K^-\pi^+$, $K^-\pi^+\pi^0$, and $K^-\pi^+\pi^+\pi^-$.

$\Gamma(D_s^-e^+\nu_e + \text{c.c.})$	Γ_{total}	Γ_{366}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-6}$	90	ABLIKIM	14R BES3	$e^+e^- \rightarrow J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.6 \times 10^{-5}$	90	¹ ABLIKIM	06M BES2	$e^+e^- \rightarrow J/\psi$
-----------------------	----	----------------------	----------	-----------------------------

¹ Using $B(D_s^- \rightarrow \phi\pi^-) = 4.4 \pm 0.5\%$.

$\Gamma(D_s^{*-} e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.8 \times 10^{-6}$	90

 $\Gamma(D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$< 7.5 \times 10^{-5}$	90

 $\Gamma(\bar{D}^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.7 \times 10^{-4}$	90

 $\Gamma(\bar{D}^0 \bar{K}^{*0} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$< 2.5 \times 10^{-6}$	90

 $\Gamma(D_s^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.3 \times 10^{-4}$	90

 $\Gamma(D_s^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.3 \times 10^{-5}$	90

 CHARGE CONJUGATION (C), PARITY (P),
 LEPTON FAMILY NUMBER (LF) VIOLATING MODES

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE	CL%
$< 2.7 \times 10^{-7}$	90
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$< 0.5 \times 10^{-5}$	90
$< 1.6 \times 10^{-4}$	90
$< 2.2 \times 10^{-5}$	90
$< 50 \times 10^{-5}$	90

¹ WICHT 08 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [\mathcal{B}(B^+ \rightarrow J/\psi(1S) K^+)] < 0.16 \times 10^{-6}$ which we divide by our best value $\mathcal{B}(B^+ \rightarrow J/\psi(1S) K^+) = 1.020 \times 10^{-3}$.

 $\Gamma(\gamma\phi)/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.4 \times 10^{-6}$	90

 $\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.6 \times 10^{-7}$	90
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$< 1.1 \times 10^{-6}$	90

 Γ_{367}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	14R	$e^+ e^- \rightarrow J/\psi$

 Γ_{368}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	08J	$e^+ e^- \rightarrow J/\psi$

 Γ_{369}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	08J	$e^+ e^- \rightarrow J/\psi$

 Γ_{370}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	14K	$e^+ e^- \rightarrow J/\psi$

 Γ_{371}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	08J	$e^+ e^- \rightarrow J/\psi$

 Γ_{372}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	14K	$e^+ e^- \rightarrow J/\psi$

 Γ_{373}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	14Q	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •		
ADAMS	08	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
¹ WICHT	08	$B^\pm \rightarrow K^\pm \gamma\gamma$
ABLIKIM	07J	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
BARTEL	77	$e^+ e^-$

DOCUMENT ID	TECN	COMMENT
ABLIKIM	14Q	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

 Γ_{374}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	13L	$e^+ e^- \rightarrow J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •		
BAI	03D	$e^+ e^- \rightarrow J/\psi$

 Γ_{375}/Γ

$\Gamma(e^\pm\tau^\mp)/\Gamma_{\text{total}}$		Γ_{376}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.5 \times 10^{-8}$	90	ABLIKIM	21M	BES3 $e^+e^- \rightarrow J/\psi$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<8.3 \times 10^{-6}$	90	¹ ABLIKIM	04	BES $e^+e^- \rightarrow J/\psi$

¹ Superseded by ABLIKIM 21M.

$\Gamma(\mu^\pm\tau^\mp)/\Gamma_{\text{total}}$		Γ_{377}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-6}$	90	ABLIKIM	04	BES $e^+e^- \rightarrow J/\psi$

$\Gamma(\Lambda_c^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$		Γ_{378}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.9 \times 10^{-8}$	90	ABLIKIM	19AF	BES3 $e^+e^- \rightarrow J/\psi \rightarrow pK^-\pi^+e^- (+ \text{c.c.})$

OTHER DECAYS

$\Gamma(\text{invisible})/\Gamma(e^+e^-)$		Γ_{379}/Γ_5		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.6 \times 10^{-2}$	90	LEES	13I	BABR $B \rightarrow K^{(*)}J/\psi$

$\Gamma(\text{invisible})/\Gamma(\mu^+\mu^-)$		Γ_{379}/Γ_7		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-2}$	90	ABLIKIM	08G	BES2 $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$J/\psi(1S)$ REFERENCES

ABLIKIM	23BD	PR D108	092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BU	PR D108	112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23S	PR D107	072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	23	JHEP	2306 196	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
GONG	23	PR D107	072008	G. Gong <i>et al.</i>	(BELLE Collab.)
LEES	23	PR D107	072001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LIAO	23	PR D107	112007	L. Liao <i>et al.</i>	
ZHU	23	PR D107	012006	W. Zhu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	22AI	PRL	129 192002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D106	072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AS	PR D106	072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D107	079901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AY	PR D106	112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22B	PRL	129 022002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22C	PR D105	072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22H	PR D105	012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	22	EPJ C82	938	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
ABLIKIM	21AM	PR D104	092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AT	JHEP	2111 226	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21C	PR D103	012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21M	PR D103	112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21Q	JHEP	2106 157	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	21	PR D103	092001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	21B	PR D104	112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	21C	PR D104	112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SARANTSEV	21	PL	B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
ABLIKIM	20	PR D101	012004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20K	PR D101	112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20Q	EPJ C80	746	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	20	JHEP	2007 112	V.V. Anashin <i>et al.</i>	(KEDR Collab.)

ABLIKIM	19A	PR D99 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D104 099901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AB	PR D99 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AC	PR D99 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AF	PR D99 072006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AN	PR D99 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19H	PR D99 012013	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19N	PR D99 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19Q	PL B791 375	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19T	PRL 122 142002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18AB	PR D98 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	18A	JHEP 1805 119	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	18E	PR D98 112015	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AH	PR D96 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17D	PR D95 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	16E	PR D93 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16K	PR D93 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16N	PR D93 112011	M. Ablikim	(BESIII Collab.)
ABLIKIM	16P	PR D94 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16Q	PL B761 98	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15K	PR D91 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	14I	PR D89 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14K	PR D89 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14N	PR D90 052009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14Q	PR D90 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14R	PR D90 112014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
AULCHENKO	14	PL B731 227	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13L	PR D87 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13P	PR D87 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12C	PR D86 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ABLIKIM	12P	CP C36 1031	M. Ablikim <i>et al.</i>	(BES II Collab.)
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)
ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DEL-AMO-SA...	10O	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
INSLER	10	PR D81 091101	J. Insler <i>et al.</i>	(CLEO Collab.)
ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES II Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSON	08	PR D78 032012	D. Besson <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Fermilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	06A	PR D73 051103	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer	
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)

BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
SETH	04	PR D69 097503	K.K. Seth	
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98G	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 and E706 Collab.)
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
HSUEH	92	PR D45 2181	S. Hsueh, S. Palestini	(FNAL, TORI)
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
BISELLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LAPO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LAPO, CLER, FRAS+)
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
HENRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LAPO+)
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LAPO, PADO)
BALTRUSAIT...	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
BALTRUSAIT...	85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
BALTRUSAIT...	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
		Translated from YAF 41 733.		
EATON	84	PR D29 804	M.W. Eaton <i>et al.</i>	(LBL, SLAC)
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
EDWARDS	83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)
EDWARDS	82B	PR D25 3065	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
Also		ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)

LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
BESCH	81	ZPHY C8 1	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)
PARTRIDGE	80	PRL 44 712	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34 1471.		
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
ALEXANDER	78	PL 72B 493	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)
BESCH	78	PL 78B 347	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)
BRANDELIK	78B	PL 74B 292	R. Brandelik <i>et al.</i>	(DASP Collab.)
PERUZZI	78	PR D17 2901	I. Peruzzi <i>et al.</i>	(SLAC, LBL)
BARTEL	77	PL 66B 489	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BURMESTER	77D	PL 72B 135	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
VANNUCCI	77	PR D15 1814	F. Vannucci <i>et al.</i>	(SLAC, LBL)
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BRAUNSCH...	76	PL 63B 487	W. Braunschweig <i>et al.</i>	(DASP Collab.)
JEAN-MARIE	76	PRL 36 291	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG
BALDINI...	75	PL 58B 471	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)
BOYARSKI	75	PRL 34 1357	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC
DASP	75	PL 56B 491	W. Braunschweig <i>et al.</i>	(DASP Collab.)
ESPOSITO	75B	LNC 14 73	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
FORD	75	PRL 34 604	R.L. Ford <i>et al.</i>	(SLAC, PENN)
