

$T_{c\bar{c}}(4020)$

$I^G(J^{PC}) = 1^+(?^-)$

was $X(4020)$

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Charged $T_{c\bar{c}}(4020)$ seen by ABLIKIM 13X from $e^+e^- \rightarrow \pi^+\pi^- h_c(1P)$ at c.m. energy from 3.90 to 4.42 GeV as a peak in the invariant mass distribution of the $\pi^\pm h_c(1P)$ system, and by ABLIKIM 14B from $e^+e^- \rightarrow (D^*\bar{D}^*)^\pm \pi^\mp$ events in $(D^*\bar{D}^*)^\pm$ mass. A neutral $T_{c\bar{c}}(4020)$ seen by ABLIKIM 14P at three c.m. energies in the same range in $e^+e^- \rightarrow \pi^0\pi^0 h_c(1P)$ as a peak in the larger of the two masses recoiling against a π^0 . ABLIKIM 15AA observes a 5.9σ signal in $(D^*\bar{D}^*)^0$ in $e^+e^- \rightarrow (D^*\bar{D}^*)^0 \pi^0$ events using collisions at two c.m. energies. Production rates and mass values support grouping neutral and charged $T_{c\bar{c}}(4020)$ together as manifestations of a single $I = 1$ particle.

$T_{c\bar{c}}(4020)$ MASS

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | CHG | COMMENT |
|--|------|------------------------|-----------|-----|---|
| 4024.1±1.9 OUR AVERAGE | | | | | |
| 4025.5 ^{+2.0} _{-4.7} ± 3.1 | 116 | ¹ ABLIKIM | 15AA BES3 | 0 | $e^+e^- \rightarrow (D^*\bar{D}^*)^0 \pi^0$ |
| 4026.3 ± 2.6 ± 3.7 | 401 | ¹ ABLIKIM | 14B BES3 | ± | $e^+e^- \rightarrow (D^*\bar{D}^*)^\pm \pi^\mp$ |
| 4023.9 ± 2.2 ± 3.8 | 61 | ^{1,2} ABLIKIM | 14P BES3 | 0 | $e^+e^- \rightarrow \pi^0\pi^0 h_c$ |
| 4022.9 ± 0.8 ± 2.7 | 253 | ¹ ABLIKIM | 13X BES3 | ± | $e^+e^- \rightarrow \pi^+\pi^- h_c$ |


¹ Neglecting interference between the $T_{c\bar{c}}(4020)$ and non-resonant continuum.

² Assuming $J^P = 1^+$ and width of 7.9 ± 2.6 MeV.

$T_{c\bar{c}}(4020)$ WIDTH

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | CHG | COMMENT |
|---|------|----------------------|-----------|-----|---|
| 13 ±5 OUR AVERAGE | | | | | |
| Error includes scale factor of 1.7. See the ideogram below. | | | | | |
| 23.0 ± 6.0 ± 1.0 | 116 | ¹ ABLIKIM | 15AA BES3 | 0 | $e^+e^- \rightarrow (D^*\bar{D}^*)^0 \pi^0$ |
| 24.8 ± 5.6 ± 7.7 | 401 | ¹ ABLIKIM | 14B BES3 | ± | $e^+e^- \rightarrow (D^*\bar{D}^*)^\pm \pi^\mp$ |
| 7.9 ± 2.7 ± 2.6 | 253 | ¹ ABLIKIM | 13X BES3 | ± | $e^+e^- \rightarrow \pi^+\pi^- h_c$ |

¹ Neglecting interference between the $T_{c\bar{c}}(4020)$ and non-resonant continuum.



$T_{c\bar{c}}(4020)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|------------------------------------|--------------------------------|
| $\Gamma_1 h_c(1P)\pi$ | seen |
| $\Gamma_2 D^*\bar{D}^*$ | seen |
| $\Gamma_3 D\bar{D}^* + c.c.$ | not seen |
| $\Gamma_4 \eta_c\pi^+\pi^-$ | not seen |
| $\Gamma_5 \eta_c(1S)\rho(770)^\pm$ | |
| $\Gamma_6 J/\psi(1S)\pi^\pm$ | not seen |

$T_{c\bar{c}}(4020)$ BRANCHING RATIOS

| $\Gamma(h_c(1P)\pi)/\Gamma_{\text{total}}$ | Γ_1/Γ |
|--|--|
| VALUE | EVTS DOCUMENT ID TECN CHG COMMENT |
| seen | 61 ABLIKIM 14P BES3 0 $e^+e^- \rightarrow \pi^0\pi^0 h_c$ |
| seen | 253 ABLIKIM 13X BES3 ± $e^+e^- \rightarrow \pi^+\pi^- h_c$ |

| $\Gamma(D^*\bar{D}^*)/\Gamma_{\text{total}}$ | Γ_2/Γ |
|--|--|
| VALUE | EVTS DOCUMENT ID TECN CHG COMMENT |
| seen | 116 ¹ ABLIKIM 15AA BES3 0 $e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$ |
| seen | 401 ¹ ABLIKIM 14B BES3 ± $e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$ |

¹ Neglecting interference between the $T_{c\bar{c}}(4020)$ and non-resonant continuum.

| $\Gamma(D\bar{D}^* + \text{c.c.})/\Gamma_{\text{total}}$ | Γ_3/Γ |
|--|--|
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u> |
| not seen | ABLIKIM 15AC BES3 ± $e^+ e^- \rightarrow \pi^\pm (D\bar{D}^*)^\mp$ |
| $\Gamma(\eta_c \pi^+ \pi^-)/\Gamma_{\text{total}}$ | Γ_4/Γ |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |
| not seen | ¹ VINOKUROVA 15 BELL $B^+ \rightarrow K^+ \eta_c \pi^+ \pi^-$ |
| ¹ VINOKUROVA 15 reports $B(B^+ \rightarrow K^+ T_{c\bar{c}}(4020)^0) \times B(T_{c\bar{c}} \rightarrow \eta_c \pi^+ \pi^-) < 1.6 \times 10^{-5}$ at 90% CL. | |
| $\Gamma(\eta_c(1S)\rho(770)^\pm)/\Gamma(h_c(1P)\pi)$ | Γ_5/Γ_1 |
| <u>VALUE</u> <u>CL%</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |
| <1.2 90 | ¹ ABLIKIM 19BC BES3 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \eta_c(1S)$ |
| ¹ Using $e^+ e^- \rightarrow \pi^\mp (Z_c(4020)^\pm \rightarrow h_c(1P)\pi^\pm)$ cross section at 4.23, 4.26 and 4.36 GeV from ABLIKIM 13X. | |
| $\Gamma(J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$ | Γ_6/Γ |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |
| not seen | ¹ ABLIKIM 17J BES3 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ |
| ¹ From Partial Wave Analysis assuming $J^P = 1^+$. | |

$T_{c\bar{c}}(4020)$ REFERENCES

| | | | |
|------------|--|--|------------------------------------|
| ABLIKIM | 19BC PR D100 111102 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17J PRL 119 072001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 15AA PRL 115 182002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 15AC PR D92 092006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| VINOKUROVA | ¹ 15 JHEP 1506 132 Also JHEP 1702 088 (errat.) | A. Vinokurova <i>et al.</i> A. Vinokurava <i>et al.</i> | (BELLE Collab.) (BELLE Collab.) |
| ABLIKIM | 14B PRL 112 132001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 14P PRL 113 212002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13X PRL 111 242001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |