

**$T_{c\bar{c}s1}(4220)^+$** 

$$I(J^P) = \frac{1}{2}(1^+)$$

OMMITTED FROM SUMMARY TABLE  
was  $Z_{cs}(4220)^+$

Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on "Heavy Non- $q\bar{q}$  Mesons."

Seen by AAIJ 21E in  $B^+ \rightarrow T_{c\bar{c}s1}(4220)^+ \phi$  with  $T_{c\bar{c}s1}(4220)^+ \rightarrow J/\psi K^+$  using an amplitude analysis of  $B^+ \rightarrow J/\psi \phi K^+$  with a significance (accounting for systematic uncertainties) of  $5.9\sigma$ . The  $J^P = 1^+$  assignment is favored over  $1^-$  with a significance of  $2\sigma$  and other assignments are disfavored by  $4.9\sigma$ .

 **$T_{c\bar{c}s1}(4220)^+$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>4216 \pm 24^{+43}_{-30}</math></b>	24k	<sup>1</sup> AAIJ	21E LHCb	$B^+ \rightarrow J/\psi \phi K^+$

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.9\sigma$ .

 **$T_{c\bar{c}s1}(4220)^+$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>233 \pm 52^{+97}_{-73}</math></b>	24k	<sup>1</sup> AAIJ	21E LHCb	$B^+ \rightarrow J/\psi \phi K^+$

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.9\sigma$ .

 **$T_{c\bar{c}s1}(4220)^+$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad J/\psi K^+$	seen

$\Gamma(J/\psi K^+)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
<b>seen</b>	<b><math>24k \quad 1 \quad AAIJ \quad 21E \quad LHCb \quad B^+ \rightarrow J/\psi \phi K^+</math></b>

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.9\sigma$ .

 **$T_{c\bar{c}s1}(4220)^+$  REFERENCES**

AAIJ

21E PRL 127 082001

R. Aaij *et al.*

(LHCb Collab.) JP