

**$\chi_{b2}(2P)$** 
 $I^G(J^{PC}) = 0^+(2^{++})$   
 $J$  needs confirmation.

Observed in radiative decay of the  $\Upsilon(3S)$ , therefore  $C = +$ . Branching ratio requires E1 transition, M1 is strongly disfavored, therefore  $P = +$ .

 **$\chi_{b2}(2P)$  MASS**VALUE (MeV)DOCUMENT ID

**10268.65±0.22±0.50 OUR EVALUATION** From  $\gamma$  energy below, using  $\Upsilon(3S)$  mass =  $10355.2 \pm 0.5$  MeV

 **$m_{\chi_{b2}(2P)} - m_{\chi_{b1}(2P)}$** VALUE (MeV)DOCUMENT IDTECNCOMMENT**13.10±0.24 OUR AVERAGE**

12.3 $\pm 2.6 \pm 0.6$	<sup>1</sup> AAIJ	14BG LHCb	$p p \rightarrow \gamma \mu^+ \mu^- X$
13.04 $\pm 0.26$	LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
13.5 $\pm 0.4 \pm 0.5$	<sup>2</sup> HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X, \ell^+ \ell^- \gamma \gamma$

<sup>1</sup> From the  $\chi_{bj}(2P) \rightarrow \Upsilon(1S)\gamma$  transition.

<sup>2</sup> From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

 **$\gamma$  ENERGY IN  $\Upsilon(3S)$  DECAY**VALUE (MeV)EVTSDOCUMENT IDTECNCOMMENT**86.19±0.22 OUR EVALUATION**

Treating systematic errors as correlated

**86.40±0.18 OUR AVERAGE**

86.04 $\pm 0.06 \pm 0.27$	ARTUSO	05	CLEO	$\Upsilon(3S) \rightarrow \gamma X$
86 $\pm 1$	101	CRAWFORD	92B CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
86.7 $\pm 0.4$	10319	<sup>3</sup> HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X$
86.9 $\pm 0.4$	157	<sup>4</sup> HEINTZ	92 CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
86.4 $\pm 0.1 \pm 0.4$	30741	MORRISON	91 CLE2	$e^+ e^- \rightarrow \gamma X$

<sup>3</sup> A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

<sup>4</sup> A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

 **$\chi_{b2}(2P)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1 \omega \Upsilon(1S)$	$(1.10^{+0.34}_{-0.30}) \%$	
$\Gamma_2 \gamma \Upsilon(2S)$	$(8.9 \pm 1.2) \%$	
$\Gamma_3 \gamma \Upsilon(1S)$	$(6.6 \pm 0.8) \%$	
$\Gamma_4 \pi \pi \chi_{b2}(1P)$	$(5.1 \pm 0.9) \times 10^{-3}$	
$\Gamma_5 D^0 X$	$< 2.4 \%$	90%
$\Gamma_6 \pi^+ \pi^- K^+ K^- \pi^0$	$< 1.1 \times 10^{-4}$	90%

$\Gamma_7$	$2\pi^+ \pi^- K^- K_S^0$	< 9	$\times 10^{-5}$	90%
$\Gamma_8$	$2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 7	$\times 10^{-4}$	90%
$\Gamma_9$	$2\pi^+ 2\pi^- 2\pi^0$	(3.9 $\pm$ 1.6 )	$\times 10^{-4}$	
$\Gamma_{10}$	$2\pi^+ 2\pi^- K^+ K^-$	(9 $\pm$ 4 )	$\times 10^{-5}$	
$\Gamma_{11}$	$2\pi^+ 2\pi^- K^+ K^- \pi^0$	(2.4 $\pm$ 1.1 )	$\times 10^{-4}$	
$\Gamma_{12}$	$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	(4.7 $\pm$ 2.3 )	$\times 10^{-4}$	
$\Gamma_{13}$	$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 4	$\times 10^{-4}$	90%
$\Gamma_{14}$	$3\pi^+ 3\pi^-$	(9 $\pm$ 4 )	$\times 10^{-5}$	
$\Gamma_{15}$	$3\pi^+ 3\pi^- 2\pi^0$	(1.2 $\pm$ 0.4 )	$\times 10^{-3}$	
$\Gamma_{16}$	$3\pi^+ 3\pi^- K^+ K^-$	(1.4 $\pm$ 0.7 )	$\times 10^{-4}$	
$\Gamma_{17}$	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	(4.2 $\pm$ 1.7 )	$\times 10^{-4}$	
$\Gamma_{18}$	$4\pi^+ 4\pi^-$	(9 $\pm$ 5 )	$\times 10^{-5}$	
$\Gamma_{19}$	$4\pi^+ 4\pi^- 2\pi^0$	(1.3 $\pm$ 0.5 )	$\times 10^{-3}$	

 **$\chi_{b2}(2P)$  BRANCHING RATIOS**

$\Gamma(\omega \Upsilon(1S))/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$			
<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.10^{+0.32+0.11}_{-0.28-0.10}</math></b>	$20.1^{+5.8}_{-5.1}$	5 CRONIN-HEN..04	CLE3	$\Upsilon(3S) \rightarrow \gamma \omega \Upsilon(1S)$

<sup>5</sup> Using  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.4 \pm 0.8)\%$  and  $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 2 (2.48 \pm 0.06)\%$ .

$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$			
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.089 \pm 0.012</math> OUR AVERAGE</b>				
0.085 $\pm$ 0.010 $\pm$ 0.010	6,7,8 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$	
0.084 $\pm$ 0.011 $\pm$ 0.010	2.5k <sup>9</sup> LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$	
0.096 $\pm$ 0.022 $\pm$ 0.012	7, <sup>10</sup> CRAWFORD	92B CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$	
0.106 $\pm$ 0.016 $\pm$ 0.013	7, <sup>11</sup> HEINTZ	92 CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$	

<sup>6</sup> LEES 14M quotes  $\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$  =  $(1.12 \pm 0.13)\%$  combining the results from samples of  $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$  with and without converted photons.

<sup>7</sup> Assuming  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$ .

<sup>8</sup> LEES 14M reports  $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (1.12 \pm 0.13) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>9</sup> LEES 11J reports  $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (1.1 \pm 0.1 \pm 0.1) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>10</sup> CRAWFORD 92B quotes  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \ell^+ \ell^-) = (4.98 \pm 0.94 \pm 0.62) 10^{-4}$ .

<sup>11</sup> Recalculated by us. HEINTZ 92 quotes  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) = (1.90 \pm 0.23 \pm 0.18) \%$  using  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$ . Supersedes HEINTZ 91.

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$					$\Gamma_3/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.066 ± 0.008 OUR AVERAGE</b>					
0.061 ± 0.004 ± 0.007	12,13,14	LEES	14M	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
0.070 ± 0.004 ± 0.008	11k	15 LEES	11J	BABR	$\Upsilon(3S) \rightarrow X\gamma$
0.077 ± 0.018 ± 0.009	13,16	CRAWFORD	92B	CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
0.061 ± 0.009 ± 0.007	13,17	HEINTZ	92	CSB2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
12 LEES 14M quotes $\Gamma(\chi_b2(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma\chi_b2(2P))/\Gamma_{\text{total}}$ = $(8.03 \pm 0.50) \times 10^{-3}$ combining the results from samples of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with and without converted photons.					
13 Assuming $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .					
14 LEES 14M reports $[\Gamma(\chi_b2(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_b2(2P))] = (8.03 \pm 0.50) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_b2(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.					
15 LEES 11J reports $[\Gamma(\chi_b2(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_b2(2P))] = (9.2 \pm 0.3 \pm 0.4) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_b2(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.					
16 CRAWFORD 92B quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_b2(2P)) \times B(\chi_b2(2P) \rightarrow \gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (5.03 \pm 0.94 \pm 0.63) 10^{-4}$ .					
17 Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_b2(2P)) \times B(\chi_b2(2P) \rightarrow \gamma \Upsilon(1S)) = (0.77 \pm 0.11 \pm 0.05)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.05)\%$ . Supersedes HEINTZ 91.					

$\Gamma(\pi\pi\chi_b2(1P))/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma$
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>5.1 ± 0.9 OUR AVERAGE</b>					
4.9 ± 0.7 ± 0.6	17k	18 LEES	11c	BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
6.0 ± 1.6 ± 1.4		19 CAWLFIELD 06	06	CLE3	$\Upsilon(3S) \rightarrow 2(\gamma\pi\ell)$
18 $(0.64 \pm 0.05 \pm 0.08) \times 10^{-3}$ . We derive the value assuming $B(\Upsilon(3S) \rightarrow \chi_b2(2P)X) = B(\Upsilon(3S) \rightarrow \chi_b2(2P)\gamma) = (13.1 \pm 1.6) \times 10^{-2}$ .					
19 CAWLFIELD 06 quote $\Gamma(\chi_b(2P) \rightarrow \pi\pi\chi_b(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$ keV assuming I-spin conservation, no $D$ -wave contribution, $\Gamma(\chi_b1(2P)) = 96 \pm 16$ keV, and $\Gamma(\chi_b2(2P)) = 138 \pm 19$ keV.					

$\Gamma(D^0 X)/\Gamma_{\text{total}}$					$\Gamma_5/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<2.4 × 10 <sup>-2</sup>					
<2.4 × 10 <sup>-2</sup>	90	20,21 BRIERE	08	CLEO	$\Upsilon(3S) \rightarrow \gamma D^0 X$
20 For $p_{D^0} > 2.5$ GeV/c.					
21 The authors also present their result as $(0.2 \pm 1.4 \pm 0.1) \times 10^{-2}$ .					

$\Gamma(\pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$					$\Gamma_6/\Gamma$
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;1.1</b>					
<1.1	90	22 ASNER	08A	CLEO	$\Upsilon(3S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$
22 ASNER 08A reports $[\Gamma(\chi_b2(2P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_b2(2P))] < 14 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_b2(2P)) = 13.1 \times 10^{-2}$ .					

$\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$					$\Gamma_7/\Gamma$
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;0.9</b>	90	23 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$	
$^{23} \text{ASNER 08A reports } [\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] < 12 \times 10^{-6} \text{ which we divide by our best value } B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}.$					

$\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$					$\Gamma_8/\Gamma$
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;7</b>	90	24 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$	
$^{24} \text{ASNER 08A reports } [\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] < 87 \times 10^{-6} \text{ which we divide by our best value } B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}.$					

$\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$					$\Gamma_9/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>3.9 \pm 1.6 \pm 0.5</math></b>	23	25 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$	
$^{25} \text{ASNER 08A reports } [\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (51 \pm 16 \pm 13) \times 10^{-6} \text{ which we divide by our best value } B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}. \text{ Our first error is their experiment's error and our second error is the systematic error from using our best value.}$					

$\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$					$\Gamma_{10}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>0.9 \pm 0.4 \pm 0.1</math></b>	11	26 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$	
$^{26} \text{ASNER 08A reports } [\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6} \text{ which we divide by our best value } B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}. \text{ Our first error is their experiment's error and our second error is the systematic error from using our best value.}$					

$\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$					$\Gamma_{11}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>2.4 \pm 1.0 \pm 0.3</math></b>	16	27 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$	
$^{27} \text{ASNER 08A reports } [\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (32 \pm 11 \pm 8) \times 10^{-6} \text{ which we divide by our best value } B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}. \text{ Our first error is their experiment's error and our second error is the systematic error from using our best value.}$					

$\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{12}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>4.7 \pm 2.2 \pm 0.6</math></b>	14	28 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	
$^{28} \text{ASNER 08A reports } [\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (62 \pm 23 \pm 17) \times 10^{-6} \text{ which we divide by our best value } B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}. \text{ Our first error is their experiment's error and our second error is the systematic error from using our best value.}$					

$\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;4</b>	90	29 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$

<sup>29</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] < 58 \times 10^{-6}$  which we divide by our best value  $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$ .

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.9±0.4±0.1</b>	14	30 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$

<sup>30</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6}$  which we divide by our best value  $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>12±4±1</b>	45	31 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$

<sup>31</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (159 \pm 33 \pm 43) \times 10^{-6}$  which we divide by our best value  $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.4±0.7±0.2</b>	12	32 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$

<sup>32</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (19 \pm 7 \pm 5) \times 10^{-6}$  which we divide by our best value  $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4.2±1.7±0.5</b>	16	33 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$

<sup>33</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (55 \pm 16 \pm 15) \times 10^{-6}$  which we divide by our best value  $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.9±0.4±0.1</b>	9	34 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$

<sup>34</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (12 \pm 5 \pm 3) \times 10^{-6}$  which we divide by our best value  $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$	$\Gamma_{19}/\Gamma$
<b>13±5±2</b>	<b>EVTS</b>
35 ASNER 08A	35 ASNER
27	08A CLEO
$\Gamma(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$	
35 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$	
$= (165 \pm 46 \pm 50) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))$	
$= (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.	

### $\chi_{b2}(2P)$ Cross-Particle Branching Ratios

$\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$	$\Gamma_3/\Gamma \times \Gamma_{20}^{\Upsilon(3S)}/\Gamma^{\Upsilon(3S)}$
<b>9.2±0.3±0.4</b>	<b>EVTS</b>
11k	LEES
DOCUMENT ID	11J BABR
COMMENT	$\Upsilon(3S) \rightarrow X\gamma$

$\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma \times \Gamma_{20}^{\Upsilon(3S)}/\Gamma^{\Upsilon(3S)}$
<b>1.1±0.1±0.1</b>	<b>EVTS</b>
2.5k	LEES
DOCUMENT ID	11J BABR
COMMENT	$\Upsilon(3S) \rightarrow X\gamma$

### $B(\chi_{b2}(2P) \rightarrow \chi_{b2}(1P)\pi^+\pi^-) \times B(\Upsilon(3S) \rightarrow \chi_{b2}(2P)X)$

$B(\chi_{b2}(2P) \rightarrow \chi_{b2}(1P)\pi^+\pi^-) \times B(\Upsilon(3S) \rightarrow \chi_{b2}(2P)X)$	$\Gamma_{20}^{\chi_{b2}(1P)\pi^+\pi^-}/\Gamma^{\chi_{b2}(1P)\pi^+\pi^-}$
<b>0.64±0.05±0.08</b>	<b>EVTS</b>
17k	LEES
DOCUMENT ID	11C BABR
COMMENT	$e^+ e^- \rightarrow \pi^+ \pi^- X$

### $B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\Upsilon(1S) \rightarrow \ell^+\ell^-)$

$B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\Upsilon(1S) \rightarrow \ell^+\ell^-)$	$\Gamma_{20}^{\gamma \chi_{b2}(2P)} \times \Gamma_{20}^{\Upsilon(1S)}/\Gamma^{\gamma \chi_{b2}(2P)} \times \Gamma^{\Upsilon(1S)}$
<b>2.02±0.18 OUR AVERAGE</b>	<b>EVTS</b>

1.95 <sup>+0.22</sup> <sub>-0.21</sub> <sup>+0.10</sup> <sub>-0.16</sub>	36 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
2.52 <sup>±0.47</sup> <sub>±0.32</sub>	48	37 CRAWFORD	92B CLE2 $\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$
1.98 <sup>±0.28</sup> <sub>±0.12</sub>		38 HEINTZ	92 CSB2 $\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$

36 From a sample of  $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$  with converted photons.

37 CRAWFORD 92B quotes  $2 \times B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma \Upsilon(nS)) B(\Upsilon(nS) \rightarrow \ell^+\ell^-)$ .

38 Calculated by us. HEINTZ 92 quotes  $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) = (0.77 \pm 0.11 \pm 0.05)\%$  using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.05)\%$ .

### $[B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] / [B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$

$[B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] / [B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$	$\Gamma_{20}^{\gamma \chi_{b2}(2P)} \times \Gamma_{20}^{\Upsilon(1S)}/\Gamma^{\gamma \chi_{b2}(2P)} \times \Gamma^{\Upsilon(1S)}$
<b>66.6±3.0</b>	<b>EVTS</b>
39 LEES	14M BABR
DOCUMENT ID	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
COMMENT	

39 From a sample of  $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$  events without converted photons.

**$B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\Upsilon(2S) \rightarrow \ell^+ \ell^-)$** 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.74±0.29 OUR AVERAGE</b>				
$3.22^{+0.58 +0.16}_{-0.53 -0.71}$	40	LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
$2.49 \pm 0.47 \pm 0.31$	53	41 CRAWFORD	92B CLE2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$
$2.74 \pm 0.33 \pm 0.18$		42 HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$
40 From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with converted photons.				
41 CRAWFORD 92B quotes $2 \times B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma \Upsilon(nS)) B(\Upsilon(nS) \rightarrow \ell^+\ell^-)$ .				
42 Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) = (1.90 \pm 0.23 \pm 0.18) \%$ using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$ .				
<b><math>[B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P))] / [B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]</math></b>				
VALUE (%)		DOCUMENT ID	TECN	COMMENT
<b>46.9±2.0</b>		43 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
43 From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ without converted photons.				

 **$\chi_{b2}(2P)$  REFERENCES**

AAIJ	14BG	JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
CAWLFIELD	06	PR D73 012003	C. Cawlfield <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN... 04	PRL 92 222002	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	
CRAWFORD	92B	PL B294 139	G. Crawford <i>et al.</i>	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)