

$\eta_c(2S)$

$I^G(J^{PC}) = 0^+(0^{-+})$

Quantum numbers are quark model predictions.

$\eta_c(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3637.7 ± 0.9 OUR AVERAGE		Error includes scale factor of 1.2.		
3637.90 ± 0.54 ± 1.40	3.7k	AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K\pi)$
3643.4 ± 2.3 ± 4.4	569	ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$
3635.1 ± 3.7 ± 2.9	106	XU	18 BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
3633.6 ± 1.7 ± 0.6	106	¹ AAIJ	17AD LHCb	$p p \rightarrow B^+ X \rightarrow p\bar{p} K^+ X$
3636.4 ± 4.1 ± 0.7	365	² AAIJ	17BB LHCb	$p p \rightarrow b\bar{b} X \rightarrow 2(K^+ K^-)X$
3637.0 ± 5.7 ± 3.4	178	^{3,4} LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^0$
3635.1 ± 5.8 ± 2.1	47	^{3,5} LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \eta$
3646.9 ± 1.6 ± 3.6	57	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
3637.6 ± 2.9 ± 1.6	127	⁶ ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K\pi, K K \pi^0$
3638.5 ± 1.5 ± 0.8	624	³ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
3640.5 ± 3.2 ± 2.5	1201	³ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
3636.1 ^{+3.9} _{-4.2} ^{+0.7} _{-2.0}	128	⁷ VINOKUROVA 11	BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
3626 ± 5 ± 6	311	⁸ ABE	07 BELL	$e^+ e^- \rightarrow J/\psi(c\bar{c})$
3645.0 ± 5.5 ^{+4.9} _{-7.8}	121	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
3642.9 ± 3.1 ± 1.5	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3639 ± 7	98	⁹ AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
3630.8 ± 3.4 ± 1.0	112	¹⁰ AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
3654 ± 6 ± 8	39	¹¹ CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
3594 ± 5		¹² EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

¹ AAIJ 17AD report $m_{\psi(2S)} - m_{\eta_c(2S)}$ = 52.5 ± 1.7 ± 0.6 MeV. We use the current value $m_{\psi(2S)} = 3686.097 \pm 0.025$ MeV to obtain the quoted mass.

² From a fit of the $\phi\phi$ invariant mass with the width of $\eta_c(2S)$ fixed to the PDG 16 value.

³ Ignoring possible interference with continuum.

⁴ With a width fixed to 11.3 MeV.

⁵ With a width fixed to 11.3 MeV. Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

⁶ From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes.

⁷ Accounts for interference with non-resonant continuum.

⁸ From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

⁹ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

¹⁰ Superseded by DEL-AMO-SANCHEZ 11M.

¹¹ Superseded by VINOKUROVA 11.

¹² Assuming mass of $\psi(2S) = 3686$ MeV.

$\eta_c(2S)$ WIDTH

VALUE (MeV)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
11.8 ± 1.6 OUR AVERAGE				
10.77 ± 1.62 ± 1.08	3.7k	AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K\pi)$
19.8 ± 3.9 ± 3.1	569	ABLIKIM	22Q BESS	$\psi(2S) \rightarrow \gamma 3(\pi^+\pi^-)$
9.9 ± 4.8 ± 2.9	57	ABLIKIM	13K BESS	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
16.9 ± 6.4 ± 4.8	127	¹ ABLIKIM	12G BESS	$\psi(2S) \rightarrow \gamma K^0 K\pi,$ $KK\pi^0$
13.4 ± 4.6 ± 3.2	624	² DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
6.6 ± 8.4 ± 2.6	128	³ VINOKUROVA 11	BELL	$B^\pm \rightarrow K^\pm(K_S^0 K^\pm \pi^\mp)$
6.3 ± 12.4 ± 4.0	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 23	90 98	⁴ AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
22 ± 14	121	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
17.0 ± 8.3 ± 2.5	112	⁵ AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
< 55	90 39	⁶ CHOI	02 BELL	$B \rightarrow KK_S K^- \pi^+$
< 8.0	95	⁷ EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

¹ From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes.

² Ignoring possible interference with continuum.

³ Accounts for interference with non-resonant continuum.

⁴ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

⁵ Superseded by DEL-AMO-SANCHEZ 11M.

⁶ For a mass value of 3654 ± 6 MeV. Superseded by VINOKUROVA 11.

⁷ For a mass value of 3594 ± 5 MeV

$\eta_c(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 hadrons	seen	
Γ_2 $K\bar{K}\pi$	(1.9 ± 1.2) %	
Γ_3 $K\bar{K}\eta$	(5 ± 4) × 10 ⁻³	
Γ_4 $2\pi^+ 2\pi^-$	< 2.1 %	90%
Γ_5 $\rho^0 \rho^0$	< 1.9 × 10 ⁻³	90%
Γ_6 $3\pi^+ 3\pi^-$	(1.3 ± 0.9) %	
Γ_7 $K^+ K^- \pi^+ \pi^-$	< 1.4 %	90%
Γ_8 $K^{*0} \bar{K}^{*0}$	< 2.9 × 10 ⁻³	90%
Γ_9 $K^+ K^- \pi^+ \pi^- \pi^0$	(1.4 ± 1.0) %	
Γ_{10} $K^+ K^- 2\pi^+ 2\pi^-$	< 1.4 %	90%
Γ_{11} $K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$	(1.0 ± 0.8) %	
Γ_{12} $2K^+ 2K^-$	< 1.3 × 10 ⁻³	90%

Γ_{13}	$\phi\phi$	<	1.1	$\times 10^{-3}$	90%
Γ_{14}	$p\bar{p}$	<	2.0	$\times 10^{-3}$	90%
Γ_{15}	$p\bar{p}\pi^+\pi^-$		seen		
Γ_{16}	$\gamma\gamma$		(1.8±1.2)	$\times 10^{-4}$	
Γ_{17}	$\gamma J/\psi(1S)$	<	1.4	%	90%
Γ_{18}	$\pi^+\pi^-\eta$		(4.3±3.2)	$\times 10^{-3}$	
Γ_{19}	$\pi^+\pi^-\eta'$		(2.6±1.9)	$\times 10^{-3}$	
Γ_{20}	$K_2^*(1430)\bar{K} + \text{c.c.}$		seen		
Γ_{21}	$K_0^*(1950)\bar{K} + \text{c.c.}$		seen		
Γ_{22}	$a_0(1710)\pi$		seen		
Γ_{23}	$a_0(1450)\pi$		seen		
Γ_{24}	$a_2(1700)\pi$		seen		
Γ_{25}	$K_0^*(2600)\bar{K} + \text{c.c.}$		seen		
Γ_{26}	$\pi^+\pi^-\eta_c(1S)$	< 25	%	90%	

 $\eta_c(2S)$ PARTIAL WIDTHS **$\Gamma(\gamma\gamma)$** **Γ_{16}**

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.44±0.14	106	¹ XU	18	BELL $e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
1.3 ± 0.6		² ASNER	04	CLEO $\gamma\gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$

¹ Assuming that the branching fraction into $\eta'\pi^+\pi^-$ is the same as for $\eta_c(1S)$.

² They measure $\Gamma(\eta_c(2S)\gamma\gamma) B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma) B(\eta_c(1S) \rightarrow K\bar{K}\pi)$. The value for $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$ is derived assuming that the branching fractions for $\eta_c(2S)$ and $\eta_c(1S)$ decays to $K_S K\pi$ are equal and using $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$ keV.

 $\Gamma(\gamma\gamma) \times \Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ **$\Gamma_{16}\Gamma_{19}/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.6^{+1.2}_{-1.1}±1.1	106	XU	18	BELL $e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$

 $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ **$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$** **$\Gamma_2\Gamma_{16}/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
41±4±6	624	¹ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$

¹ Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

 $\Gamma(2\pi^+2\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma_4\Gamma_{16}/\Gamma$**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<6.5	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+\pi^-)$

 $\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma_7\Gamma_{16}/\Gamma$**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.0	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+K^-\pi^+\pi^-$

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_9 \Gamma_{16}/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
30±6±5	1201	1 DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

¹ Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(2K^+ 2K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{12} \Gamma_{16}/\Gamma$			
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.9	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+ K^-)$

$\Gamma(\pi^+ \pi^- \eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{26} \Gamma_{16}/\Gamma$			
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<133	90	LEES	12AE	BABR $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{14}/\Gamma \times \Gamma_{16}/\Gamma$			
<u>VALUE (units 10^{-8})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 5.6	90	1,2,3 AMBROGIANI	01	E835 $\bar{p}p \rightarrow \gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 8.0	90	1,2,4 AMBROGIANI	01	E835 $\bar{p}p \rightarrow \gamma\gamma$
<12.0	90	2,4 AMBROGIANI	01	E835 $\bar{p}p \rightarrow \gamma\gamma$

¹ Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.

² For a total width $\Gamma=5$ MeV.

³ For the resonance mass region 3589 – 3599 MeV/ c^2 .

⁴ For the resonance mass region 3575 – 3660 MeV/ c^2 .

$\eta_c(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$	Γ_1/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	ABREU	980 DLPH	$e^+ e^- \rightarrow e^+ e^- + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	¹ EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

¹ For a mass value of 3594 ± 5 MeV

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$	Γ_2/Γ			
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.9±0.4±1.1	59 ± 12	¹ AUBERT	08AB	BABR $B \rightarrow \eta_c(2S) K \rightarrow K\bar{K}\pi K$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	127 ± 18	ABLIKIM	12G	BES3 $\psi(2S) \rightarrow \gamma K\bar{K}\pi$
seen	39 ± 11	² CHOI	02	BELL $B \rightarrow K K_S K^- \pi^+$

¹ Derived from a measurement of $[B(B^+ \rightarrow \eta_c(2S) K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$ and using $B(B^+ \rightarrow \eta_c(2S) K^+) = (3.4 \pm 1.8) \times 10^{-4}$, and $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$.

² For a mass value of 3654 ± 6 MeV

$\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$ Γ_3/Γ_2

<u>VALUE</u> (units 10^{-2})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
27.3±7.0±9.0	225	1 LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \gamma\gamma$

¹ LEES 14E reports $B(\eta_c(2S) \rightarrow K^+ K^- \eta)/B(\eta_c(2S) \rightarrow K^+ K^- \pi^0) = 0.82 \pm 0.21 \pm 0.27$, which we divide by 3 to account for isospin symmetry.

$\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	UEHARA	08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$
----------	--------	----	------	---------------------------------------

$\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$
----------	---------	----------	---

$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	UEHARA	08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$
----------	--------	----	------	---------------------------------------

$\Gamma(K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
----------	---------	----------	---

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$ Γ_9/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.73±0.17±0.17	1201	1 DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
-----------------------	------	-------------------	------	--

¹ We have multiplied the value of $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$ reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$. Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
------	-------------	---------	----------	---

$\Gamma(2K^+ 2K^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	UEHARA	08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$
----------	--------	----	------	---------------------------------------

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_{13}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$
----------	---------	----------	---

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	106	¹ AAIJ	17AD LHCb	$p p \rightarrow B^+ X \rightarrow p\bar{p} K^+ X$
¹ AAIJ 17AD report a 6.4 standard deviation signal, with $B(B^+ \rightarrow \eta_c(2S) K^+ \rightarrow p\bar{p} K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p} K^+) = (1.58 \pm 0.33 \pm 0.09) \times 10^{-2}$.				

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	110	¹ CHILIKIN	19	$e^+ e^- \rightarrow \gamma(4S)$
¹ CHILIKIN 19 reports signals in $B^+ \rightarrow \eta_c(2S) K^+$ and $B^0 \rightarrow \eta_c(2S) K_S^0$ with 12.3 and 5.9 standard deviations, respectively.				

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<4 $\times 10^{-4}$	90	¹ WICHT	08	$B^\pm \rightarrow K^\pm \gamma\gamma$
not seen		AMBROGIANI	01	$\bar{p}p \rightarrow \gamma\gamma$
<0.01	90	LEE	85	$\psi' \rightarrow \text{photons}$
¹ WICHT 08 reports $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S) K^+)] < 0.18 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c(2S) K^+) = 4.4 \times 10^{-4}$.				

$\Gamma(\pi^+\pi^-\eta_c(1S))/\Gamma(K\bar{K}\pi)$ Γ_{26}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.33	90	¹ LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

¹ We divided the reported limit by 3 to take into account isospin relations.

$\eta_c(2S)$ CROSS-PARTICLE BRANCHING RATIOS

$\Gamma(\eta_c(2S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<11.8 $\times 10^{-6}$	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \eta$
¹ CRONIN-HENNESSY 10 reports a limit of $< 5.9 \times 10^{-6}$ for the decay $\eta_c(2S) \rightarrow K^+ K^- \eta$ which we multiply by 2 account for isospin symmetry. It assumes $\Gamma(\eta_c(2S)) = 14$ MeV. It also gives the analytic dependence of limits on width.				

$\Gamma(\eta_c(2S) \rightarrow 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<14.6 $\times 10^{-6}$	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \rho^0 \rho^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_5/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<12.7 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

$$\Gamma(\eta_c(2S) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_6/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$9.2 \pm 1.0 \pm 1.2$	569		ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13.2	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$
---------	----	-----------------------------	------	---

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_7/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.6 \times 10^{-6}$	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_8/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<19.6 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_9/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<43.0 \times 10^{-6}$	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{10}/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-6}$	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{11}/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$7.03 \pm 2.10 \pm 0.7$	60		ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$$< 15.2 \quad 90 \quad {}^1 \text{CRONIN-HEN..10} \quad \text{CLEO} \quad \psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$$

¹ Assuming $\Gamma(\eta_c(2S)) = 14 \text{ MeV}$. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{13}/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.8 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

$$\Gamma(\eta_c(2S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{14}/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.4 \times 10^{-6}$	90	ABLIKIM	13V BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$

$$\Gamma(\eta_c(2S) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{17}/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 9.7 \times 10^{-6}$	90	33	{}^1 \text{ABLIKIM}	17N BES3	$\psi(2S) \rightarrow \gamma\gamma J/\psi$

¹ Uses $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{18}/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.97 \pm 0.81 \pm 0.26$	106		ABLIKIM	23Q BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$$< 4.3 \quad 90 \quad {}^1 \text{CRONIN-HEN..10} \quad \text{CLEO} \quad \psi(2S) \rightarrow \gamma\pi^+\pi^-\eta$$

¹ Assuming $\Gamma(\eta_c(2S)) = 14 \text{ MeV}$. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{19}/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
				• • • We do not use the following data for averages, fits, limits, etc. • • •

• • • We do not use the following data for averages, fits, limits, etc. • • •

$$< 14.2 \times 10^{-6} \quad 90 \quad {}^1 \text{CRONIN-HEN..10} \quad \text{CLEO} \quad \psi(2S) \rightarrow \gamma\pi^+\pi^-\eta'$$

¹ Assuming $\Gamma(\eta_c(2S)) = 14 \text{ MeV}$. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(K_2^*(1430)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$$

$$\Gamma_{20}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	{}^1 \text{AAIJ}	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K\pi)$

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$.

$\Gamma(a_0(1710)\pi)/\Gamma_{\text{total}}$	Γ_{22}/Γ			
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>			
seen	¹ AAIJ 23AH LHCb $B^+ \rightarrow K^+(K_S^0 K\pi)$			
¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$				
$\Gamma(a_0(1450)\pi)/\Gamma_{\text{total}}$	Γ_{23}/Γ			
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>			
seen	¹ AAIJ 23AH LHCb $B^+ \rightarrow K^+(K_S^0 K\pi)$			
¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$				
$\Gamma(a_2(1700)\pi)/\Gamma_{\text{total}}$	Γ_{24}/Γ			
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>			
seen	¹ AAIJ 23AH LHCb $B^+ \rightarrow K^+(K_S^0 K\pi)$			
¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$				
$\Gamma(K_0^*(2600)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{25}/Γ			
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>			
seen	¹ AAIJ 23AH LHCb $B^+ \rightarrow K^+(K_S^0 K\pi)$			
¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$				
$\Gamma(K_0^*(1950)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{21}/Γ			
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>			
seen	¹ AAIJ 23AH LHCb $B^+ \rightarrow K^+(K_S^0 K\pi)$			
¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$				
$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$	$\Gamma_{26}/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma^{\psi(2S)}$			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.7 × 10⁻⁴	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta_c(1S)$
¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.				

$\eta_c(2S)$ REFERENCES

AAIJ	23AH PR D108 032010	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	23Q PR D107 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22Q PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
CHILIKIN	19 PR D100 012001	K. Chilikin <i>et al.</i>	(BELLE Collab.)
XU	18 PR D98 072001	Q.N. Xu <i>et al.</i>	(BELLE Collab.)
AAIJ	17AD PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BB EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17N PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16 CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
LEES	14E PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13K PR D87 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13V PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12G PRL 109 042003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	12AE PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	11H PR D84 091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DEL-AMO-SA... 11M PR D84 012004	P. del Amo Sanchez <i>et al.</i>		(BABAR Collab.)

VINOKUROVA	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05C	PR D72 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)