

**$f_0(2020)$** 

$$I^G(J^{PC}) = 0^+(0^{++})$$

Needs confirmation.

 **$f_0(2020)$  T-MATRIX POLE  $\sqrt{s}$** Note that  $\Gamma = -2 \text{Im}(\sqrt{s})$ .

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1870–2080) – <math>i</math> (120–240) OUR ESTIMATE</b>			
$(2038 \pm 48) - i(156 \pm 41)$	<sup>1</sup> RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K})$
$(1925 \pm 25) - i(160 \pm 18)$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$(1910 \pm 50) - i(199 \pm 40)$	<sup>2</sup> ROPERTZ	18	RVUE $\bar{B}_s^0 \rightarrow J/\psi(\pi^+\pi^-/K^+K^-)$
$(1992 \pm 16) - i(221 \pm 30)$	<sup>3</sup> BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_s$
$(2020 \pm 35) - i(205 \pm 25)$	BARBERIS	97B	OMEG 450 $pp \rightarrow pp2(\pi^+\pi^-)$

<sup>1</sup> T-matrix pole from coupled channel K-matrix fit to data on  $J/\psi \rightarrow \gamma\pi^0\pi^0$  (ABLIKIM 15AE) and  $J/\psi \rightarrow \gamma K_S^0 K_S^0$  (ABLIKIM 18AA).

<sup>2</sup> T-matrix pole of 3 channel unitary model fit to data from AAIJ 14BR and AAIJ 17V extracted using Pade approximants.

<sup>3</sup> Average between  $\pi^+\pi^-2\pi^0$  and  $2(\pi^+\pi^-)$ .

 **$f_0(2020)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$1982 \pm 3_{-0}^{+54}$		<sup>1</sup> ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$
$2010 \pm 6_{-4}^{+6}$		<sup>2</sup> ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
$2037 \pm 8$	80k	<sup>3</sup> UMAN	06 E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
$2040 \pm 38$		ANISOVICH	00J SPEC	
$2010 \pm 60$		ALDE	98 GAM4	$100 \pi^- p \rightarrow \pi^0\pi^0 n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> From a partial wave analysis of the systems  $(\gamma X)$ , with  $X \rightarrow \eta'\eta'$ , and  $(\eta' X)$ , with  $X \rightarrow \gamma\eta'$  in the decay  $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

<sup>2</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta\eta'$   $P$ -wave.

<sup>3</sup> Statistical error only.

 **$f_0(2020)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$436 \pm 4_{-49}^{+46}$		<sup>1</sup> ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$203 \pm 9^{+13}_{-11}$		<sup>2</sup> ABLIKIM	22AS	BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
$296 \pm 17$	80k	<sup>3</sup> UMAN	06	E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
$405 \pm 40$		ANISOVICH	00J	SPEC	
$240 \pm 100$		ALDE	98	GAM4	$100 \pi^- p \rightarrow \pi^0 \pi^0 n$

<sup>1</sup>From a partial wave analysis of the systems  $(\gamma X)$ , with  $X \rightarrow \eta'\eta'$ , and  $(\eta' X)$ , with  $X \rightarrow \gamma\eta'$  in the decay  $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

<sup>2</sup>From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta\eta'$   $P$ -wave.

<sup>3</sup>Statistical error only.

## $f_0(2020)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\rho\pi\pi$	seen
$\Gamma_2$ $\pi^0\pi^0$	seen
$\Gamma_3$ $\rho\rho$	seen
$\Gamma_4$ $\omega\omega$	seen
$\Gamma_5$ $\eta\eta$	seen
$\Gamma_6$ $\eta'\eta'$	seen

## $f_0(2020)$ BRANCHING RATIOS

$\Gamma(\rho\rho)/\Gamma(\omega\omega)$				$\Gamma_3/\Gamma_4$
VALUE	DOCUMENT ID	TECN	COMMENT	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\sim 3$	BARBERIS	00F	450	$pp \rightarrow p_f \omega \omega p_s$
----------	----------	-----	-----	--

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$				$\Gamma_5/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	

seen	UMAN	06	E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
------	------	----	------	--

$\Gamma(\eta'\eta')/\Gamma_{\text{total}}$				$\Gamma_6/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	

seen	<sup>1</sup> ABLIKIM	22C	BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$
------	----------------------	-----	------	---

<sup>1</sup>From a partial wave analysis of the systems  $(\gamma X)$ , with  $X \rightarrow \eta'\eta'$ , and  $(\eta' X)$ , with  $X \rightarrow \gamma\eta'$  in the decay  $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

**$f_0(2020)$  REFERENCES**

ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D107 079901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ROPERTZ	18	EPJ C78 1000	S. Ropertz, C. Hanhart, B. Kubis	(BONN, JULI)
AAIJ	17V	JHEP 1708 037	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	14BR	PR D89 092006	R. Aaij <i>et al.</i>	(LHCb Collab.)
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00F	PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 62 446.		
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)

---