

$\phi(2170)$

$I^G(J^{PC}) = 0^-(1^{--})$

See the review on "Spectroscopy of Light Meson Resonances."

$\phi(2170)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2164 ± 6 OUR AVERAGE				
2178 ± 20 ± 5	1	ABLIKIM	23AX BES3	$e^+ e^- \rightarrow \phi \pi^+ \pi^-$
2190 ± 19 ± 37	2	ABLIKIM	22L BES3	$e^+ e^- \rightarrow K^+ K^- \pi^0$
2176 ± 24 ± 3	3	ABLIKIM	21A BES3	$e^+ e^- \rightarrow \omega \eta$
2163.5 ± 6.2 ± 3.0	4	ABLIKIM	21T BES3	$e^+ e^- \rightarrow \phi \eta$
2177.5 ± 4.8 ± 19.5	5	ABLIKIM	20M BES3	$e^+ e^- \rightarrow \eta' \phi$
2126.5 ± 16.8 ± 12.4	6	ABLIKIM	20S BES3	$e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2215.7 ± 8.3	7	LICHARD	23 RVUE	$e^+ e^- \rightarrow \gamma(nS) \rightarrow \phi \eta \gamma$
2169 ± 5 ± 6	8	ZHU	23A RVUE	$e^+ e^- \rightarrow \eta \phi$
2273.7 ± 5.7 ± 19.3	9	ABLIKIM	21AP BES3	$e^+ e^- \rightarrow K_S^0 K_L^0$
2135 ± 8 ± 9	95	ABLIKIM	19I BES3	$e^+ e^- \rightarrow \eta \phi f_0(980)$
2239.2 ± 7.1 ± 11.3	10	ABLIKIM	19L BES3	$e^+ e^- \rightarrow K^+ K^-$
2200 ± 6 ± 5	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta \phi \pi^+ \pi^-$
2180 ± 8 ± 8	11,12	LEES	12F BABR	$10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$
2079 ± 13 +79 -28	4.8k	13 SHEN	09 BELL	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
2186 ± 10 ± 6	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta \phi f_0(980)$
2125 ± 22 ± 10	483	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \phi \eta \gamma$
2192 ± 14	116	14 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
2169 ± 20	149	14 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$
2175 ± 10 ± 15	201	12,15 AUBERT,BE	06D BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi \pi \gamma$

¹ From a fit to the $e^+ e^-$ cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a non-resonant contribution.

² By a simultaneous fit of the $K_2^*(1430)^+ K^-$ and $K^*(892)^+ K^-$ intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

³ From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from $\omega(1420)$ and $\omega(1650)/\phi(1680)$.

⁴ From a fit to the cross section below 3.5 GeV measured by BaBar and BESIII with a coherent sum of two modified Breit-Wigner amplitudes ($\phi(1680)$ and $\phi(2170)$) and a nonresonant term.

⁵ From a fit using a coherent sum of a phase-space modified Breit-Wigner function and a phase-space term.

⁶ By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

- ⁷ From a VDM fit to ZHU 23 $\eta\phi\gamma$ data with two resonances, $\phi(1680)$, $\phi(2170)$, and a third resonance with mass 1850.7 ± 5.3 MeV and width 25 ± 35 MeV of 1.7σ statistical evidence.
- ⁸ From the analysis of the combined measurements of $\sigma(e^+e^- \rightarrow \eta\phi)$ from BaBar, Belle, BESIII, CMD3. The statistical significance for $\phi(2170)$ is 7.2σ .
- ⁹ From a fit to the cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a nonresonant contribution. The observed structure can be also due to $\rho(2150)$.
- ¹⁰ The observed structure can be due to both the $\phi(2170)$ and $\rho(2150)$.
- ¹¹ Fit includes interference with the $\phi(1680)$.
- ¹² From the $\phi f_0(980)$ component.
- ¹³ From a fit with two incoherent Breit-Wigners.
- ¹⁴ From the $K^+K^-f_0(980)$ component.
- ¹⁵ Superseded by LEES 12F.


$\phi(2170)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
106 $^{+24}_{-18}$ OUR AVERAGE				Error includes scale factor of 2.0. See the ideogram below.
140 ± 36 ± 16		1 ABLIKIM	23AX BES3	$e^+e^- \rightarrow \phi\pi^+\pi^-$
191 ± 28 ± 60		2 ABLIKIM	22L BES3	$2.0\text{--}3.08 e^+e^- \rightarrow K^+K^-\pi^0$
89 ± 50 ± 5		3 ABLIKIM	21A BES3	$e^+e^- \rightarrow \omega\eta$
$31.1^{+21.1}_{-11.6} \pm 1.1$		4 ABLIKIM	21T BES3	$e^+e^- \rightarrow \phi\eta$
$149.0 \pm 15.6 \pm 8.9$		5 ABLIKIM	20M BES3	$e^+e^- \rightarrow \eta'\phi$
$106.9 \pm 32.1 \pm 28.1$		6 ABLIKIM	20S BES3	$e^+e^- \rightarrow K^+K^-\pi^0\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
35 ± 23		7 LICHARD	23 RVUE	$e^+e^- \rightarrow \gamma(nS) \rightarrow \phi\eta\gamma$
96^{+17}_{-14} ± 9		8 ZHU	23A RVUE	$e^+e^- \rightarrow \eta\phi$
86 ± 44 ± 51		9 ABLIKIM	21AP BES3	$e^+e^- \rightarrow K_S^0 K_L^0$
104 ± 24 ± 12	95	ABLIKIM	19I BES3	$e^+e^- \rightarrow \eta\phi f_0(980)$
$139.8 \pm 12.3 \pm 20.6$		10 ABLIKIM	19L BES3	$e^+e^- \rightarrow K^+K^-$
104 ± 15 ± 15	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta\phi\pi^+\pi^-$
77 ± 15 ± 10		11,12 LEES	12F BABR	$10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
192 ± 23 $^{+25}_{-61}$	4.8k	13 SHEN	09 BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
65 ± 23 ± 17	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$
61 ± 50 ± 13	483	AUBERT	08S BABR	$10.6 e^+e^- \rightarrow \phi\eta\gamma$
71 ± 21	116	14 AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
102 ± 27	149	14 AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
58 ± 16 ± 20	201	12,15 AUBERT,BE	06D BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi\pi\gamma$

¹ From a fit to the e^+e^- cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a non-resonant contribution.

² By a simultaneous fit of the $K_2^*(1430)^+K^-$ and $K^*(892)^+K^-$ intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

- ³ From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from $\omega(1420)$ and $\omega(1650)/\phi(1680)$.
⁴ From a fit to the cross section below 3.5 GeV measured by BaBar and BESIII with a coherent sum of two modified Breit-Wigner amplitudes ($\phi(1680)$ and $\phi(2170)$) and a nonresonant term.
⁵ From a fit using a coherent sum of a phase-space modified Breit-Wigner function and a phase-space term.
⁶ By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.
⁷ From a VDM fit to ZHU 23 $\eta\phi\gamma$ data with two resonances, $\phi(1680)$, $\phi(2170)$, and a third resonance with mass 1850.7 ± 5.3 MeV and width 25 ± 35 MeV of 1.7σ statistical evidence.
⁸ From the analysis of the combined measurements of $\sigma(e^+e^- \rightarrow \eta\phi)$ from BaBar, Belle, BESIII, CMD3. The statistical significance for $\phi(2170)$ is 7.2σ .
⁹ From a fit to the cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a nonresonant contribution. The observed structure can be also due to $\rho(2150)$.
¹⁰ The observed structure can be due to both the $\phi(2170)$ and $\rho(2150)$.
¹¹ Fit includes interference with the $\phi(1680)$.
¹² From the $\phi f_0(980)$ component.
¹³ From a fit with two incoherent Breit-Wigners.
¹⁴ From the $K^+K^- f_0(980)$ component.
¹⁵ Superseded by LEES 12F.



$\phi(2170)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 e^+ e^-$	seen
$\Gamma_2 \phi\eta$	seen
$\Gamma_3 \omega\eta$	seen
$\Gamma_4 \phi\eta'$	seen
$\Gamma_5 \phi\pi\pi$	seen
$\Gamma_6 \phi f_0(980)$	seen
$\Gamma_7 K_S^0 K_L^0$	
$\Gamma_8 K^+ K^- \pi^+ \pi^-$	
$\Gamma_9 K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-$	seen
$\Gamma_{10} K^+ K^- \pi^0 \pi^0$	
$\Gamma_{11} K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0$	seen
$\Gamma_{12} K^{*0} K^\pm \pi^\mp$	not seen
$\Gamma_{13} K^*(892)^0 \bar{K}^*(892)^0$	not seen
$\Gamma_{14} K^*(892)^+ K^*(892)^-$	
$\Gamma_{15} K^*(892)^+ K^- + \text{c.c.}$	
$\Gamma_{16} K(1460)^+ K^- + \text{c.c.}$	
$\Gamma_{17} K_1(1270)^+ K^- + \text{c.c.}$	
$\Gamma_{18} K_1(1400)^+ K^- + \text{c.c.}$	
$\Gamma_{19} K_2^*(1430)^+ K^- + \text{c.c.}$	

$\phi(2170) \Gamma(i) \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(\phi\eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_2 \Gamma_1 / \Gamma$
<i>VALUE (eV)</i>	<i>CL%</i>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.17	90
$0.36^{+0.05}_{-0.03} \pm 0.07$ to $41 \pm 2 \pm 6$	¹ ZHU 23 BELL $e^+ e^- \rightarrow \gamma(nS) \rightarrow \phi\eta\gamma$
$0.24^{+0.12}_{-0.07}$	² ZHU 23A RVUE $e^+ e^- \rightarrow \eta\phi$
$1.7 \pm 0.7 \pm 1.3$	483 AUBERT 08S BABR $10.6 e^+ e^- \rightarrow \phi\eta\gamma$

¹ From a solution of the fit using a vector meson dominance model with contributions from $\phi(1680)$, $\phi(2170)$ and non resonant contribution with mass and width of $\phi(2170)$ fixed at 2163.5 MeV and 31.1 MeV respectively. Four solutions are found with equal fit quality giving 0.17 eV (solution I and II) and 18.6 eV (III and IV) at 90% CL.

² From the analysis of the combined measurements of $\sigma(e^+ e^- \rightarrow \eta\phi)$ from BaBar, Belle, BESIII, CMD3. The statistical significance for $\phi(2170)$ is 7.2σ . Four solutions are found, with equal fit quality: $(0.56^{+0.03}_{-0.02} \pm 0.07)$ eV, $(0.36^{+0.05}_{-0.03} \pm 0.07)$ eV, $(38 \pm 1 \pm 5)$ eV, $(41 \pm 2 \pm 6)$ eV.

³ From a solution of the fit to the cross section below 3.5 GeV measured by BaBar and BESIII with a coherent sum of two modified Breit-Wigner amplitudes ($\phi(1680)$ and $\phi(2170)$) and a nonresonant term. The other solution gives $10.11^{+3.87}_{-3.13}$ eV.

$\Gamma(\omega\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_3\Gamma_1/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
0.43±0.15±0.04	¹ ABLIKIM	21A BES3	$e^+e^- \rightarrow \omega\eta$

¹ For constructive interference with $\omega(1420)$ and $\omega(1650)/\phi(1680)$. For destructive interference: $1.25 \pm 0.48 \pm 0.18$ eV.

$\Gamma(\phi\eta') \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_4\Gamma_1/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
7.1±0.7±0.7	¹ ABLIKIM	20M BES3	$e^+e^- \rightarrow \eta'\phi$

¹ From a fit using a coherent sum of a phase-space modified Breit-Wigner function and a phase-space term.

$\Gamma(\phi f_0(980)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_6\Gamma_1/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2.3 \pm 0.3 \pm 0.3$	1,2 LEES	12F BABR	$10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$	

$2.5 \pm 0.8 \pm 0.4$ 201 2,³ AUBERT,BE 06D BABR $10.6 e^+e^- \rightarrow K^+K^-\pi\pi\gamma$

¹ From a fit with constructive interference with the $\phi(1680)$. In a fit with destructive interference, the value is larger by a factor of 12.

² For $f_0(980) \rightarrow \pi\pi$.

³ Superseded by LEES 12F.

$\Gamma(K_S^0 K_L^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_7\Gamma_1/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.9 \pm 0.6 \pm 0.7$	¹ ABLIKIM	21AP BES3	$e^+e^- \rightarrow K_S^0 K_L^0$

¹ From a fit to the cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a nonresonant contribution. The observed structure can be also due to $\rho(2150)$.

$\Gamma(K^*(892)^+ K^*(892)^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{14}\Gamma_1/\Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.9	90	¹ ABLIKIM	20S BES3	$e^+e^- \rightarrow K^+K^-\pi^0\pi^0$

¹ By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

$\Gamma(K^*(892)^+ K^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{15}\Gamma_1/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.0 ± 0.3	¹ ABLIKIM	22L BES3	$2.0-3.08 e^+e^- \rightarrow K^+K^-\pi^0$

¹ From a solution of a simultaneous fit of the $K_2^*(1430)^+ K^-$ and $K^*(892)^+ K^-$ intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. The other solution gives 7.1 ± 0.9 eV. Significance 3.7 σ .

$\Gamma(K(1460)^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{16}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
3.0 \pm 3.8	¹ ABLIKIM	20S BES3	$e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0$

¹ By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

 $\Gamma(K_1(1270)^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{17}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<12.5	90	¹ ABLIKIM	20S BES3	$e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0$

¹ By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. A second solution of the fit with equal fit quality gives an upper limit value of 297.6 eV.

 $\Gamma(K_1(1400)^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{18}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			

4.7 \pm 3.3 ¹ ABLIKIM 20S BES3 $e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0$

¹ By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. A second solution of the fit with equal fit quality gives a value of 98.8 ± 7.8 eV.

 $\Gamma(K_2^*(1430)^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{19}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			

12.6 \pm 2.4 ¹ ABLIKIM 22L BES3 2.0–3.08 $e^+ e^- \rightarrow K^+ K^- \pi^0$

¹ From a solution of a simultaneous fit of the $K_2^*(1430)^+ K^-$ and $K^*(892)^+ K^-$ intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. The other solution gives 161.1 ± 20.6 eV.

 $\phi(2170) \Gamma(i)\Gamma(e^+ e^-)/\Gamma^2(\text{total})$
 $\Gamma(\phi\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma \times \Gamma_1/\Gamma$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				

1.65 \pm 0.15 \pm 0.18 4.8k ¹ SHEN 09 BELL 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

¹ Multiplied by 3/2 to take into account the $\phi\pi^0\pi^0$ mode. Using $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$.

$\phi(2170)$ BRANCHING RATIOS

$\Gamma(\phi\pi\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
seen	1 ABLIKIM	23AX BES3	$e^+ e^- \rightarrow \phi\pi^+\pi^-$	

¹ From a fit to the $e^+ e^-$ cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a non-resonant contribution.

$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
seen	AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$	

$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
seen	AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$	

$\Gamma(K^{*0} K^\pm \pi^\mp)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ
not seen	AUBERT	07AK BABR	$10.6 \text{ GeV } e^+ e^-$	

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
not seen	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$	

$\phi(2170)$ REFERENCES

ABLIKIM	23AX PR D108 032011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LICHARD	23 PR D108 092005	P. Lichard	(OPAV, CTUP)
ZHU	23 PR D107 012006	W. Zhu <i>et al.</i>	(BELLE Collab.)
ZHU	23A CP C47 113003	W. Zhu, X. Wang	(RVUE)
ABLIKIM	22L JHEP 2207 045	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21A PL B813 136059	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AP PR D104 092014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21T PR D104 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20M PR D102 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20S PRL 124 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19I PR D99 012014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19L PR D99 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15H PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	12F PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	10C PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
SHEN	09 PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08F PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	08S PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AK PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)