

$\psi_2(3823)$	$I^G(J^{PC}) = 0^-(2^{--})$
	I, J, P need confirmation.

was $\psi(3823)$, $X(3823)$

Seen by BHARDWAJ 13 in $B \rightarrow \chi_{c1}\gamma K$ and ABLIKIM 15S in $e^+e^- \rightarrow \pi^+\pi^-\gamma\chi_{c1}$ decays as a narrow peak in the invariant mass distribution of the $\chi_{c1}\gamma$ system. Properties consistent with the $\psi_2(1^3D_2)$ $c\bar{c}$ state.

$\psi_2(3823)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3823.51 ± 0.34 OUR AVERAGE				
3824.5 ± 2.4 ± 1.0	30	¹ ABLIKIM	23J BES3	$e^+e^- \rightarrow \pi^0\pi^0\chi_{c1}\gamma$
$3823.12 \pm 0.43 \pm 0.13$	120	ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
$3824.08 \pm 0.53 \pm 0.14$	137	² AAIJ	20S LHCb	$B^+ \rightarrow J/\psi\pi^+\pi^-K^+$
$3823.1 \pm 1.8 \pm 0.7$	33 ± 10	³ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1}\gamma K^\pm$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3821.7 $\pm 1.3 \pm 0.7$ 19 ± 5 ⁴ ABLIKIM 15S BES3 $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$

¹ Using the measured $m_{\psi_2(3823)} - m_{\psi(2S)}$ and assuming $m_{\psi(2S)} = 3686.097$ MeV from PDG 22.

² Using the measured $m_{\psi_2(3823)} - m_{\psi(2S)} = 137.98 \pm 0.53 \pm 0.14$ MeV.

³ From a simultaneous fit to $B^\pm \rightarrow (\chi_{c1}\gamma)K^\pm$ and $B^0 \rightarrow (\chi_{c1}\gamma)K_S^0$ with significance 4.0σ including systematics. Corrected for the measured $\psi(2S)$ mass using $B \rightarrow \psi(2S)K \rightarrow (\gamma\chi_{c1})K$ decays.

⁴ From a simultaneous unbinned maximum likelihood fit of $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$ data (the $\pi^+\pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c1}\gamma$ and $\psi_2(3823) \rightarrow \chi_{c1}\gamma$ together, with floating mass scale offset for $\psi(2S)$, floating $\psi_2(3823)$ mass, and zero $\psi_2(3823)$ width, resulting in a significance of 5.9σ when including systematic uncertainties. Superseded by ABLIKIM 22R.

$m_{\psi_2(3823)} - m_{\psi(2S)}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$137.98 \pm 0.53 \pm 0.14$ 137 ¹ AAIJ 20S LHCb $B^+ \rightarrow J/\psi\pi^+\pi^-K^+$

¹ AAIJ 20S also reports $m_{\chi_{c1}(3872)} - m_{\psi_2(3823)} = 47.50 \pm 0.53 \pm 0.13$ MeV.

$\psi_2(3823)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 2.9	90	120	ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<18.8	90	30	¹ ABLIKIM	23J BES3	$e^+e^- \rightarrow \pi^0\pi^0\chi_{c1}\gamma$
< 5.2	90		² AAIJ	20S LHCb	$B^+ \rightarrow J/\psi\pi^+\pi^-K^+$
<16	90		³ ABLIKIM	15S BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
<24	90		⁴ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1}\gamma K^\pm$

- ¹ From a fit of $e^+ e^- \rightarrow \pi^0 \pi^0 \chi_{c1} \gamma$ data at \sqrt{s} values from 4.23 to 4.70 GeV to a Breit-Wigner function with floating width, using the Bayesian approach.
- ² AAIJ 20S also provides a limit of < 6.6 MeV with 95% CL.
- ³ From a fit of $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$ data (the $\pi^+ \pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to a Breit-Wigner function with the mass fixed from the likelihood fit above, Gaussian resolution smearing, and floating width.
- ⁴ From a simultaneous fit to $B^\pm \rightarrow (\chi_{c1} \gamma) K^\pm$ and $B^0 \rightarrow (\chi_{c1} \gamma) K_S^0$ with significance 4.0σ including systematics.

$\psi_2(3823)$ DECAY MODES

Branching fractions are given relative to the one **DEFINED AS 1**.

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $J/\psi(1S)\pi^+\pi^-$	<0.06	90%
Γ_2 $J/\psi(1S)\pi^0\pi^0$	<0.11	90%
Γ_3 $J/\psi(1S)\pi^0$	<0.030	90%
Γ_4 $J/\psi(1S)\eta$	<0.14	90%
Γ_5 $\chi_{c0}\gamma$	<0.24	90%
Γ_6 $\chi_{c1}\gamma$	DEFINED AS 1	
Γ_7 $\chi_{c2}\gamma$	$0.28^{+0.14}_{-0.11}$	

$\psi_2(3823)$ BRANCHING RATIOS

$$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen		¹ ABLIKIM	210 BES3	$e^+ e^- \rightarrow \pi^+ \pi^- X$
seen	137 ± 26	AAIJ	20S LHCb	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$

- ¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+ \pi^-$ recoil mass distributions of seven decay channels in the process $e^+ e^- \rightarrow \pi^+ \pi^- X$.

$$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(\chi_{c1}\gamma) \quad \Gamma_1/\Gamma_6$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.06	90	¹ ABLIKIM	210 BES3	$e^+ e^- \rightarrow \pi^+ \pi^- X$

- ¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+ \pi^-$ recoil mass distributions of seven decay channels in the process $e^+ e^- \rightarrow \pi^+ \pi^- X$.

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(\chi_{c1}\gamma) \quad \Gamma_2/\Gamma_6$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.11	90	¹ ABLIKIM	210 BES3	$e^+ e^- \rightarrow \pi^+ \pi^- X$

- ¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+ \pi^-$ recoil mass distributions of seven decay channels in the process $e^+ e^- \rightarrow \pi^+ \pi^- X$.

$$\Gamma(J/\psi(1S)\pi^0)/\Gamma(\chi_{c1}\gamma) \quad \Gamma_3/\Gamma_6$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.03	90	¹ ABLIKIM	210 BES3	$e^+ e^- \rightarrow \pi^+ \pi^- X$

- ¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+ \pi^-$ recoil mass distributions of seven decay channels in the process $e^+ e^- \rightarrow \pi^+ \pi^- X$.

$\Gamma(J/\psi(1S)\eta)/\Gamma(\chi_{c1}\gamma)$	Γ_4/Γ_6			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.14	90	¹ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$	Γ_4/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
$4.4^{+2.5}_{-1.9} \pm 0.9$	¹ AAIJ	22D LHCb	$B^+ \rightarrow J/\psi(1S)\eta K^+$

¹ Using the branching ratio for $B^+ \rightarrow \psi_2(3823)K^+$ with $\psi_2(3823) \rightarrow J/\psi(1S)\pi^+\pi^-$ from AAIJ 20S.

$\Gamma(\chi_{c0}\gamma)/\Gamma_{\text{total}}$	Γ_5/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

not seen	¹ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.			

$\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$	Γ_6/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	120	¹ ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
seen	63 ± 9	² ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
seen	16 ± 5	³ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^0\pi^0X$
seen	33 ± 10	⁴ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1}\gamma K^\pm$

¹ From a fit to the $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak R with mass $4417.5 \pm 26.2 \pm 3.5$ MeV and width $245 \pm 48 \pm 13$ MeV, which leads to $\Gamma(e^+e^-)B(R \rightarrow \pi^+\pi^-\psi_2(3823))B(\psi_2(3823) \rightarrow \chi_{c1}\gamma) = 0.57 \pm 0.08$ eV.

² From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$. Signal has a 11.8σ significance.

³ From a fit of the invariant $\pi^0\pi^0$ recoil-mass distribution. Signal has a 4.3σ significance.

⁴ BHARDWAJ 13 reports $B(B^\pm \rightarrow \psi_2(3823)K^\pm) \times B(\psi_2(3823) \rightarrow \gamma\chi_{c1}) = (9.7 \pm 2.8 \pm 1.1) \times 10^{-6}$ with statistical significance 3.8σ .

$\Gamma(\chi_{c0}\gamma)/\Gamma(\chi_{c1}\gamma)$	Γ_5/Γ_6			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.24	90	¹ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

$\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$	Γ_7/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	¹ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
not seen	² ABLIKIM	15S BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$
not seen	³ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c2}\gamma K^\pm$

- ¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^- X$. Signal has a 3.2σ significance.
- ² From a simultaneous unbinned maximum likelihood fit of $e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$ data (the $\pi^+\pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c2}\gamma$ and $\psi_2(3823) \rightarrow \chi_{c2}\gamma$ together, with floating mass scale offset for $\psi(2S)$, $\psi_2(3823)$ mass floating (fixed to that above), and zero $\psi_2(3823)$ width.
- ³ BHARDWAJ 13 reports $B(B^\pm \rightarrow \psi_2(3823)K^\pm) \times B(\psi_2(3823) \rightarrow \gamma\chi_{c2}) < 3.6 \times 10^{-6}$ at 90% CL.

$\Gamma(\chi_{c2}\gamma)/\Gamma(\chi_{c1}\gamma)$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ_6
$0.28^{+0.14}_{-0.11} \pm 0.02$		9 ± 4	¹ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.42	90	² ABLIKIM	15S BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$
<0.41	90	BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1/c2}\gamma K^\pm$

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^- X$.

² From a simultaneous unbinned maximum likelihood fit of $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1(2)}\gamma$ data (the $\pi^+\pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c1(2)}\gamma$ and $\psi_2(3823) \rightarrow \chi_{c1(2)}\gamma$ together, with floating mass scale offset for $\psi(2S)$, $\psi_2(3823)$ mass floating (fixed to that above), and zero $\psi_2(3823)$ width.

$\psi_2(3823)$ REFERENCES

ABLIKIM	23J	JHEP 2302 171	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	22D	JHEP 2204 046	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	22R	PRL 129 102003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)
ABLIKIM	21O	PR D103 L091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	20S	JHEP 2008 123	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)