

$\psi(4415)$

$I^G(J^{PC}) = 0^-(1^{--})$

$\psi(4415)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4415 ± 5 OUR AVERAGE			
4414.6 ± 3.4 ± 6.1	ABLIKIM	23BH BES3	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-}$
4415.1 ± 7.9	¹ ABLIKIM	08D BES2	$e^+ e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4412 ± 15	² MO	10 RVUE	$e^+ e^- \rightarrow \text{hadrons}$
4411 ± 7	³ PAKHLOVA	08A BELL	$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$
4425 ± 6	⁴ SETH	05A RVUE	$e^+ e^- \rightarrow \text{hadrons}$
4429 ± 9	⁵ SETH	05A RVUE	$e^+ e^- \rightarrow \text{hadrons}$
4417 ± 10	BRANDELIK	78C DASP	$e^+ e^-$
4414 ± 7	SIEGRIST	76 MRK1	$e^+ e^-$
¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.			
² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
³ Systematic uncertainties not estimated.			
⁴ From a fit to Crystal Ball (OSTERHELD 86) data.			
⁵ From a fit to BES (BAI 02C) data.			

$\psi(4415)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
110 ± 22 OUR AVERAGE			
Error includes scale factor of 2.3.			
122.5 ± 7.5 ± 8.1	ABLIKIM	23BH BES3	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-}$
71.5 ± 19.0	⁶ ABLIKIM	08D BES2	$e^+ e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
118 ± 32	⁷ MO	10 RVUE	$e^+ e^- \rightarrow \text{hadrons}$
77 ± 20	⁸ PAKHLOVA	08A BELL	$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$
119 ± 16	⁹ SETH	05A RVUE	$e^+ e^- \rightarrow \text{hadrons}$
118 ± 35	¹⁰ SETH	05A RVUE	$e^+ e^- \rightarrow \text{hadrons}$
66 ± 15	BRANDELIK	78C DASP	$e^+ e^-$
33 ± 10	SIEGRIST	76 MRK1	$e^+ e^-$
⁶ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.			
⁷ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
⁸ Systematic uncertainties not estimated.			
⁹ From a fit to Crystal Ball (OSTERHELD 86) data.			
¹⁰ From a fit to BES (BAI 02C) data.			

$\psi(4415)$ DECAY MODES

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 D\bar{D}$	seen	
$\Gamma_2 D^0\bar{D}^0$	seen	
$\Gamma_3 D^+D^-$	seen	
$\Gamma_4 D^*\bar{D} + \text{c.c.}$	seen	
$\Gamma_5 D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen	
$\Gamma_6 D^*(2010)^+D^- + \text{c.c.}$	seen	
$\Gamma_7 D^*\bar{D}^*$	seen	
$\Gamma_8 D^*(2007)^0\bar{D}^*(2007)^0 + \text{c.c.}$	seen	
$\Gamma_9 D^*(2010)^+D^*(2010)^- + \text{c.c.}$	seen	
$\Gamma_{10} D^0 D^- \pi^+ (\text{excl. } D^*(2010)^+ D^- + \text{c.c.})$	< 2.3 %	90%
$\Gamma_{11} D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.}$	(10 ± 4) %	
$\Gamma_{12} D^0\bar{D}^* \pi^+ + \text{c.c.}$	< 19 %	90%
$\Gamma_{13} D_1(2420)\bar{D} + \text{c.c.}$	possibly seen	
$\Gamma_{14} D_s^+ D_s^-$	not seen	
$\Gamma_{15} \omega\chi_{c2}$	possibly seen	
$\Gamma_{16} D_s^{*+} D_s^- + \text{c.c.}$	seen	
$\Gamma_{17} D_s^{*+} D_s^{*-}$	seen	
$\Gamma_{18} \psi_2(3823)\pi^+\pi^-$	possibly seen	
$\Gamma_{19} \psi(3770)\pi^+\pi^-$	possibly seen	
$\Gamma_{20} J/\psi\eta$	< 6 × 10 ⁻³	90%
$\Gamma_{21} \chi_{c1}\gamma$	< 8 × 10 ⁻⁴	90%
$\Gamma_{22} \chi_{c2}\gamma$	< 4 × 10 ⁻³	90%
$\Gamma_{23} \Lambda\bar{\Lambda}$	< 3.1 × 10 ⁻⁶	90%
$\Gamma_{24} \Xi^-\bar{\Xi}^+$	< 4 × 10 ⁻⁵	90%
$\Gamma_{25} pK^-\bar{\Lambda} + \text{c.c.}$	< 6 × 10 ⁻⁶	90%
$\Gamma_{26} \omega\pi^0$	not seen	
$\Gamma_{27} \omega\eta$	not seen	
$\Gamma_{28} e^+e^-$	(5.3 ± 1.2) × 10 ⁻⁶	
$\Gamma_{29} \mu^+\mu^-$	(1.1 ± 0.5) × 10 ⁻⁵	

$\psi(4415)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$		Γ_{28}
VALUE (keV)	DOCUMENT ID	TECN COMMENT
0.58±0.07 OUR ESTIMATE		
0.35±0.12	¹¹ ABLIKIM	08D BES2 $e^+e^- \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.4 to 0.8	¹² MO	10	RVUE	$e^+ e^- \rightarrow$	hadrons
0.72 ± 0.11	¹³ SETH	05A	RVUE	$e^+ e^- \rightarrow$	hadrons
0.64 ± 0.23	¹⁴ SETH	05A	RVUE	$e^+ e^- \rightarrow$	hadrons
0.49 ± 0.13	BRANDELIK	78C	DASP	$e^+ e^-$	
0.44 ± 0.14	SIEGRIST	76	MRK1	$e^+ e^-$	

¹¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.

¹² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different $e^+ e^-$ partial widths. We quote only the range of values.

¹³ From a fit to Crystal Ball (OSTERHELD 86) data.

¹⁴ From a fit to BES (BAI 02c) data.

$\Gamma(\mu^+ \mu^-)$	Γ_{29}			
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_{29}
$1.25 \pm 0.28 \pm 0.35$	^{15,16} ABLIKIM	20AG BES3	$e^+ e^- \rightarrow \mu^+ \mu^-$	
¹⁵ From a fit to the $e^+ e^- \rightarrow \mu^+ \mu^-$ cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming $\Gamma(\mu^+ \mu^-) = \Gamma(e^+ e^-)$.				
¹⁶ From solution 1 of 8 with equal fit quality. Other solutions range from $1.24 \pm 0.28 \pm 0.35$ to $1.27 \pm 0.41 \pm 0.36$ keV.				

$\psi(4415) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(J/\psi \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{20}\Gamma_{28}/\Gamma$				
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{20}\Gamma_{28}/\Gamma$
<3.6	90	WANG	13B	BELL	$e^+ e^- \rightarrow J/\psi \eta \gamma$

$\Gamma(\chi_{c1} \gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{21}\Gamma_{28}/\Gamma$				
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{21}\Gamma_{28}/\Gamma$
<0.47	90	¹⁷ HAN	15	BELL	$10.58 e^+ e^- \rightarrow \chi_{c1} \gamma$

¹⁷ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\chi_{c2} \gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{22}\Gamma_{28}/\Gamma$				
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{22}\Gamma_{28}/\Gamma$
<2.3	90	¹⁸ HAN	15	BELL	$10.58 e^+ e^- \rightarrow \chi_{c2} \gamma$

¹⁸ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{23}\Gamma_{28}/\Gamma$				
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{23}\Gamma_{28}/\Gamma$
$<1.8 \times 10^{-3}$	90	¹⁹ ABLIKIM	21AS BES3	$e^+ e^- \rightarrow \psi(4415)$	

¹⁹ From a measurement of the $e^+ e^- \rightarrow \Lambda\bar{\Lambda}$ cross section between 3.5 and 4.6 GeV.

$\Gamma(\Xi^-\bar{\Xi}^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{24}\Gamma_{28}/\Gamma$			
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0217	90	20 ABLIKIM	23BK BES3	$e^+e^- \rightarrow \psi(4415)$

20 From a fit to $e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$ cross sections.

$\Gamma(pK^-\bar{\Lambda}+\text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{25}\Gamma_{28}/\Gamma$			
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.4 × 10 ⁻³	90	21 ABLIKIM	23BL BES3	$e^+e^- \rightarrow \psi(4415)$

21 From a fit to $e^+e^- \rightarrow pK^-\bar{\Lambda}+\text{c.c.}$ cross sections.

$\psi(4415) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma^2(\text{total})$

$\Gamma(D^0D^{*-}\pi^++\text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{12}/\Gamma \times \Gamma_{28}/\Gamma$			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.99 × 10 ⁻⁶	90	22 PAKHLOVA	09	BELL $e^+e^- \rightarrow D^0D^{*-}\pi^+$

22 Using 4421 ± 4 MeV for the mass of $\psi(4415)$.

$\psi(4415)$ BRANCHING RATIOS

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$	Γ_2/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	PAKHLOVA	08	BELL $e^+e^- \rightarrow D^0\bar{D}^0\gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
not seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$

$\Gamma(D^+D^-)/\Gamma_{\text{total}}$	Γ_3/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	PAKHLOVA	08	BELL $e^+e^- \rightarrow D^+D^-\gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
not seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^+D^-\gamma$

$\Gamma(D^*(2007)^0\bar{D}^0+\text{c.c.})/\Gamma_{\text{total}}$	Γ_5/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$

$\Gamma(D^*(2010)^+D^-+\text{c.c.})/\Gamma_{\text{total}}$	Γ_6/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	ZHUKOVA	18	BELL $e^+e^- \rightarrow D^{*+}D^-\gamma$
seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^{*+}D^-\gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
seen	PAKHLOVA	07	BELL $e^+e^- \rightarrow D^{*+}D^-\gamma$

23 Supersedes PAKHLOVA 07.

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$	Γ_1/Γ_7		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.14 ± 0.12 ± 0.03	AUBERT	09M BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

$\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$ Γ_4/Γ_7

VALUE	DOCUMENT ID	TECN	COMMENT
0.17±0.25±0.03	AUBERT	09M BABR	$e^+ e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

 $\Gamma(D^*(2007)^0\bar{D}^*(2007)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^{*0}\bar{D}^{*0}\gamma$

 $\Gamma(D^*(2010)^+ D^*(2010)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	24 ZHUKOVA	18 BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	PAKHLOVA	07 BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$

24 Supersedes PAKHLOVA 07.

 $\Gamma(D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
10.5±2.4±3.8	25 PAKHLOVA	08A BELL	$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

25 Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of $\psi(4415)$. $\Gamma(D^0 D^- \pi^+ (\text{excl. } D^*(2010)^+ D^- + \text{c.c.}))/\Gamma(D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.})$ Γ_{10}/Γ_{11}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.22	90	26 PAKHLOVA	08A BELL	$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

26 Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of $\psi(4415)$. $\Gamma(D_1(2420)\bar{D} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	27 ABLIKIM	19AR BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$

27 Evidence for $e^+ e^- \rightarrow D_1(2420)\bar{D} + \text{c.c.}$ between $\sqrt{s} = 4.3$ and 4.6 GeV, not necessarily resonant. $\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA	11 BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$

 $\Gamma(\omega\chi_{c2})/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	ABLIKIM	16A BES3	$e^+ e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$

 $\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA	11 BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$

$\Gamma(D_s^{*+} D_s^{*-})/\Gamma_{\text{total}}$				Γ_{17}/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
seen	ABLIKIM	23BH BES3	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	PAKHLOVA	11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$
not seen	DEL-AMO-SA..10N	BABR		$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$

$\Gamma(\psi_2(3823)\pi^+\pi^-)/\Gamma_{\text{total}}$				Γ_{18}/Γ
<u>VALUE</u>	<u>EVTs</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
possibly seen	19	28 ABLIKIM	15S BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
28 From a fit of $e^+ e^- \rightarrow \pi^+ \pi^- \psi_2(3823)$, $\psi_2(3823) \rightarrow \chi_{c1} \gamma$ cross sections taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to the $\psi(4415)$ line shape.				

$\Gamma(\psi(3770)\pi^+\pi^-)/\Gamma_{\text{total}}$				Γ_{19}/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
possibly seen	29 ABLIKIM	19AR BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$	
29 Observe $e^+ e^- \rightarrow \pi^+ \pi^- \psi(3770)$ at $\sqrt{s} = 4.26, 4.36$, and 4.42 GeV but cannot establish if continuum or resonant.				

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$				Γ_{26}/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
not seen	ABLIKIM	22K BES3	$e^+ e^- \rightarrow \omega\pi^0$	
$\Gamma(\omega\eta)/\Gamma_{\text{total}}$				Γ_{27}/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
not seen	ABLIKIM	22K BES3	$e^+ e^- \rightarrow \omega\eta$	

$\psi(4415)$ REFERENCES

ABLIKIM	23BH PRL 131 151903	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BK JHEP 2311 228	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BL JHEP 2312 027	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22K JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AS PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20AG PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AR PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ZHUKOVA	18 PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	16A PR D93 011102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15S PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
HAN	15 PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
WANG	13B PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	11 PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SA...10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO	10 PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT	09M PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
PAKHLOVA	09 PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08D PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA	08 PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	07 PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
SETH	05A PR D72 017501	K.K. Seth	
BAI	02C PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00 PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86 SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)
SIEGRIST	76 PRL 36 700	J.L. Siegrist <i>et al.</i>	(LBL, SLAC)